Molecular Cloning, Genomic Organization, Developmental Regulation, and a Knock-out Mutant of a Novel Leu-rich Repeats-containing G Protein-coupled Receptor (DLGR-2) from *Drosophila melanogaster*

Kathrine Krageskov Eriksen,¹ Frank Hauser,^{1,3} Morten Schiøtt,²
Karen-Marie Pedersen,² Leif Søndergaard,² and Cornelis J.P. Grimmelikhuijzen^{1,4}

Departments of ¹Cell Biology, Zoological Institute and ²Genetics, Institute of Molecular Biology, University of Copenhagen, DK-2100 Copenhagen Ø, Denmark

After screening the Berkeley *Drosophila* Genome Project database with sequences from a recently characterized Leu-rich repeats-containing G protein-coupled receptor (LGR) from *Drosophila* (DLGR-1), we identified a second gene for a different LGR (DLGR-2) and cloned its cDNA. DLGR-2 is 1360 amino acid residues long and shows a striking structural homology with members of the glycoprotein hormone [thyroid-stimulating hormone (TSH); follicle-stimulating hormone (FSH); luteinizing hormone/choriogonadotropin (LH/CG)] receptor family from mammals and with two additional, recently identified mammalian orphan LGRs (LGR-4 and LGR-5). This homology includes the seven transmembrane region (e.g., 49% amino acid identity with the human TSH receptor) and the very large extracellular amino terminus. This amino terminus contains 18 Leu-rich repeats—in contrast with the 3 mammalian glycoprotein hormone receptors and DLGR-1 that contain 9 Leu-rich repeats, but resembling the mammalian LGR-4 and LGR-5 that each have 17 Leu-rich repeats in their amino termini. The DLGR-2 gene is >18.6 kb pairs long and contains 15 exons and 14 introns. Four intron positions coincide with the intron positions of the three mammalian glycoprotein hormone receptors and have the same intron phasing, showing that DLGR-2 is evolutionarily related to these mammalian receptors. The DLGR-2 gene is located at position 34E-F on the left arm of the second chromosome and is expressed in embryos and pupae but not in larvae and adult flies. Homozygous knock-out mutants, where the DLGR-2 gene is interrupted by a P element insertion, die around the time of hatching. This finding, together with the expression data, strongly suggests that DLGR-2 is exclusively involved in development.

[The nucleotide sequence(s) reported in this paper has been submitted to the GenBank/EMBL database with accession no. AF142343.]

Insects, which comprise 75% of all animal species, are of extreme ecological and economical importance, because 70% of all flowering plants depend on insects for their pollination and because insects can be severe pests, destroying about 30% of our potential annual harvest and transmitting major diseases such as malaria.

Despite the importance of insects, however, the molecular mechanisms controlling their reproduction are largely unknown. This is in contrast to the situation in mammals, where reproduction is known to be controlled by the hypothalamic neuropeptide GnRH (gonadotropin-releasing hormone), the pituitary gly-

coprotein hormones LH (luteinizing hormone) and FSH (follicle-stimulating hormone), and the gonadal steroid hormones.

Recently, using homology screening, we found that *Drosophila* produces a receptor that was structurally and evolutionarily related to the LH/FSH receptors from mammals (Hauser et al. 1997). This was an exciting finding, because it opened the possibility that insects use the same hormonal mechanisms as mammals to steer their reproduction. Subsequently, we cloned a receptor from *Drosophila* that was structurally and evolutionarily related to the GnRH receptors from mammals (Hauser et al. 1998), making the finding of a glycoprotein hormone receptor in *Drosophila* even more interesting.

Mammals have at least four glycoprotein hormones [LH, FSH, choriogonadotropin (CG), and thy-

³Present address: H. Lundbeck A/S, Ottiliavej 9, DK-2500 Valby, Denmark.

⁴Corresponding author.

E-MAIL cgrimmelikhuijzen@zi.ku.dk; FAX 0045 35 32 12 00.

roid-stimulating hormone (TSH)] and at least three glycoprotein hormone receptors (the LH/CG, FSH, and TSH receptors) that are all closely related, forming a subfamily of the large family of G protein-coupled (seven transmembrane) receptors. A characteristic of these glycoprotein hormone receptors is the presence of a very large, extracellular amino terminus that constitutes about half of the receptor protein and that contains 9 Leu-rich repeats, each measuring about 24 amino acid residues. These nine Leu-rich repeats probably form a horseshoe-like structure to which the glycoprotein hormone ligand binds (Jiang et al. 1995; Kajava et al. 1995). Similar Leu-rich repeats have been found in the *Drosophila* receptor (which is called *Drosophila* Leu-rich repeats-containing G protein-coupled receptor, or DLGR-1) (Hauser et al. 1997).

In addition to the three mammalian glycoprotein hormone receptors and DLGR-1, other LGRs have been cloned from sea anemones (Nothacker and Grimmelikhuijzen 1993), nematodes, and snails (Tensen et al. 1994). Based on the conserved sequences between the sea anemone and the *Drosophila* LGRs, two other LGRs (LGR-4 and LGR-5) were recently cloned from rat and man (Hsu et al. 1998). Similar receptors have been found by other groups (McDonald et al. 1998; Hermey et al. 1999). All of these novel invertebrate and mammalian LGRs are orphan receptors, that is, receptors for which the ligands are unknown.

In this paper we describe a second *Drosophila* receptor (DLGR-2) that is closely related to the three mammalian glycoprotein hormone receptors. Instead of 9 Leu-rich repeats, however, its amino terminus contains 18 Leu-rich sequences. We also describe the developmental expression of the DLGR-2 gene and the

isolation of a knock-out mutant. These data suggest that DLGR-2 is exclusively involved in development.

RESULTS

Identification of the *Drosophila* Receptor Protein

In an attempt to find a second *Drosophila* LGR, we screened (Spring 1997) the database of the Berkeley *Drosophila* Genome Project with the sequences of various transmembrane helices of the first *Drosophila* LGR, using the tblastn program provided by the National Center for Biotechnology Information (NCBI). This resulted in the identification of two overlapping genomic clones, P1 DS00180 (GenBank accession no. AC001660) and P1 DS01514 (GenBank accession no. AC002515), containing open reading frames resembling the genes of the three mammalian glycoprotein hormone receptors (McFarland et al. 1989; Sprengel et al. 1990; Heckert et al. 1992; Koo et al. 1991; Tsai-Morris et al. 1991). Using GENESCAN software, we predicted the exons of this presumed *Drosophila* receptor gene and designed oligonucleotide probes that could be used in PCR to amplify the *Drosophila* receptor cDNA. After PCR and 5'- and 3'-RACE, we obtained the composite cDNA coding for the *Drosophila* receptor, which we named DLGR-2 (Fig. 1A,B). Subsequently, the presence of the actual full-length transcript was confirmed with several long-range PCR experiments (Fig. 1A).

The receptor cDNA is 5399 bp long and is shown in Figure 2. Its transcription start site was determined by 5'-RACE to be at nucleotide position -628 relative to the start codon, although some 5'-RACE products indicated an additional transcription start site at position -624. Similarly, the 3'-RACE experiments indi-

Figure 1 Schematic representation of the DLGR-2 cDNA and genomic clones and the organization of the DLGR-2 gene. (A) Positions of the PCR clones. (B) Schematic drawing of the composite cDNA (top) and the organization of the receptor gene (bottom). The exons are given as bars and numbered 1–15. We named the introns after the preceding exons (e.g., intron 1 follows exon 1). The narrow and broad bars represent noncoding and coding regions, respectively. The DNA region coding for the transmembrane domain is black and that coding for the Leu-rich repeats are gray. (C) The positions of the genomic P1 clones, DS00180 and DS01514, from the Berkeley *Drosophila* Genome Project.

		CTTTAGTTCGCTTTTGTGCGTTTCGACGCGTGGC		-596
GTTCGCTCTGACGATAAAAACAAAAACGTAGAACATATATGTCCAGTTGGCGCGCTGATTTGG3CAGTAAACGGATTAGTAAAGTGCACCGAGATTTCGAGCGCGTGAACCTGGTT				-477
ACGCATATATTATATTTCTGCGCGCTGACCGCGATTTTAAACGGAATCCACCGTGGCAGCAACTTTCGCCCAAGGAGTCGGCAGTACCGGCTGTGCTCTCTCTCTGTTGGAGCGGGTTT				-358
1↓				
TGGCCGCTGTTTGTGACAGCATCAATCAACCCGATAAACCGAGAAATTCGCTGCATTTTGAGCCACATTTGAGCGTATAAAACGCTAGCAATCACAGTGAATGCCATTTTCCGCTCGTCAATA				-239
TATATGTAAAGAGTGCAAAAAACAAAAACCAATAAAGTGCAGCTGAGGCGGAAAGTGAATGCAAACTGAATGAAATTAATTCGAACTGCGGCGAGAGAGTGAAGTGAAGCTGAGTTT				-120
AAGGAACCGCAACGCGAGCGAGAAATCATGAGCGAGAAACGAATCGCACACAAAACCTGTCTCAATGACAAATGCTTTGACGACATCAATCAACAAAAGCGACGACGACATCGAGC				-1
ATG GCA GCA CGT TGC CGG TGG AGC TGG CGC CTG GCG CTC TGC CCT CTG CTG TTG CAA TTG CTT CTG CAA CTG CTC CTG TTG CCG CCA TCG				90
Met Ala Ala Arg Cys Arg Trp Ser Trp Arg Leu Ala Leu Cys Pro Leu Leu Leu Gln Leu Leu Leu Gln Leu Leu Leu Pro Pro Ser				30
GCA ATG GGC CAT GAT GAA ACC AAA GAA AAT CCC GCT CCG GAT ATG CAA AAT AGC CAG GAG CAA GAG CCA TAT GTC CAC CTG CAA CAT CTG				180
Ala Met Gly His Asp Glu Thr Lys Glu Asn Pro Ala Pro Asp Met Gln Asn Ser Gln Glu Gln Glu Pro Tyr Val His Leu Gln His Leu				60
CAG CAG CAG CAG CAG CAA AAT CCA CAA ACC GTC CAA CAA CTT AGC CAA ATT ACC GTG AAT AGA ACC TCG AAA TCA GCC AGT GTC ACT CCT				270
Gln Gln Gln Gln Gln Gln Asn Pro Gln Thr Val Gln Gln Leu Ser Gln Ile Thr Val Asn Arg Thr Ser Lys Ser Ala Ser Val Thr Pro				90
ACG GGC ATC AGG GAG AAC GTG ATG CTG CCA TCG GCC GAT CCC GAG AAG GAG GCC CAG ATC CTG TAC GAG AAG TCC TTG CAG GAA TAC CAT				360
Thr Gly Ile Arg Glu Asn Val Met Leu Pro Ser Ala Asp Pro Gln Glu Lys Glu Ala Gln Ile Leu Tyr Glu Lys Ser Gln Gln Glu Tyr His				120
GGC AGT CAG CTC TCC ACC CCC TCA ACC GCC ACT GAT GTC ATC GCA GGC AAG CAG CCG CTG CAC TCC ATT TGT Ser CCG TGG TTG CAG AAG				450
Gly Ser Gln Leu Ser Thr Ala Ser Thr Thr Asp Val Ile Ala Gly Lys Arg CAG ACC CTG CAC Ser Ile Cys Glu Arg Trp Leu Gln Lys				150
CAC TGT CAC TGC ACC GGC AGC CTG GAG GTC CTG CCG CTC AGT TGT CGA GGC ATA GGG ATC CTG GCC GTG CCC GTC AAT CTT CCC AAC GAG				540
His Cys His Cys Thr Gly Ser Leu Glu Val Leu Arg Leu Ser Cys Arg Gly Ile Gly Ile Leu Ala Val Pro Val Asn Leu Pro Asn Glu				180
2↓				
GTG GTC GTC CTA GAT TTG GGT AAC AAC AAC TTA ACC AAA TTG GAA GCG AAC TCA TTT TTT ATG GCA CCC AAT CTG GAG GAT CTA ACT TTG				630
Val Val Val Leu Asp Leu Gly Asn Asn Asn Leu Thr Lys Leu Ala Asn Ser Phe Phe Met Ala Pro Asn Leu Glu Asp Leu Thr Leu				210
TCC GAC AAT AGC ATC ATT AAT ATG GAT CCC AAT GCA TTC TAT GGC CTG GCC AAA TTG AAA CGA TTG AGC CTG CAG AAC TGT GGC CTC AAG				720
Ser Asp Asn Ser Ile Ile Asn Met Asp Pro Asn Ala Phe Tyr Gly Leu Ala Lys Leu Lys Arg Leu Ser Leu Gln Asn Cys Gly Leu Lys				240
TCC TTG CCA CCG CAG TCG TTC CAA GGA CTC GCT CAG CTG ACC AGC ATA CTA CAG CTG AAT GGC AAC GCC CTG GTC AGT CTG GAT GGC GAT TGT				810
Ser Leu Pro Pro Gln Ser Phe Gln Gly Leu Ala Gln Leu Thr Ser Leu Gln Leu Asn Gly Asn Ala Leu Val Ser Leu Asp Gly asp Cys				270
CTG GGC CAC CTG CAG AAA CTG CGC ACC TTG CGA TTG GAG GGC AAT CTC TTC TAT CGC ATT CCC ACG AAT GCC TTG GCC GGA CTC AGA ACC				900
Leu Gly His Leu Gln Lys Leu Arg Thr Leu Arg Leu Gly Asn Leu Phe Tyr Arg Ile Pro Thr Asn Ala Leu Ala Gly Leu Arg Thr				300
5↓				
CTA GAA GCA CTC AAT TTG GGC AAT TTG TCG ACA ATA AAT GAC GAG CAG TTT CCG CGA ATG CCA AAC TTG ATC CTG CTT TTG CTG				990
Leu Glu Ala Leu Asn Leu Gly Ser Asn Leu Leu Thr Ile Ile Asn Asp Glu Asp Phe Pro Arg Met Pro Asn Leu Ile Val Leu Leu Leu				330
AAG CGA AAT CAA ATC ATG AAA ATC TCC GCG GGA GCT CTT AAA AAT TTA ACA GCC TTA AAA GTT TTA GAG CTG GAC GAT AAT TTA ATA AGC				1080
Lys Arg Asn Gln Ile Met Lys Ile Ser Ala Gly Ala Leu Lys Asn Leu Thr Ala Leu Lys Val Leu Glu Leu Asp Asp Asn Leu Ile Ser				360
8↓				
AGC CTA CCC GAG GGA CTC AGC AAA TTG TCG CAA CTG CAG GAG CTT TCC ATC ACC AGC AAT CGA CTG CGC TGG ATA AAC GAC ACG GAA CTG				1170
Ser Leu Pro Glu Gly Leu Ser Lys Leu Ser Gln Leu Gln Glu Leu Ser Ile Thr Ser Asn Arg Leu Arg Trp Ile Asn Asp Thr Glu Leu				390
CCC AGG AGC ATG CAA ATG CTG GAC ATG AGG GCC AAT CCT CTG TCC ACA ATT TOG GCA GGG GCG TTT CGG GGC ATG TCC AAG TTG CCG AAG				1260
Pro Arg Ser Met Gln Met Leu Asp Met Arg Ala Asn Pro Leu Ser Thr Thr Ile Ser Ala Gly Ala Phe Arg Gly Leu Ser Lys Leu Arg Lys				420
9↓				
CTG ATT TTA TCG GAT GTG CGA ACT TTG CGC TCG TTT CCC GAA CTT GAG GCC TGC CAT GCG CTG GAA ATA CTG AAA CTG GAT CGT GCC GGC				1350
Leu Ile Leu Ser Asp Val Arg Thr Leu Arg Ser Phe Pro Glu Leu Glu Ala Cys His Ala Leu Glu Ile Leu Lys Leu Asp Arg Ala Gly				450
10↓				
ATT CAG GAG GTG CCA GCG AAT CTA TGC CGC CAA ACG CCA AGA CTG AAG AGT TTA GAA CTA AAA ACC AAT TCC CTA AAA CGC ATC CCA AAT				1440
Ile Gln Glu Val Pro Ala Asn Leu Cys Arg Gln Thr Pro Arg Leu Lys Ser Leu Glu Leu Lys Thr Asn Ser Leu Lys Arg Ile Pro Asn				480
11↓				
TTA AGT AGC TGC AGA GAC TTA AGG CTG TTA GAT CTA TCG AGC AAT CAA ATT GAA AAG ATT CAG GGA AAG CCC TTC AAT GGA TTG AAG CAA				1530
Leu Ser Ser Cys Arg Asp Leu Arg Leu Leu Asp Leu Ser Ser Asn Gln Ile Glu Lys Ile Gln Gly Lys Pro Phe Asn Gln Lys Lys Gln				510
12↓				
TTG AAC GAT CTG TTG CTC TCC TAC AAT CGA ATC AAG GCA CTG CCA CAA GAT GCC TTT CAG GGC ATT CCG AAG CTC CAA CTT CTT GAC CTG				1620
Leu Asn Asp Leu Leu Leu Ser Tyr Asn Arg Ile Lys Ala Leu Pro Gln Asp Ala Phe Gln Gly Ile Pro Lys Leu Gln Leu Leu Asp Leu				540
GAG GGC AAC GAG ATC TCG TAC ATA CAC AAG GAA GCC TTT TCC GGT TTC ACA GCC CTC GAG GAT CTC AAT TTG GGC AAC AAT ATT TTT CCA				1710
Glu Gly Asn Glu Ile Ser Tyr Ile His Lys Glu Ala Phe Ser Gly Phe Thr Ala Leu Glu Asp Leu Asn Leu Gly Asn Asn Ile Phe Pro				570
GAG TTG CCC GAA TCG GGA CTG CGC GCT TTG CTA CAC CTA AAG ACC TTC AAC AAT CCA AAA CTT AGG GAG TTC CCG CCA CCG GAC ACC TTT				1800
Glu Leu Pro Glu Ser Gly Leu Arg Ala Leu Leu His Leu Lys Thr Phe Asn Asn Pro Lys Leu Arg Glu Phe Pro Pro Ser Pro Asp Thr Phe				600
CCC AGG ATC CAA ACT CTT ATA CTC TCT TAT GGC TAC CAC TGC TGC GCG TTT CTC CCA CTG GTT GCC ATG TCG TCC CAG AAG AAA ACC TCC				1890
Pro Arg Ile Gln Thr Leu Ile Leu Ser Tyr Ala Tyr His Cys Cys Ala Phe Leu Pro Leu Val Ala Met Ser Ser Gln Lys Lys Thr Ser				630
CAG GTC CAG GAG GCT GTA CTC TTT CCC TCC GAT GCC GAG TTC GAT ATG ACT TTG TGG AAC AAT AGC ATG ATG AAT ATT TGG CCG CAA ATG				1980
Gln Val Gln Glu Ala Val Leu Phe Pro Ser Asp Ala Glu Phe Asp Met Thr Leu Trp Asn Asn Ser Met Met Asn Ile Trp Pro Gln Met				660
13↓				
CAT AAT CTC AGC AAG CAG TTG GGT GCT TCC ATG CAT GAT CCT TGG GAG AOG GCT ATT AAT TTC AAC GAA GAG CAG CTG CAA ACA CAA ACG				2070
His Asn Asn Lys Lys Gln Leu Gly Ala Ser Met His Asp Pro Trp Glu Thr Ala Ile Asn Phe Asn Glu Glu Gln Leu Thr Gln Thr				690
GGA GGC CAA ATC GCC ACC AGC TAC ATG GAG GAG TAC TTC GAG GAG CAC GAT GTG AGT GGT CCT GCC ACG GGA TAC GGC TTT GGT ACT GGA				2160
Gly Gly Gln Ile Ala Thr Ser Tyr Met Glu Glu Tyr Phe Glu Glu His Asp Val Ser Gly Pro Ala Thr Gly Tyr Gly Phe Gly Thr Gly				720
CTA TTC TCT GGT ATG TCC ACA GAG GAC TTT CAA CCG GGT TCG GTG CAG TGC CTG CCA ATG CCA GGA CCT TTT CTA CCC TGC GCC GAT CTC				2250
Leu Phe Ser Gly Met Ser Thr Glu Asp Phe Gln Pro Gly Ser Val Gln Cys Leu Pro Met Pro Gly Pro Phe Leu Pro Cys Ala Asp Leu				750

Figure 2 cDNA and deduced amino acid sequence of DLGR-2. This figure is compiled from the sequences of the overlapping cDNA clones 5'C-2, 5B, and 3'4-J (Fig. 1A). Nucleotides are numbered from 5' to 3' end, and the amino acid residues are numbered starting with the first ATG in the open reading frame. Introns are indicated by arrows and numbered 1–14. The seven membrane-spanning domains are boxed and labeled TM I–VII. The proposed signal sequence is shaded. Spades indicate potential amino-glycosylation sites. The inframe stop codon in the 5' region, upstream of the assigned start codon is underlined twice; the translation termination codon is indicated by an asterisk (*). Putative polyadenylation sites at the 3' end are underlined.

Drosophila G Protein-coupled Receptor

TM I																												2340	
TTC	GAT	TGG	TGG	ACT	CTG	CGC	TGC	GGA	GTG	TGG	GTG	GTT	TTT	CTG	TTG	TCT	CTC	CTG	GGC	AAT	GGA	ACG	GTG	GTG	TTT	GTG	CTG	CTT	TGC
Phe	Asp	Trp	Trp	Thr	Leu	Arg	Cys	Gly	Val	Trp	Val	Val	Phe	Leu	Leu	Ser	Leu	Leu	Gly	Asn	Gly	Thr	Val	Val	Phe	Val	Leu	Leu	Cys
TM II																												2430	
TCG	CGA	TCT	AAG	ATG	GAT	GTG	CCC	CGA	TTT	CTA	GTC	TGT	AAT	CTG	GCC	GCG	GCC	GAC	TTC	TTC	ATG	GGC	ATC	TAC	CTC	GGT	ATC	CTG	GCC
Ser	Arg	Ser	Lys	Met	Asp	Val	Pro	Arg	Phe	Leu	Val	Cys	Asn	Leu	Ala	Ala	Ala	Asp	Phe	Phe	Met	Gly	Ile	Tyr	Leu	Gly	Ile	Leu	Ala
TM III																												2520	
ATT	GTG	GAT	GCG	GCG	ACG	TTA	GCG	GAA	TTT	CGC	ATG	TTT	GCC	ATT	CCT	TGG	CAG	ATG	TCC	GTG	CTC	TGC	CAG	TTG	TCC	GGT	TTT	CTG	GCC
Ile	Val	Asp	Ala	Ala	Thr	Leu	Gly	Glu	Phe	Arg	Met	Phe	Ala	Ile	Pro	Trp	Gln	Met	Ser	Val	Leu	Cys	Gln	Leu	Ser	Gly	Phe	Leu	Ala
TM IV																												2610	
GTA	CTC	AGC	TCA	GAA	TTA	TCC	GTT	TAC	ACG	TTG	GCT	GTG	ATC	ACT	TTG	GAG	CGC	AAT	TAC	GCC	ATC	ACT	CAT	GCC	ATT	CAC	CTG	AAC	AAG
Val	Leu	Ser	Ser	Glu	Leu	Ser	Val	Tyr	Thr	Leu	Ala	Val	Ile	Thr	Leu	Glu	Arg	Asn	Tyr	Ala	Ile	Thr	His	Ala	Ile	His	Leu	Asn	Lys
TM V																												2700	
AGG	CTC	TCC	TTG	AAG	CAG	GCG	GGA	TAT	ATA	ATG	AGT	GTA	GGA	TGG	GTT	TTT	GCC	CTG	ATC	ATG	GCT	TIG	ATG	CCT	TTG	GTT	GGT	GTT	TCG
Arg	Leu	Ser	Leu	Lys	Gln	Ala	Gly	Tyr	Ile	Met	Ser	Val	Gly	Trp	Val	Phe	Ala	Leu	Ile	Met	Ala	Leu	Met	Pro	Leu	Val	Gly	Val	Ser
TM VI																												2790	
GAC	TAC	AGG	AAG	TTT	GCC	GTG	TGT	TTG	CCA	TTC	GAG	ACC	ACC	ACT	GGA	CCG	GCC	AGT	TTG	ACC	TAT	GTA	ATC	TCA	CTG	ATG	TTC	ATC	AAC
Asp	Tyr	Arg	Lys	Phe	Ala	Val	Cys	Leu	Pro	Phe	Glu	Thr	Thr	Thr	Gly	Pro	Ala	Ser	Leu	Thr	Tyr	Val	Ile	Ser	Leu	Met	Phe	Ile	Asn
TM VII																												2930	
GGA	TGC	GCA	TTT	CTC	ACT	CTG	ATG	GGT	TGC	TAC	CTG	AAG	ATG	TAC	TGG	GCC	ATA	AGG	GGC	AGC	CAG	GCA	TGG	AAC	ACG	AAT	GAT	TCG	CGA
Gly	Cys	Ala	Phe	Leu	Thr	Leu	Met	Gly	Cys	Tyr	Leu	Lys	Met	Tyr	Trp	Ala	Ile	Arg	Gly	Ser	Gln	Ala	Trp	Asn	Thr	Asn	Asp	Ser	Arg
TM VIII																												2880	
ATT	GCC	AAG	CGA	ATG	GCT	CTG	TTG	GTC	TTT	ACT	GAC	TTC	CTC	TGC	TGG	TCA	CCT	ATC	GCC	TTC	TCG	ATC	ACT	GCC	ATC	TTT	GGC	CTG	2970
Ile	Ala	Lys	Arg	Met	Ala	Leu	Leu	Val	Phe	Thr	Asp	Phe	Leu	Cys	Trp	Ser	Pro	Ile	Ala	Phe	Phe	Ser	Ile	Thr	Ala	Ile	Phe	Gly	Leu
TM IX																												2990	
CAA	CTG	ATT	TCG	CTG	GAG	CAG	GCC	AAG	ATC	TTT	ACG	GTA	TTT	GTG	CTG	CCC	TTG	AAT	AGT	TGC	TGC	AAT	CCC	TTT	CTC	TAC	GCC	ATA	ATG
Gln	Leu	Ile	Ser	Leu	Glu	Gln	Ala	Lys	Ile	Phe	Thr	Val	Phe	Val	Leu	Leu	Leu	Asn	Ser	Cys	Cys	Asn	Pro	Phe	Leu	Tyr	Ala	Ile	Met
TM X																												3060	
ACC	AAG	CAG	TTC	AAG	AAG	GAC	TGT	GTG	ACG	TTG	TGC	AAA	CAT	TTC	GAA	GAA	TCC	CGC	GTT	GTG	GGA	GGC	GGT	GGT	CCG	GGT	GGA	CGT	GGT
Thr	Lys	Gln	Phe	Lys	Lys	Asp	Cys	Val	Thr	Leu	Cys	Lys	His	Phe	Glu	Glu	Ser	Arg	Val	Val	Gly	Gly	Gly	Gly	Pro	Gly	Gly	Arg	Gly
TM XI																												3150	
GCA	GTG	GCA	AGA	ACC	AAG	AGG	GGT	GAT	CTC	CCG	CCA	CCA	CTC	TTA	CCC	GCT	GCA	GCG	GTG	GCA	CAT	CCG	CCA	GGC	TGT	CGA	TGT	CTA	CGG
Ala	Val	Ala	Arg	Thr	Lys	Arg	Gly	Asp	Leu	Pro	Pro	Pro	Leu	Leu	Pro	Ala	Ala	Ala	Val	Ala	His	Pro	Pro	Gly	Cys	Arg	Cys	Leu	Arg
TM XII																												3240	
ATG	CTG	CCC	AGC	GAA	ATG	CCC	AAC	TGG	CAT	AAA	ATG	GAG	CAG	ACG	CCG	AGC	ATG	TGG	CAG	AGA	TTG	AGA	ACA	TTT	TGC	TGC	GGG	GAG	AAC
Met	Leu	Pro	Ser	Glu	Met	Pro	Asn	Trp	His	Lys	Met	Glu	Gln	Thr	Pro	Ser	Met	Trp	Gln	Arg	Leu	Arg	Thr	Phe	Cys	Cys	Gly	Glu	Asn
TM XIII																												3330	
CGA	AGG	CGT	CGC	AAA	CAG	CGA	CGG	CAA	CCG	CAA	CAG	CGA	CGC	CAG	AGA	GCC	TAT	ACT	GCC	GCA	GCC	GCA	AAT	CCG	TAT	CAG	TAT	CAG	TTT
Arg	Arg	Arg	Arg	Lys	Gln	Arg	Arg	Gln	Pro	Gln	Gln	Arg	Arg	Gln	Arg	Ala	Tyr	Thr	Ala	Ala	Ala	Ala	Asn	Pro	Tyr	Gln	Tyr	Gln	Phe
TM XIV																												3420	
GCT	GAA	CTG	CGT	CAG	CAA	CGC	AAT	CGA	GCC	AGT	TCC	ATT	TCC	AGC	GAG	AAC	TTC	TGC	AGC	TCA	AGA	TCA	TCC	AGC	TGG	CGA	CAT	GGT	3510
Ala	Glu	Leu	Arg	Gln	Gln	Arg	Gln	Asn	Arg	Ala	Ser	Ser	Ile	Ser	Ser	Glu	Asn	Phe	Cys	Ser	Ser	Arg	Ser	Ser	Ser	Trp	Arg	His	Gly
TM XV																												3510	
CCA	CCA	TCC	TCA	GCA	CCC	GTG	CCG	CGG	GGT	AAC	TGC	AGC	ATG	CCG	CTG	AAG	ATG	CTG	GAG	CCA	CAT	GCT	CAT	CCG	CAC	GGT	CAT	GGG	CGC
Pro	Pro	Ser	Ser	Ala	Pro	Val	Pro	Pro	Gly	Asn	Cys	Ser	Met	Pro	Leu	Lys	Met	Leu	Glu	Pro	His	Ala	His	Pro	His	Gly	His	Gly	Arg
TM XVI																												3600	
AGA	CGA	CAC	TCC	GCC	TGG	CTG	ATC	ACG	CGG	AAA	ACG	TCG	CAG	GAC	TCG	AAC	TTG	TCC	AGC	TCG	CGC	AAT	GAC	TCC	TCC	GCC	TCG	GCG	ACC
Arg	Arg	His	Ser	Ala	Trp	Leu	Ile	Thr	Arg	Lys	Thr	Ser	Gln	Asp	Ser	Gln	Asn	Leu	Ser	Ser	Ser	Arg	Asn	Asp	Ser	Ser	Ala	Ser	Ala
TM XVII																												3690	
ACC	GCC	AGC	ACC	TCC	ACG	TTT	CGG	TTA	TCC	CGG	TCA	AGT	GCC	GGC	AGC	AGC	ACA	CCC	CTG	CCC	TCG	ATC	ATA	GCC	CAC	AAT	GGG	AAG	GCC
Thr	Ala	Ser	Thr	Ser	Thr	Phe	Arg	Leu	Ser	Arg	Ser	Ser	Ala	Gly	Ser	Ser	Thr	Pro	Leu	Pro	His	Ala	His	Pro	His	Gly	His	Gly	Arg
TM XVIII																												3780	
CAG	TTG	GAT	GCA	GTT	AAA	CCA	CGA	TTG	GTC	CGC	CAG	GAA	GCA	GTC	CAG	GAA	GAG	GAC	TCA	TCA	CCA	CCT	CGG	CTG	GGA	GTC	CGT	TTC	3870
Gln	Leu	Asp	Ala	Val	Lys	Pro	Arg	Leu	Val	Arg	Gln	Glu	Ala	Val	Gln	Glu	Glu	Gly	Asp	Ser	Ser	Pro	Pro	Arg	Leu	Gly	Val	Arg	Phe
TM XIX																												1290	
CTG	CCC	ACC	ATA	CCC	TGC	GCT	GCC	GAC	AGT	TCT	GTC	GTC	ATG	GAA	GAT	GGC	GAC	TCG	GCC	AAC	ACG	GGA	GTA	GCC	AGT	TTT	CTA	GGC	ATG
Leu	Pro	Thr	Ile	Pro	Ser	Ala	Ala	Asp	Ser	Ser	Val	Val	Met	Glu	Asp	Gly	Asp	Ser	Ala	Asn	Thr	Gly	Val	Ala	Ser	Phe	Leu	Gly	Met
TM XX																												3960	
CCA	CTT	CCT	GGG	GCG	TCC	AGC	GGC	TTT	CTT	ATA	GCT	CCA	ACC	ACC	GCT	GCC	ACC	ACC	TCC	CCA	CCC	CCG	GTG	GTC	CTG	CAG	CCG	GCA	AAG
Pro	Leu	Pro	Gly	Ala	Ser	Ser	Gly	Phe	Leu	Ile	Ala	Pro	Thr	Ala	Ala	Ala	Ala	Thr	Ser	Pro	Pro	Pro	Val	Val	Leu	Gln	Pro	Ala	Lys
TM XXI																												4050	
CCA	CCT	CCT	GAT	CCC	AAC	GAC	GCA	CCA	CTA	TGA	TGGTGGGACTGGCGCTGGCGAAGATTGGGTGGGGCCCAACAGGAGACCTTTCTATATAAAGTAATGGGGAACAGGA	4158																	
Pro	Pro	Pro	Asp	Pro	Asn	Asp	Ala	Pro	Leu	*																			
TM XXII																												1360	
CAAAAGCTGACTGTGTTTCATAGATTGTATTGTGGGGTGGGCTGGGGGAGGTGATGAGGGCTGTTCGATTGGGTTGGCACTTAGGCTAAGTAAGACACAAATTAATAAAATTCACATTAG																												4277	
GCATATACAAAAATATAATGTATTGTAGCATTAAGTAGCCCAATTGTGTTAAGTAAGAACCAACTGAAGATCCGGATCCCAATTAAATCAACAAACAAACAAAGGATAACACCGTTGAAA																												4396	
GGATCAGTACATAAATCCAGTTTCTCAAAATGATTGTACATTTCTCAGATGTGTGTCATCAACATAAAATTTGATTGATCATTTGTTAGATAAAATACTTTAAAGTGAACCTAGATCTGGTAT																												4515	
ATGAAACAACACTTTGAGAAACTGAAGCCGACAATGAATATTAAGTATTACCTAAATTTATACCGGTAGATTATGTTTTTAACTTTTAAACCATATAATGTTTAACTTAACTTATATATATATA																												4634	
TATTTATATACGTGTGTGTTTATTTTCGATGCCAAGCATCTCTCTGTGTAAACTTAACTCGAACGAGAAGATTGTAAAGAAACCAAGCCATATAAAATTTGAATTTAAATATTTCCACAG																												4753	
ACAAAAATTTTCTCGCCTT(A) ₁₆																												4772	

Figure 2 (Continued.)

cated various transcription termination points. The longest identified transcript extended 689 bp beyond the stop codon and the transcription stop at this point indicated the use of an imperfect polyadenylation signal (Proudfoot 1991), AATTAA, at nucleotide positions 4735–4740 of Figure 2, that is, 32 bp upstream of the poly(A)⁺ tail. Other 3'-RACE experiments indicated additional transcription termination points at nucleotide positions 4753 and 4623 of Figure 2 and, hence, twice the use of another imperfect polyadenylation signal, CATAAA. All putative polyadenylation signals are underlined in Figure 2.

Although the flanking regions of the ATG codon at nucleotide positions 1–3 of Figure 2 do not fully match the *Drosophila* consensus sequence for a translation start site (Cavener 1987), it is probably the start codon, because there is an in-frame stop codon shortly upstream of this codon (at position –42 to –40 of Fig. 2). Furthermore, this potential start codon is followed by a series of nucleotides that code for the signal sequence, which probably is cleaved-off between amino acid positions 33 and 34 of Figure 2 (von Heijne 1986, 1990).

The cDNA of Figure 2 codes for a protein of 1360 amino acid residues, which corresponds to a molecular

Figure 3 (See facing page for legend.)

Figure 3 Amino acid sequence comparison between DLGR-2, LGR-5, LGR-4, DLGR-1, the human TSH receptor (TSHR), and the LGR from the sea anemone *Anthopleura elegantissima* (ALGR). Broken lines represent spaces introduced to optimize alignment. The Gly-rich repeats of ALGR (Nothacker and Gimmelikhuijzen 1993) were omitted to facilitate alignment. Amino acid residues that are identical between DLGR-2 and at least one of the other receptors are boxed. Known intron–exon transitions in the genes coding for the receptors are shaded at the corresponding amino acid residues. The positions of the aliphatic and aromatic residues characteristic for the Leu-rich repeats in DLGR-2 are marked by asterisks (*) or solid circles (●). The solid circles also mark intron–exon transitions in the DLGR-2 gene that occur at the same positions and have the same intron phasing as in the DLGR-1, TSHR, and ALGR genes. The seven membrane-spanning domains are indicated by I–VII. The open circles (○) mark conserved cysteine residues, flanking the region containing the Leu-rich repeats. The amino acid residue positions are given at right. The amino acid sequences of DLGR-1 and the mammalian and sea anemone receptors as well as the intron–exon positions in their genes are from Mirahi et al. (1990); Gross et al. (1991); Nothacker and Gimmelikhuijzen (1993); Hauser et al. (1997); Hsu et al. (1998); and Vibede et al. (1998).

mass of 150.8 kD. Several potential amino glycosylation sites on the extracellular portion of the protein (seven on the amino terminus, but none on the extracellular loops; Fig. 2) suggest that the true mass might be considerably higher, although removal of the signal sequence will, of course, somewhat reduce the molecular mass of the mature protein.

The receptor protein has seven putative transmembrane domains (Fig. 2) characteristic for G protein-coupled receptors. The intracellular loops in between these transmembrane domains and the carboxy-terminal intracellular part of the protein, contain multiple serine and threonine residues, which are po-

tential phosphorylation sites (e.g., Ser-1204 is part of the consensus sequence RRHS for A-kinase phosphorylation; see Kemp and Pearson 1990).

Comparison of DLGR-2 with Other Related Receptors

Figure 3 shows a comparison of the amino acid sequence of DLGR-2 with that of DLGR-1, the sea anemone LGR, the human TSH receptor, and the two newly discovered mammalian orphan receptors, LGR-4 and LGR-5. The transmembrane region of DLGR-2 (amino acid positions 758–1021 of Fig. 2) shows a high sequence identity with the human TSH receptor (49%),

the sea anemone LGR (47%), and DLGR-1 (45%). Less sequence identity is found with LGR-4 and LGR-5 (30%–31%). When the region of the amino terminus of DLGR-2, containing the Leu-rich repeats (amino acid positions 205–617 of Fig. 2) is compared, however, most sequence identity is found with LGR-4 and LGR-5 (36%–37%), which is higher than when the transmembrane regions are compared. Less sequence identity is found between the Leu-rich repeats of DLGR-2 (amino acid positions 393–617 of Fig. 2) and the human TSH receptor (27%), DLGR-1 (25%), and the sea anemone LGR (22%). Furthermore, a long Gln-rich region of about 100 amino acid residues that is located shortly after the signal sequence is only found in DLGR-2 (Fig. 2).

The amino terminus of DLGR-2 contains a region with 18 Leu (or Ile/Val/Ala/Phe)-rich repeats (marked by solid circles and asterisks in Fig. 3). This region is flanked by clusters of cysteine residues, which is a typical combination also found in other Leu-rich repeats-containing proteins (Kobe and Deisenhofer 1994, 1995). The aliphatic and aromatic residues of the Leu-

rich repeats of DLGR-2 form a pattern that strongly resembles that of the other LGRs (Figs 3 and 4). Compared with the other LGRs, however, DLGR-2 has the largest number of these repeats, although it resembles LGR-4 and LGR-5, which have 17 repeats (Hsu et al. 1998).

DLGR-2 is considerably larger than the other known LGRs. This is not only due to the large amino terminus, but also to the intracellular carboxyl terminus that is 340 amino acid residues in length, which is 2–4 times longer than the carboxyl termini of the other receptors of Figure 3. Homology screening with the receptor carboxyl terminus did not reveal the existence of other known proteins with resembling sequences.

Genomic Organization of the *Drosophila* Receptor Gene

The genomic organization of the gene coding for DLGR-2 is summarized in Figure 1B. Figure 1C shows the position of the gene on the genomic P1 clones DS00180 and DS01514 from the Berkeley *Drosophila* Genome Project. The total length of the transcribed portion of the gene is 18.6 kb. No conventional TATA box could be recognized on the P1 clones within a region of 1 kb upstream of the identified transcription start site.

The gene contains 14 introns and 15 exons. Four introns in the gene occur at exactly the same positions and have the same intron phasing (intron phase 2) as four introns in the rat TSH, FSH, and LH/CG receptor genes; the DLGR-1 gene; and the sea anemone LGR gene (indicated by solid circles in Fig. 3), strongly suggesting that these genes are evolutionarily related. The introns have a length ranging from 55 bp to more than 7 kb (Table 1), and most of them are located in the region coding for the receptor amino terminus, in such a way that each of the resulting exons codes for one or more Leu-rich repeats. There are no introns in the gene region coding for the seven-transmembrane domain, but there is one intron in the gene region coding for the long intracellular carboxyl terminus (Fig. 1B).

Figure 4 Leu (Ile/Val/Ala/Phe)-rich repeats in the amino-terminal region of DLGR-2 and the rat LH receptor. (A) Consecutive segments (L1–L18) within the amino terminus of DLGR-2 were aligned and small gaps (–) were introduced to show that many of the aliphatic and aromatic residues in one segment occur at a similar position in the other segments. These residues are boxed. In addition, Asn residues at positions typical for Leu-rich repeats (Kobe and Deisenhofer 1994, 1995) are also marked. The shaded aliphatic residues correspond to intron–exon transitions in the receptor gene and are given at the start of the repeating segments. Most of the repeating segments, therefore, are coded for by distinct exons. Only complete segments, lying within the cluster of Cys residues bordering the Leu-rich repeats of Leu-rich repeats-containing proteins (Kobe and Deisenhofer 1994, 1995), are taken into account. However, 0.8 repeat flanking L1 and 0.3 repeat flanking L18 might contribute to an additional repeat (altogether 19 Leu-rich repeats) in DLGR-2. (B) Similar alignment of the rat LH receptor. Data from Bhowmick et al. (1996).

Table 1. Intron/Exon Boundaries of the DGLR-2 Gene

Intron	5' Donor	Intron size (bp)	3' Acceptor	Intron phase
1 (variant1)	TTT gtaaggt....	~7 kb	cccttag ACA	–
1 (variant2)	AAG gtgccac....	~7 kb	cccttag ACA	–
1 (variant3)	TTG gtgccac....	~7 kb	cccttag ACA	–
1 (variant4)	TTT gtaaggt....	~7 kb	taaataag ATA	–
2	CT gtgagta.... Leu	1223 bp	ccaacag A Leu	2
3	CT gtgagta.... Leu	583 bp	ttgcag A Leu	2
4	CT gtaagta.... Leu	325 bp	cacacag A Leu	2
5	CT gtgagcc.... Leu	591 bp	tcgacag C Leu	2
6	CT gtaagta.... Leu	74 bp	ttgcag T Leu	2
7	TT gtaagta.... Leu	1747 bp	ttgcag A Leu	2
8	CT gtaagta.... Leu	1083 bp	ccttcag T Leu	2
9	CT gtgagta.... Leu	55 bp	ttttcag G Leu	2
10	TT gtgagtt.... Leu	65 bp	actgcag A Leu	2
11	TT gtaagtt.... Leu	57 bp	caccag A Leu	2
12	CT gtgcgt.... Leu	61 bp	ttcacag T Leu	2
13	C gtaagt.... His	56 bp	tccccag AT His	1
14	G gtaagt.... Ala	57 bp	tgtacag CG Ala	1

The sequence of each of the intron-exon boundaries is shown, as well as the codons for the amino acid residues. Uppercase and lowercase letters represent nucleotides in the exons and introns, respectively. For intron 1, the four splicing variants that we have isolated are given. These variants differ from each other, depending on which 5' donor and 3' acceptor sites are used. The sequences of the introns are given in our GenBank/EMBL Database submission. The overall positions of the introns are shown in Figs. 1 and 2. Introns 9-12 occur at the same positions within the gene and have the same intron phasing as four introns in the genes of the mammalian glycoprotein hormone receptors (● in Fig. 3).

Alternative Splicing

During the analysis of several clones, containing the 5' noncoding region of the receptor cDNA, four splicing variants were identified that all concerned exons 1 and 2 and intron 1. Three donor and two acceptor sites were

found to exist (Fig. 5), which means that there are six possible transcripts in which the contributions from exon 1 can be either 45, 50, or 217 nucleotides and from exon 2 either 894 or 1037 nucleotides. Four of these transcripts have been identified (see legend to Fig. 5).

```

                                d1    d2
AGTTGGGCGGCTGATTGGGCAGTAAACGGATTAgtaagtgccaccgagatttGTAGGCGCGTGAACTCGTTACGCATAT -469
ATTATATTTCTGGGCGCTGACCGCGATTTTAAACGGAATCCAACGGTGGCAGCAACTTGCAGCAAGGAGTCGGCACTACCG -389
CCTGTGCTCTCTTCTGTGGAGCGGGTTTGTGGCCGCTGTTTGTgtgtgtgtttattggcgttaaagggtcgttactt
                                d3
..... INTRON 1 .....
                                a1
caaaatatagctacttttttaaaattttgcatacaatttaatttatattgccataaatagaataataatcattaaatttgt
aatagcataatgtgctactgtgggataatggcatgcattagccgcaaatcatcaagagcccatataacaaaaattataata
aattacgaaccctgactgacacctcattatcgatccccttagACAGCATCAATCACCCGATAAACCGAGAATTCGCTG -307
.... EXON 2 .....
                                a2

```

Figure 5 Partial nucleotide sequence of the genomic DNA around intron 1. The numbers in this figure refer to the nucleotide positions of the cDNA of Fig. 2. Uppercase and lowercase letters represent the nucleotides in exons 1 and 2 and in intron 1, respectively. The three gt splicing donor sites and the two ag acceptor sites are underlined and printed in boldface type. The three donor and two acceptor sites give six possible mRNAs, of which four have been identified. The four identified mRNAs have the combinations d1/a1 (donor 1/acceptor 1), d1/a2, d2/a2, and d3/a2. Fig. 2 corresponds to d3/a2.

Sequence Deviations Between the Cloned Receptor cDNA and the Genomic Database DNA

The coding region of our cDNA (Fig. 2) is identical to the corresponding region in the genomic sequence from the *Drosophila* Genome Project, except for 36 nucleotides. Most of these substitutions do not lead to changes in the amino acid residues, but three of them do, two of them being conservative (Glu²⁰⁷ → Asp²⁰⁷ and Met⁴⁴⁶ → Leu⁴⁴⁶), whereas one is not (Pro⁴⁰⁹ → Ala⁴⁰⁹). The nucleotide substitutions within the coding region are given in Table 2. In the noncoding region of the cDNA, 16 nucleotide differences were identified. Most of these differences were replacements, but in the 3' noncoding region, one nucleotide insertion

(one within the stretch of Gs at nucleotide positions 4200–4205 of Fig. 2) and eight nucleotide deletions occurred (four TA sequences were deleted within the repetitive TA sequence at nucleotide positions 4622–4636 of Fig. 2, resulting in the TA motif only being repeated 7 times compared with 11 times in the genomic sequence). In the 5' noncoding region, the cDNA sequence contains one A less within the stretch of As starting at nucleotide position –573 of Figure 2. All above-mentioned sequence deviations are probably due to a small genetic difference (~1%) between our own laboratory *D. melanogaster* Canton S. population and the one used in the *Drosophila* Genome Project. The differences are probably not due to PCR artefacts, because they were found in several independent cDNA clones.

Table 2. Codon Differences Between the Coding Region of our Cloned cDNA and that of the Genomic Sequences from the Berkeley *Drosophila* Genome Project

Position of the different nucleotides	Codon in the gene	Codon in the cDNA residue	Change in amino acid
498	CGG	CGA	–
507	GGT	GGG	–
585	GAG	GAA	–
621	GAG	GAT	Glu → Asp
750	CTT	CTC	–
1017	TCT	TCC	–
1086	CTG	CTA	–
1209	CCA	CCT	–
1225	CCA	GCA	Pro → Ala
1336	ATG	CTG	Met → Leu
1806	AGA	AGG	–
1827	TCG	TCT	–
1854	CTA	CTC	–
2313	AAC	AAT	–
2337	CTC	CTT	–
2349	TCA	TCT	–
2418	CTG	CTC	–
2424	ATT	ATC	–
2451	TTG	TTA	–
3120	GTC	GTT	–
3198	CCA	CCC	–
3285	ACA	ACG	–
3399	GCC	GCA	–
3474	AAT	AAC	–
3534	CCA	CCG	–
3687	GCC	GCG	–
3711	TTC	TTT	–
3771	AAC	AAT	–
3855	CGC	CGG	–
3864	GTA	GTC	–
3867	CGC	CGT	–
3873	CTA	CTG	–
3981	AGT	AGC	–
4053	CCT	CCA	–
4059	CCC	CCT	–
4071	GAT	GAC	–

The position of the changed nucleotide (Fig. 2) is given in the first column, the affected codon in the genomic sequence in the second column, and the cDNA in the third column. Most nucleotide differences do not lead to a difference in amino acid residue (fourth column).

Chromosomal Localization

The chromosomal localization of the two P1 clones (Fig. 1C) have been determined by the Berkeley *Drosophila* Genome Project to be on chromosome 2L, position 34E-F (<http://www.fruitfly.org/>).

Southern Blot Analysis

A Southern blot analysis, using a cDNA probe representing most of the coding region of the receptor (including the Leu-rich repeats 10–18 and the seven-transmembrane domain) and *Bam*HI-, *Eco*RV-, *Sac*I-, or *Sal*I-digested fragments of *Drosophila* genomic DNA, showed single hybridizing bands (Fig. 6). The sizes of these single bands fully agreed with the genomic restriction maps of the P1 clones DS01514 and DS00180, suggesting that a single gene codes for DGLR-2. Digestion with *Xba*I yielded, in addition to the expected band of ~11 kb, a slightly weaker hybridizing band of ~30 kb (Fig. 6). This extra band might be due to partial

Figure 6 Southern blot analysis. Genomic DNA from *D. melanogaster* Canton S. was digested with one of five restriction enzymes (*Bam*HI, *Eco*RV, *Sac*I, *Sal*I, and *Xba*I). After electrophoresis and blotting, the genomic fragments were hybridized with a cDNA fragment coding for the Leu-rich repeats 10–18 and the transmembrane domain of DLGR-2. The size of the markers (left) is in kb. All lanes show a single hybridization band with exception of the lane containing the *Xba*I fragment.

digestion of the genomic DNA, or to genotypic variations, with the result that one of the *Xba*I sites in the DLGR-2 gene region is only present in part of our *Drosophila* population.

Developmental Regulation of the *Drosophila* Receptor
mRNA isolated from different developmental stages was analyzed in a Northern blot for the presence of DLGR-2 mRNA, using a cDNA probe coding for the seven-transmembrane domain. This showed that the receptor gene was only expressed in embryos and pupae, but not in larvae or adult flies (Fig. 7). The transcript size of ~5.5 kb corresponded well with the size of the cloned receptor cDNA (Fig. 2). The blot was also hybridized with a probe coding for the ribosomal protein RP49 to check for uniform mRNA loading (Fig. 7).

Isolation of a Knock-Out Mutant

We screened several *Drosophila* mutants from the Bloomington Stock Center (University of Indiana, Bloomington) that were known to have a P element insertion close to the chromosomal region of the DLGR-2 gene. This screening was performed using a P element-specific primer together with one of several gene-specific primers, covering the complete receptor sequence. One mutant (P919) had a P element inserted in exon 1 of the DLGR-2 gene (between nucleotide positions –584 and –583 of Fig. 2). In a cross between flies that were heterozygous for this mutation, about one-quarter of the offspring died around the time of

hatching, suggesting that the mutation is homozygous lethal and that it roughly follows mendelian genetics (Table 3). This was confirmed by PCR, showing that all animals that were homozygous for the mutation indeed died around the time of hatching. On the other hand, all larvae that survived were either heterozygous or did not carry the mutation.

We also investigated whether the observed lethality was caused by the mutated DLGR-2 gene, or by an eventual second mutation located on the same chromosome, which, therefore, would always accompany the DLGR-2 gene during our crossing experiments. To establish this, we carried out classical complementation experiments (Fig. 8), where heterozygous DLGR-2 mutant flies (containing one chromosome 2 with the P element inserted into the DLGR-2 gene and the other chromosome 2 with the dominant CyO mutation, which gives the heterozygous flies a “curly wing” phenotype) were crossed with 15 different heterozygous mutants, carrying single, well-defined deletions close to or including the DLGR-2 gene region (these flies contained one chromosome 2 with a well-defined deletion and the other chromosome 2 without a deletion, but with the dominant CyO mutation). We found that the various deletion mutants could not be rescued by our P element insertion mutant P919 if their deletions included the chromosome 2L map coordinates 34E5-F1, which is exactly the region where the DLGR-2 gene is located. In fact, this independent genetic localization of the defect in mutant P919 is even more precise than the chromosomal localization carried out by the Berkeley *Drosophila* Genome Project of the two P1 clones, containing the DLGR-2 gene, which was 34E-F. These results, therefore, strongly indicate that the le-

Figure 7 Northern blot analysis of the expression of the DLGR-2 gene at several developmental stages. Marker size (left) is given in kb. (A) Poly(A)⁺ RNA from each developmental stage was hybridized with a cDNA fragment coding for the seven-transmembrane domain of DLGR-2. This Northern blot shows that mRNA is only present in embryos and pupae, not in larvae and adult (mixed male and female) flies. (B) Hybridization of the same blot as in A with a cDNA probe coding for RP49. The RP49 gene is regarded to be expressed in all developmental stages (O’Connell and Rosbash 1984; Kerrebrock et al. 1995).

Table 3. Distribution of Genotypes Among the Offspring from a Cross Between Male and Female Flies, Being Heterozygous for the P Element Insertion in Exon 1 of the DLGR-2 Gene

	Dead embryos	Third instar larvae	Total
P/P	23	0	23
P/+	2	42	44
+/+	1	20	21

Of 100 eggs collected from a cross, 26 animals died during embryonic development (including both unhatched eggs and larvae that died within 1 hr after hatching), 69 developed into third instar larvae. Five animals disappeared during the experiments. All animals were investigated by PCR to reveal their genotype. For seven of the third instar larvae, the PCR gave no products, so the genotype of these animals is unknown. The dead embryos of the genotype P/+ and +/+ could easily be distinguished from the P/P animals by their less-developed appearance. Dead P/P animals had the gross anatomy of normal, wild-type first-instar larvae. The crossing experiment shows that the mutation is homozygous lethal and that it roughly follows mendelian genetics.

Figure 8 Results of crossing 15 different deficiency stains (carrying a well-defined deletion in an area of chromosome 2 close to or including the DLGR-2 gene, see Table 4) with mutant P919 that carries a P element insertion in the DLGR-2 gene. (A) Schematic description of the crossings. (P) The DLGR-2 gene, carrying the P element insertion; (Cy⁻) curly wing mutation; (D) deletion mutation. The Cy⁻ mutation is dominant and homozygous lethal. Therefore, if the offspring only consists of curly wing flies (P/Cy⁻ or Cy⁻/D), the P/D combination is nonviable and the deletion can not be rescued by the P element-inserted DLGR-2 region. On the other hand, if the offspring contains flies with normal flat wings (P/D), the deletion mutant can be rescued. (B) Map of the deletion strains used in the rescue experiments (horizontal lines). The abscissa shows the region 34A–35F of *Drosophila* chromosome 2L. The stock numbers of the mutants are given at right (ordinate), together with the information about whether the deletion mutant can be rescued (+) or not (–) in a cross with mutant fly P919. For stock numbers 45490 and 42850 only the left ends of the deletions are shown, as these deletions are very large. The vertical shaded bar indicates the borders of the chromosomal region determined by the inability of mutant P919 to rescue the deletion mutants. This region is 34E5–F1, which is exactly the region where the DLGR-2 gene has been located. This is an independent and strong indication that the P element that we have earlier shown to be inserted in the DLGR-2 gene, is the only cause of lethality in mutant P919.

thality found in our homozygous DLGR-2 gene mutants (Table 3) is indeed caused by a defect in the DLGR-2 gene itself.

When we investigated the gross anatomy of the homozygous DLGR-2 mutants around hatching, we found that the animals looked like fully developed first instar larvae, having normal segmentation, tracheal trees, mouth hooks, denticle belts, and no obvious defects in the gastrointestinal tract.

DISCUSSION

Mammals have at least three glycoprotein hormone receptors (the TSH, FSH, and LH/CG receptors) and two other LGRs for which the ligands yet are unknown (Hsu et al. 1998; McDonald et al. 1998; Hermey et al.

1999). It was, therefore, interesting to find that *Drosophila* also contains at least two LGRs. The first *Drosophila* LGR (DLGR-1) is both structurally and evolutionarily closely related to the mammalian glycoprotein hormone receptors (Hauser et al. 1997). The same is true of DLGR-2 for a number of reasons: (1) There is 49% amino acid residue identity between its seven-transmembrane domain and that of the human TSH receptor (Fig. 3). This identity is high, if one takes into account that the sequence identity between the seven-transmembrane domains of the mammalian TSH, FSH, and LH/CG receptors is only 67%–70% (Salesse et al. 1991); (2) the amino terminus has Leu-rich repeats, similar to those of the mammalian glycoprotein hormone receptors (Fig. 4); and (3) four introns in its gene

occur at the same positions and have the same intron phasing as four introns in the mammalian glycoprotein hormone receptors.

DLGR-2 is not more related to DLGR-1 than to the mammalian glycoprotein hormone receptors. This is evident from both the sequence of its transmembrane domain (45% amino acid residue identity between the two *Drosophila* receptors), the unusual large number of Leu-rich repeats in its extracellular amino terminus (18 in DLGR-2 and 9 in DLGR-1; Fig. 9), and its genomic organization (no introns present in the gene region coding for the seven-transmembrane domain of DLGR-2 and four introns in the corresponding region of DLGR-1; Fig. 9).

The genomic organization of the two *Drosophila* receptor genes gives some interesting information. Previously, it has been assumed that the glycoprotein hormone receptor genes originated from exon shuffling of an intronless gene region coding for the seven-transmembrane domain and an intron-rich region coding for the Leu-rich repeats and the rest of receptor amino terminus (Gross et al. 1991; Tsai-Morris et al. 1991; Heckert et al. 1992). The gene structures of all mammalian glycoprotein hormone receptors (Fig. 9C) and the gene structure of the sea anemone LGR, which should closely resemble that of the ancestral LGR gene (Vibede et al. 1998; Fig. 9D) are in accordance with this idea. Also, the gene structure of DLGR-2 fits the original idea of the evolution of the LGRs (Fig. 9A). The gene structure of DLGR-1, however, conflicts with this model, because it contains four introns in its region coding for the seven-transmembrane domain (Fig. 9B). If one accepts the hypothesis that the number of introns tends to decrease, instead of increase during eukaryote evolution (Gilbert et al. 1986), then both DLGR-1 and DLGR-2 would have originated from a single gene containing at least four exons in its region coding for the seven-transmembrane domain. The same would be true for the mammalian and the sea anemone receptors (Vibede et al. 1998; Fig. 9). Thus, *Drosophila* contains at least two classes of LGR genes, one class (to which the DLGR-1 gene belongs) would be more closely related to the ancestral gene, whereas the other class (to which the DLGR-2 gene belongs) would have lost most of its introns in the region coding for the transmembrane domain, thereby resembling the known genes of the other LGRs.

DLGR-2 is considerably larger than the mammalian TSH, FSH, and LH/CG receptors (Fig. 3 and Fig. 9; Salesse et al. 1991). This is mainly due to the presence of extra Leu-rich repeats in the amino terminus and the presence of a large intracellular carboxyl terminus (Fig. 9A). The extra Leu-rich repeats have probably originated by exon shuffling, because they are all (in single or multiple copies) represented by separate exons in the gene (Fig. 1B). The length of the receptor

amino terminus and the number and amino acid sequence of its Leu-rich repeats place DLGR-2 structurally closer to the two mammalian orphan receptors LGR-4 and LGR-5 than to the three mammalian glycoprotein hormone receptors. When the transmembrane regions are compared, however, DLGR-2 is more closely related to the mammalian TSH, FSH, and LH/CG receptors (~50% sequence identity). DLGR-2, therefore, appears to be a naturally occurring chimera, with its amino terminus more closely resembling LGR-4 and LGR-5 and its transmembrane region more closely resembling the three mammalian glycoprotein hormone receptors.

The Leu-rich repeats of the mammalian glycoprotein hormone receptors probably form a horseshoe-like structure to which the glycoprotein hormone binds at the inner, concave side (Jiang et al. 1995; Kajava et al. 1995). We assume that the Leu-rich repeats of DLGR-2 also form such a horseshoe, because these repeats very closely resemble those of the mammalian receptors (Fig. 4). Because DLGR-2 has twice as many Leu-rich repeats as the mammalian glycoprotein hormone receptors, we assume that the horseshoe is correspondingly larger. This might indicate that the ligand is somewhat different from the mammalian glycoprotein hormones.

Two of the three mammalian glycoprotein hormone receptors are involved in reproduction. Therefore, we monitored the expression of the DLGR-2 gene during several developmental stages of *Drosophila* to see whether it was expressed in adult male or female flies. We found, however, that the receptor was only expressed in embryos and pupae, that is, in stages where there are active cell division, cell differentiation, and other forms of development but not in the three larval stages of *Drosophila* or adult male or female flies (Fig. 7). This exclusive expression of DLGR-2 in embryos and pupae already points to an important role for the receptor in development. This idea was confirmed when we investigated the phenotype of the homozygous knock-out mutants of the DLGR-2 gene. These mutant flies have a P element insertion in exon 1, which is the noncoding 5' region of the DLGR-2 gene, meaning that the homozygous mutants are devoid of DLGR-2 (because the P element contains a gene, including stop codons; Hazelrigg et al. 1984). We observed that all mutants died around the time of hatching. It is interesting that the homozygous mutants had the overall appearance of normal larvae. This means that DLGR-2 does not play a role in morphogenesis but rather is important in a more subtle developmental process. This process must be absent in hatched first-, second-, and third-instar larvae and in adult flies, because the receptor is not expressed in these stages. This exclusive role of an LGR in development is unique and has not been described previ-

ously for the other known glycoprotein hormone receptors.

METHODS

Animals

Wild-type *Drosophila melanogaster* Canton S. were reared under standard conditions (Roberts 1986).

Database Screening

Amino acid residues 703–728 of DLGR-1 (Hauser et al. 1997) were used as a probe for the electronic screening of the Berkeley *Drosophila* Gene Project database, using the NCBI Search Engine BLAST (Altschul et al. 1990). The same search engine was used for homology screening.

Preparation of Poly(A)⁺ RNA and cDNA Synthesis

Poly(A)⁺ RNA from various *Drosophila* stages was purified with the Oligotex Direct mRNA Kit from Qiagen. Oligo(dT)-primed cDNA was synthesized from 0.4 µg poly(A)⁺ RNA, as recommended by the manufacturer, using the RT-PCR Kit (Stratagene) and an oligo (dT) primer supplied with the kit.

PCR

The PCR reactions used for cloning of the cDNA were carried out in a 50 µl volume as described in Sambrook et al. (1989), with the exception that a final concentration of 1.75 mM MgCl₂ was used. The template was 5 µl of a first-strand cDNA reaction mixture. Various primers were used, based on the predicted exon sequences of the DLGR-2 gene (see Results). The PCR products were separated on 2% agarose gel, and bands of the expected size were isolated (Qiaquick extraction kit, Qiagen), subcloned into PCR 2.1 with the Original TA Cloning Kit (Invitrogen), and sequenced.

The final full-length products were generated by use of the Expand Long Template PCR System (Boehringer Mannheim). Two independent PCR reactions were carried out. In each of these PCRs, the 50-µl reaction mixture consisted of Buffer System 1, 2 mM final concentration of MgCl₂, 500 µM of each dNTP, 0.3 µM of each primer (corresponding to positions –560 to –537 and 4661 to 4682 of Fig. 2 for the first reaction, and positions –1 to 21 and 4068 to 4087 for the second reaction), 275 mU of Long Range Polymerase, and cDNA from *Drosophila* embryos as a template. Cycling parameters were 2 min of initial denaturation at 94°C; 10 cycles of the following step program: 94°C for 15 sec, 60°C for 30 sec, 68°C for 4 min; then 25 cycles of 94°C for 15 sec, 60°C for 30 sec, 68°C for 4 min. This last 4-min period was increased with 20 sec for every new cycle.

3'-RACE

First-strand cDNA synthesis was performed according to the protocol for the 5'/3'-RACE Kit (Boehringer Mannheim), using 2 µg of poly(A)⁺ RNA from 16–24-hr-old *Drosophila* embryos as a template, and the oligo d(T) primer from the kit. This cDNA was used directly for PCR amplification with a sense DLGR-2-specific primer (positions 2985 to 3006 from Fig. 2) in combination with a PCR anchor primer supplied with the kit. A second round of PCR using 2 µl of the first PCR reaction as a template and employing the anchor primer and a second DLGR-2-specific sense primer (positions 3013 to

3033 of Fig. 2) further to the 3' end, was necessary to obtain a specific PCR product. The reaction mixture was in both cases as recommended by the manufacturer. Cycling parameters were as follows 3 min of initial denaturation at 95°C; 10 cycles of 95°C for 30 sec, 62°C for 45 sec, 72°C for 2 min. The reactions were held at 72°C while the anchor primer was added, and then further 25 cycles were run as described above.

5'-RACE

For the first strand cDNA synthesis, 2 µg of poly(A)⁺ RNA from 16–24-hour-old embryos and a DLGR-2-specific antisense primer (position 1149 to 1170 of Fig. 2) were used. 5'-RACE PCR was carried out using the 5'/3'-RACE Kit (Boehringer Mannheim), following the instructions of the manufacturer, and employing two nested antisense primers (positions 837 to 858 and –318 to –297 of Fig. 2).

DNA Sequencing and Sequence Analysis

DNA sequences were determined by the chain termination method (Sanger et al. 1977), using the T7 Sequenase Version 2.0 DNA Sequencing Kit (Amersham Life Science). GC-rich sequences were determined, using the Thermo Sequenase Radiolabeled Terminator Cycle Sequencing Kit (Amersham Life Science). DNA sequence compilation and nucleotide and amino acid sequence comparisons were performed using the Lasergene DNA Software (DNASTAR Inc.).

Promotor Analysis

The genomic sequence was searched for promotor sequences up to 1 kb upstream of the identified transcription start site, employing the TSSG promotor search engine at <http://dot.imgen.bcm.tmc.edu:9331/genefinder/gf.html>.

Prediction of Gene Structures in Genomic DNA

To predict exons, introns, and polyadenylation signals, the GENSCAN web server at <http://stanford.edu/~chis/GENSCAN.html> was used.

Radioactive Labeling of DNA Probes

DNA fragments to be labeled were excised from vector DNA by restriction enzymes and purified by agarose gel electrophoresis. Probes were labeled with [α -³²P]dCTP (3000 Ci/mmol) from Amersham by use of the Ready-To-Go DNA Labeling Beads (Pharmacia Biotech), according to the manual.

Northern Blot Analysis

Poly(A)⁺ (2.5 µg) from various stages of *Drosophila* (obtained from Clontech or purified as described above) were electrophoresed on a gel containing 1% agarose and 0.22 M formaldehyde. The 0.24–9.5-kb RNA Ladder (GIBCO-BRL) was included as a size marker. RNA was capillary transferred onto ZetaProbe membranes (BioRad) and cross-linked as recommended by the manufacturer. Hybridization was carried out as recommended by BioRad for 18 hr at 65°C in the presence of heat-denatured radioactive probe in a final concentration of 1–2 × 10⁶ cpm/ml. This probe corresponded to nucleotide positions 1806 to 3204 of Figure 2. The ribosomal protein 49 (RP49) probe was prepared as described in Hauser et al. (1997).

Southern Blot Analysis

Genomic DNA (10 µg) from *D. melanogaster* was digested with one of the restriction enzymes, *Bam*HI, *Eco*RV, *Sac*I, *Sal*I, or

*Xba*I, and separated on a 0.7% agarose gel. DNA was capillary transferred to a Hybond-N nylon membrane, hybridized, and washed as recommended by the supplier (Amersham). The radioactive probe corresponded to nucleotide positions 1149 to 3083 of Figure 2.

Isolation of a Knock-Out Mutant

Various *Drosophila* mutants from the Bloomington Stock Center were screened by PCR for a P element insertion in the DLGR-2 gene. This screen was performed using a P element-specific primer, 5'-CGACGGGACCACCTTATGTTATTCATCATG-3', along with various DLGR-2 gene-specific primers, covering the complete receptor sequence. For one mutant, P919 with the genotype w[1118]; P{w+[tAR]ry[+t7.2AR]=wA[R]}4-34/CyO, a 1200-bp PCR product was seen, using the exon 1-specific primer 5'-GGCTGTGCCGACAATTGAAC-3' together with the P element-specific primer. The location of the P element in exon 1 of the receptor gene (between nucleotide positions -584 and -583 of Fig. 2), was confirmed by sequencing the PCR product with the primer 5'-GAAATCTCTGTGCCACTG-3'.

To obtain homozygous mutants, male P919 flies were crossed with female virgin Oregon R. flies to get P{w+[tAR]ry[+t7.2AR]=wA[R]}4-34/+ flies. Mating between flies of this last genotype gave the results of Table 3. To investigate the presence of a P-element insertion in the offspring of Table 3, DNA was extracted from single animals by grinding the animal in 100 µl 5% chelex-100 (Bio-Rad Laboratories) in an Eppendorf tube (Sweet et al. 1996). The tube was then incubated for 30 min at 56°C and 8 min at 100°C. After a 3-min centrifugation at 13000 rpm, the supernatant was transferred to a clean Eppendorf tube and stored at -20°C. DNA extract (5–10 µl) was used in a PCR reaction, using the primers 5'-GAGCATAACCCTCTTCTTGT-3' and 5'-TGATCTGGGAGT-TTGGAGTG-3', that spans a 564-bp region in intron 1 of the receptor gene. For unknown reasons, the P919 strain has an insertion of ~184 bp in intron 1 (2120 bp downstream of the transcription start) of that allele that bears the P-element insertion, whereas in the allele that does not bear the P-element insertion, this 184-bp insertion is lacking. Because our two primers lie at each side of the 184-bp insertion, an animal that is homozygous for the P-element insertion will give a PCR product of 751 bp, whereas a homozygous wildtype will give

a PCR product of 567 bp and a heterozygous animal both PCR products.

Complementation Experiments

The deficiency strains of *Drosophila* that were used in our complementation experiments of Figure 8 are given in Table 4. The deficiencies in the heterozygous flies were balanced with the CyO chromosome (see also legend to Fig. 8). After crossing, we verified by PCR that the Cy⁺ offspring (flat wings; P/D, see Fig. 8A) did indeed carry the wild type DLGR-2 gene (D) together with the P element-inserted DLGR-2 gene (P).

Two deficiency strains from the Umeå *Drosophila* Stock Center (42450 and 41624) could be rescued by mutant P919 (which carries a P-element insertion in the DLGR-2 gene), despite the fact that their records indicated a deletion in the DLGR-2 gene area. PCR tests of the Cy⁺ offspring (see Fig. 8A) demonstrated that these flies carried a wild-type DLGR-2 gene together with the P element-inserted DLGR-2 gene, showing that the two deficiency strains either have been mapped incorrectly or have lost their deficient chromosomes. These two strains were, therefore, withdrawn from the complementation experiments.

ACKNOWLEDGMENTS

This work was supported by the Danish Biotechnological Research and Development Programme of the Danish Research Agency, Læge- og Naturvidenskabelige Komité of the Novo Nordisk Foundation and Carlsberg Foundation. We thank Lotte Steffensen for typing the manuscript and for help with Fig. 3.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

Figure 9 Schematic representation of four LGR cDNAs. The regions coding for the seven-transmembrane loops are given as black bars, those for the Leu-rich repeats as gray bars. Only complete Leu-rich repeats (see Fig. 4) are taken into account. Intron positions in the four LGR genes are indicated by arrows. (A) The cDNA coding for DLGR-2. (B) The cDNA coding for DLGR-1 (Hauser et al. 1997). (C) The cDNA coding for the rat FSH receptor (Heckert et al. 1992). (D) The cDNA coding for the LGR from the sea anemone *A. elegantissima* (Nothacker and Grimmelikhuijzen 1993; Vibede et al. 1998).

Table 4. Deficiency Strains of *Drosophila* Used in the Complementation Experiments of Figure 8 to Independently Localize the Deficiency in Mutant P919, which Carries a P Element Insertion in the DLGR-2 Gene

Stock number	Deficient chromosomal region
45490	35E1-2 to 36A6-7
3588	35B4-6 to 35F1-7
42850	35B3 to 38D3-5
3213	35B2-3 to 35D5-7
3078	35B1-3 to 35E6
41612	34E5-F1 to 35C3-9
80842	34F1-2 to 35A2
3211	34E3 to 35D7
41622	34E1-2 to 35B3-5
41900	34D4-6 to 34E5-6
3897	34D2 to 34E3
41600	34D1-2 to 35B9-C1
41623	34C6-7 to 35B9-C1
42800	34C4 to 35A4
3138	34B12-C1 to 35B10-C1

Stock numbers with five digits are from the Umeå *Drosophila* Stock Center (Umeå University, Sweden), those with four digits from the Bloomington *Drosophila* Stock Center (Indiana University, Bloomington). The deficient chromosomal region refers to the left arm of chromosome 2.

REFERENCES

- Altschul, S.F., Gish, W., Miller, W., Meyers, E.W., and Lipman, D.J. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**: 403–410.
- Bhowmick, N., Huang, J.N., Puett, D., Isaacs, N.W., and Lapthorn, A.J. 1996. Determination of residues important in hormone binding to the extracellular domain of the luteinizing hormone/chorionic gonadotropin receptor by site-directed mutagenesis and modeling. *Mol. Endocrinol.* **10**: 1147–1159.
- Cavener, D. 1987. Comparison of the consensus sequence flanking translational start sites in *Drosophila* and vertebrates. *Nucleic Acids Res.* **15**: 1353–1361.
- Gilbert, W., Marchionni, M., and McKnight, G. 1986. On the antiquity of introns. *Cell* **46**: 151–154.
- Gross, B., Misrahi, M., Sar, S., and Milgrom, E. 1991. Composite structure of the human thyrotropin receptor gene. *Biochem. Biophys. Res. Commun.* **177**: 679–687.
- Hauser, F., Nothacker, H.-P., and Grimmelikhuijzen, C.J.P. 1997. Molecular cloning, genomic organization and developmental regulation of a novel receptor from *Drosophila melanogaster* structurally related to members of the thyroid-stimulating hormone, follicle-stimulating hormone, luteinizing hormone/choriogonadotropin receptor family from mammals. *J. Biol. Chem.* **272**: 1002–1010.
- Hauser, F., Søndergaard, L., and Grimmelikhuijzen, C.J.P. 1998. Molecular cloning, genomic organization and developmental regulation of a novel receptor from *Drosophila melanogaster* structurally related to gonadotropin-releasing hormone receptors from vertebrates. *Biochem. Biophys. Res. Commun.* **249**: 822–828.
- Hazelrigg, T., Levis, R., and Rubin, G.M. 1984. Transformation of white locus DNA in *Drosophila*: Dosage compensation, zeste interaction, and position effects. *Cell* **36**: 469–481.
- Heckert, L.L., Daley, I.J., and Griswold, M.D. 1992. Structural organization of the follicle-stimulating hormone receptor gene. *Mol. Endocrinol.* **6**: 70–80.
- Hermey, G., Methner, A., Schaller, H.C., and Hermans-Borgmeyer, I. 1999. Identification of a novel seven-transmembrane receptor with homology to glycoprotein receptors and its expression in the adult and developing mouse. *Biochem. Biophys. Res. Commun.* **254**: 273–279.
- Hsu, S.Y., Liang, S.-G., and Hsueh, A.J.W. 1998. Characterization of two LGR genes homologous to gonadotropin and thyrotropin receptors with extracellular leucine-rich repeats and a G protein-coupled, seven-transmembrane region. *Mol. Endocrinol.* **12**: 1830–1845.
- Jiang, X., Dreano, M., Buckler, D.R., Cheng, S., Ythier, A., Wu, H., Hendrickson, W.A., and El Tayar, N. 1995. Structural predictions for the ligand-binding region of glycoprotein hormone receptors and the nature of hormone-receptor interactions. *Structure* **3**: 1341–1353.
- Kajava, A.V., Vassart, G., and Wodak, S.J. 1995. Modeling of the three-dimensional structure of proteins with the typical leucine-rich repeats. *Structure* **3**: 867–877.
- Kemp, B.E., and Pearson, R.B. 1990. Protein-kinase recognition sequence motifs. *Trends Biochem. Sci.* **15**: 342–346.
- Kerrebrock, A.W., Moore, D.P., Wu, J.S., and Orr-Weaver, T.L. 1995. Mei-S332, a *Drosophila* protein required for sister-chromatid cohesion, can localize to meiotic centromere regions. *Cell* **83**: 247–256.
- Kobe, B., and Deisenhofer, J. 1994. The leucine-rich repeat—A versatile binding motif. *Trends Biochem. Sci.* **19**: 415–421.
- . 1995. Proteins with leucine-rich repeats. *Curr. Opin. Struct. Biol.* **5**: 409–416.
- Koo, Y.B., Ji, I., Slaughter, R.G., and Ji, T.H. 1991. Structure of the luteinizing hormone receptor gene and multiple exons of the coding sequence. *Endocrinology* **128**: 2297–2308.
- McFarland, K.C., Sprengel, R., Phillips, H.S., Köhler, M., Rosembly, N., Nikolics, K., Segaloff, D.L., and Seeburg, P.H. 1989. Lutropin-choriogonadotropin receptor: An unusual member of the G protein-coupled receptor family. *Science* **245**: 494–499.
- McDonald, T., Wang, R., Bailey, W., Xie, G., Chen, F., Caskey, C.T., and Liu, Q. 1998. Identification and cloning of an orphan G protein-coupled receptor of the glycoprotein hormone receptor subfamily. *Biochem. Biophys. Res. Commun.* **247**: 266–270.
- Misrahi, M., Loosfelt, H., Atger, M., Sar, S., Guiochon-Mantel, A., and Milgrom, E. 1990. Cloning, sequencing and expression of human TSH receptor. *Biochem. Biophys. Res. Commun.* **166**: 394–403.
- Nothacker, H.-P., and Grimmelikhuijzen, C.J.P. 1993. Molecular cloning of a novel, putative G protein-coupled receptor from sea anemones structurally related to members of the FSH, TSH, LH/CG receptor family from mammals. *Biochem. Biophys. Res. Commun.* **197**: 1062–1069.
- O'Connell, P.O., and Rosbash, M. 1984. Sequence, structure, and codon preference of the *Drosophila* ribosomal protein 49 gene. *Nucleic Acids Res.* **12**: 5495–5513.
- Proudfoot, N. 1991. Poly(A⁺) signals. *Cell* **64**: 671–674.
- Roberts, D.B. 1986. *Drosophila—A practical approach*. IRL Press, Oxford, UK.
- Salesse, R., Remy, J.J., Levin, J.M., Jallat, B., and Garnier, J. 1991. Towards understanding the glycoprotein hormone receptors. *Biochimie* **73**: 109–120.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. 1989. *Molecular cloning: A laboratory manual*, 2nd edition. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Sanger, F., Nicklen, S., and Coulson, A.R. 1977. DNA sequencing with chain-termination inhibitors. *Proc. Natl. Acad. Sci. USA* **74**: 5463–5467.
- Sprengel, R., Braun, T., Nikolics, K., Segaloff, D.L., and Seeburg, P.H. 1990. The testicular receptor for follicle stimulating hormone: Structure and functional expression of cloned cDNA. *Mol. Endocrinol.* **4**: 525–530.
- Sweet, D., Lorente, M., Valenzuela, A., Lorente, J.A., and Alvarez, J.C. 1996. Increasing DNA extraction yield from saliva stains with a modified Chelex method. *Forensic Sci. Int.* **83**: 167–177.
- Tensen, C.P., van Kesteren, E.R., Planta, R.J., Cox, K.J.A., Burke, J.F., van Heerikhuizen, H., and Vreugdenhil, E. 1994. A G protein-coupled receptor with low density lipoprotein-binding motifs suggests a role for lipoproteins in G-linked signal transduction. *Proc. Natl. Acad. Sci. USA* **91**: 4816–4820.
- Tsai-Morris, C.H., Buczko, E., Wang, W., Xie, X.-Z., and Dufau, M.L. 1991. Structural organization of the rat luteinizing hormone (LH) receptor gene. *J. Biol. Chem.* **266**: 11355–11359.
- Vibede, N., Hauser, F., Williamson, M., and Grimmelikhuijzen, C.J.P. 1998. Genomic organization of a receptor from sea anemones, structurally and evolutionarily related to glycoprotein hormone receptors from mammals. *Biochem. Biophys. Res. Commun.* **252**: 497–501.
- von Heijne, G. 1986. A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* **14**: 4683–4690.
- . 1990. The signal peptide. *J. Membrane Biol.* **115**: 195–201.

Received November 10, 1999; accepted in revised form May 17, 2000.

Molecular Cloning, Genomic Organization, Developmental Regulation, and a Knock-out Mutant of a Novel Leu-rich Repeats-containing G Protein-coupled Receptor (DLGR-2) from *Drosophila melanogaster*

Kathrine Krageskov Eriksen, Frank Hauser, Morten Schiøtt, et al.

Genome Res. 2000 10: 924-938

Access the most recent version at doi:[10.1101/gr.10.7.924](https://doi.org/10.1101/gr.10.7.924)

References

This article cites 33 articles, 5 of which can be accessed free at:
<http://genome.cshlp.org/content/10/7/924.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Affordable, Accurate
Sequencing.

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>