Research

Genome-wide patterns of natural variation reveal strong selective sweeps and ongoing genomic conflict in *Drosophila mauritiana*

Viola Nolte, Ram Vinay Pandey, Robert Kofler, and Christian Schlötterer¹

Institut für Populationsgenetik, Vetmeduni Vienna, 1210 Wien, Austria

Although it is well understood that selection shapes the polymorphism pattern in *Drosophila*, signatures of classic selective sweeps are scarce. Here, we focus on *Drosophila mauritiana*, an island endemic, which is closely related to *Drosophila melanogaster*. Based on a new, annotated genome sequence, we characterized the genome-wide polymorphism by sequencing pooled individuals (Pool-seq). We show that the interplay between selection and recombination results in a genome-wide polymorphism pattern characteristic for *D. mauritiana*. Two large genomic regions (>500 kb) showed the signature of almost complete selective sweeps. We propose that the absence of population structure and limited geographic distribution could explain why such pronounced sweep patterns are restricted to *D. mauritiana*. Further evidence for strong adaptive evolution was detected for several nucleoporin genes, some of which were not previously identified as genes involved in genomic conflict. Since this adaptive evolution is continuing after the split of *D. mauritiana* and *Drosophila simulans*, we conclude that genomic conflict is not restricted to short episodes, but rather an ongoing process in *Drosophila*.

[Supplemental material is available for this article.]

Intragenomic conflict describes the phenomenon that within an organism some genetic elements (e.g., segregation distorters) increase their transmission at the expense of others (Werren 2011). Due to the preferential transmission, such elements spread in the population and can leave a characteristic trace of strongly reduced variability in the genome that resembles a selective sweep (Derome et al. 2004). Population genetic analyses of segregation distortion systems in *Drosophila* did not find a molecular signature similar to a classic selective sweep (Derome et al. 2004, 2008; Presgraves et al. 2009; Kingan et al. 2010; Bastide et al. 2011). The patterns of variability instead resembled partial selective sweeps, suggesting that the genetic element increased in frequency but did not reach fixation. This observation is consistent with the fact that elements of intragenomic conflict are frequently deleterious when homozygous (Wallace 1948; Curtsinger and Feldman 1980) or that suppressors of the intragenomic conflict have evolved (Hamilton 1967).

In the *Drosophila melanogaster* complex, only a small number of genes involved in intragenomic conflict have been identified within natural populations (e.g., Sandler et al. 1959; Mercot et al. 1995). While this may suggest that intragenomic conflict is a relatively rare event, it needs to be considered that there is a strong ascertainment bias: The rapid spread of driver alleles is either prevented by a quick fixation of suppressor alleles, or, in case of sex chromosome-linked segregation distorters, populations with an advanced intragenomic conflict become extinct (Gershenson 1928; Hamilton 1967; Lyttle 1977). In both cases, past episodes of genomic conflict cannot be recognized in an intraspecific polymorphism analysis.

Indeed, consistent with the idea that genomic conflict is a common phenomenon, detailed analysis of hybrids showed that

“speciation” genes tend to be involved in intragenomic conflict, but their effect could be only detected in hybrids (Perez et al. 1993; Dermitzakis et al. 2000; Tao et al. 2001; Presgraves et al. 2003; Phadnis and Orr 2009; Tang and Presgraves 2009).

Several genes involved in intragenomic conflict in *Drosophila* were discovered in the *Drosophila simulans* clade that consists of three recently diverged species, the cosmopolitan *D. simulans* and the island endemics *Drosophila mauritiana* and *Drosophila sechellia*. *D. mauritiana* was the first species for which a “speciation” gene could be characterized at the molecular level: In hybrid crosses with *D. simulans*, the *Odysseus* (*OdsH*) allele of *D. mauritiana* together with additional tightly linked factors causes hybrid male sterility in the F₁ generation (Perez and Wu 1995; Ting et al. 1998) and has been later identified as a gene involved in genomic conflict (Bayes and Malik 2009). Another *D. mauritiana* gene, *too much yin* (*tmy*), causes both, hybrid male sterility and segregation distortion in crosses between *D. mauritiana* and *D. simulans* (Tao et al. 2001), whereas the heterochromatic *hlx* locus causes hybrid lethality between *D. mauritiana* and both of its sister species (Cattani and Presgraves 2009).

Additional elements of intragenomic conflict have been identified between the more distantly related *D. melanogaster* and *D. simulans*, in which the interaction between the genes *Hmr* and *Lhr* contributes to hybrid male lethality in crosses between *D. melanogaster* and *D. simulans* (Brideau et al. 2006; Maheshwari and Barbash 2012).

The *D. simulans* alleles of two nucleoporin genes, *Nup96*² and *Nup160*, cause recessive male lethality when crossed to a *D. melanogaster* X chromosome (Presgraves et al. 2003; Tang and Presgraves 2009), a phenomenon that has been also linked to

¹Corresponding author

E-mail christian.schloetterer@vetmeduni.ac.at

Article published online before print. Article, supplemental material, and publication date are at <http://www.genome.org/cgi/doi/10.1101/gr.139873.112>. Freely available online through the *Genome Research* Open Access option.

²Throughout the manuscript, we refer to “*Nup96* gene” as the part of the *Nup98-96* gene that corresponds to amino acid residues 1029–1961 in the resulting protein; this part of the protein is frequently referred to as “NUP96.” Similarly, we refer to “*Nup98* gene” as the part of the *Nup98-96* gene that corresponds to amino acid residues 1–1028 in the resulting protein; this part of the protein is frequently referred to as “NUP98,” e.g., in Presgraves et al. (2003).

genomic conflict (Presgraves 2007; Presgraves and Stephan 2007). While the NUP96 protein is highly conserved between *D. simulans* and *D. mauritiana*, the *D. mauritiana* allele of *Nup96* has no hybrid-lethal effect, which suggests more complex genetic interactions leading to *Nup96*-dependent incompatibility (Barbash 2007).

Despite the importance of *D. mauritiana* as a model for understanding the genetic basis of speciation, an annotated genome sequence is not yet available. Using de novo assembly, we generated a draft genome of *D. mauritiana* and estimated genome-wide polymorphism patterns from Pool-seq data. Our data show the impact of genes involved in genomic conflict on the evolution of the *D. mauritiana* lineage. Nucleoporin genes, implicated in hybrid incompatibilities that have evolved between *D. simulans* and *D. melanogaster*, are possible targets of recurrent positive selection due to ongoing genomic conflict (Presgraves and Stephan 2007). Unlike previous genome-wide polymorphism surveys of *D. simulans* and *D. melanogaster* (Begun et al. 2007; Langley et al. 2012), we find that in the *D. mauritiana* lineage, nucleoporins are among the genes showing the strongest evidence of recurrent adaptive evolution. Furthermore, the presence of a pair of meiotic drive genes and a “speciation” gene at the center of two valleys of strongly reduced variability suggests that these sweeps have been caused by genes involved in genomic conflict.

Results

The recent advances in sequencing technology provide the opportunity to perform population genetic analyses on a genome scale. Even for species with no available reference genome, it has become feasible to generate draft genomes that can be used for population genomic analysis. Here we pursue this strategy for *D. mauritiana*, for which no annotated reference genome is available yet. We sequenced the *D. mauritiana* strain MS17 using a mixture of single-end and paired-end Illumina reads (Supplemental Table S1), and assembled and annotated the draft genome (for further details, see Supplemental Results). To study the impact of selection on the polymorphism pattern in *D. mauritiana*, we sequenced a pool of 107 isofemale lines (Supplemental Table S2).

Faster rate of evolution on the X chromosome

Since the X chromosome is hemizygous in males, rates of sequence evolution can be contrasted between the X chromosome and the autosomes to shed some light on the operating selective forces. Under the assumption that new mutations are recessive, population genetics theory predicts a higher rate of evolution on the X chromosome than on the autosomes (Maynard Smith and Haigh 1974; Charlesworth et al. 1987).

Figure 1. Mean pairwise divergence (D_{xy}) along each major chromosomal arm between species of the *D. melanogaster* complex. The following species pairs are shown: *D. mauritiana*–*D. simulans* (yellow), *D. sechellia*–*D. simulans* (orange), *D. mauritiana*–*D. sechellia* (green), *D. simulans*–*D. melanogaster* (gray), and *D. mauritiana*–*D. melanogaster* (red). The sliding window analysis was performed using a window size of 500 kb and a step size of 100 kb; chromosomal coordinates are those of *D. mauritiana*.

Table 1. Mean pairwise divergence (D_{xy}) between *D. mauritiana*, *D. simulans*, and *D. melanogaster* based on alignments of repeat-masked genomes

	<i>D. mauritiana</i> – <i>D. simulans</i> ^a	<i>D. mauritiana</i> – <i>D. melanogaster</i>	<i>D. simulans</i> ^a – <i>D. melanogaster</i>
X	0.0168	0.0548	0.0540
2L	0.0216	0.0503	0.0496
2R	0.0201	0.0481	0.0474
3L	0.0211	0.0496	0.0488
3R	0.0198	0.0482	0.0474
4	0.0121	0.0781	0.0785
Mean D_{xy} autosomes	0.0206	0.0491	0.0483
% D_{xy} X of autosomes	81.5	111.7	111.9
P-value for difference between X chromosome and autosomes (Wilcoxon rank-sum test)	$<2.2 \times 10^{-16}$	$<2.2 \times 10^{-16}$	$<2.2 \times 10^{-16}$

Means are based on nonoverlapping 10-kb windows.

^aAssembly based on the African *D. simulans* strain Kib32.

Mean pairwise divergence (D_{xy}) between *D. melanogaster* and *D. mauritiana* is significantly higher on the X chromosome (mean D_{xy} = 0.0548) than on the major autosomes (mean D_{xy} of the major autosomal arms = 0.0491, two-tailed Wilcoxon rank-sum test based on nonoverlapping 10-kb windows, $P < 2.2 \times 10^{-16}$). The same pattern is observed for the species pair *D. melanogaster* and *D. simulans* (mean D_{xy} on the X chromosome = 0.0540, mean D_{xy} on the major autosomes = 0.0483) (Fig. 1; Table 1), which is consistent with the genome-wide data of Begun et al. (2007).

Interestingly, comparisons within the *D. simulans* clade show the opposite pattern: Mean pairwise divergence between *D. mauritiana* and *D. simulans*, for example, is higher on the major autosomal arms (D_{xy} = 0.0206) than on the X chromosome (D_{xy} = 0.0168, two-tailed Wilcoxon rank-sum test based on nonoverlapping 10-kb windows, $P < 2.2 \times 10^{-16}$). This pattern of a higher divergence on the autosomes holds for all comparisons among species of the *D. simulans* clade (Fig. 1; Table 1; Supplemental Table S3) and has been noted previously in the *D. simulans*–*D. sechellia* comparison (Singh et al. 2008). Reduced divergence on the X chromosome compared with the autosomes could be explained by hybridization between species of the *D. simulans* clade (Ballard 2000; Morton et al. 2004; Nunes et al. 2010). If the X chromosome experiences more interspecific gene flow than the autosomes, this would result in a higher divergence on the autosomes. Nevertheless, since Garrigan et al. (2012) found twice as many fragments with a putative introgression signal on the autosomes than on the X chromosome (Garrigan et al. 2012), we consider this scenario not very likely.

Another cause of lower divergence on the X chromosome could be less ancestral polymorphism on the X chromosome than on the autosomes (Singh et al. 2008). Alternatively, selection on the short time scale could be mainly operating on standing variation rather than on new mutations (Orr and Betancourt 2001). Hence, assuming that in the *D. simulans* clade selection acts mainly on shared standing variation, the time scale may be too short to notice a higher sub-

stitution rate on the X chromosome. In contrast, comparisons involving *D. melanogaster* encompass longer time intervals allowing for more novel mutations and fewer shared mutations, which makes the higher substitution rate on the X chromosome visible.

Impact of the recombination landscape on the partitioning of variation

When comparing levels of polymorphism in *D. mauritiana* to those in *D. melanogaster*, we find that *D. mauritiana* is 40%–50% more variable than a cosmopolitan *D. melanogaster* population (Table 2; for further details, see Supplemental Results). While a higher level of overall

polymorphism has been suggested previously based on a small number of loci (Hey and Kliman 1993; Moriyama and Powell 1996), our Pool-seq data allow us to address the distribution of variability along all chromosomal arms.

It is well understood that the recombination landscape in *D. melanogaster* varies along the chromosomes. Both telomeres and centromeres have a reduced recombination rate, but while the drop in recombination rate is abrupt at the telomeres, a gradual decrease in recombination rate over several megabases is observed toward the centromere on all major autosomal arms (True et al. 1996). *D. mauritiana* not only has a higher genome-wide recombination rate but also shows an important difference in the recombination landscape: Instead of an extended gradual decrease in recombination rate near the centromere, the suppression of recombination is restricted to a very small pericentric region (True et al. 1996).

Since the correlation between recombination rate and variability is well-studied in *D. melanogaster* (Begun and Aquadro 1992; Hudson 1994), we were interested if the change in recombination landscape affects the pattern of variability in genomic regions toward the centromere. Figure 2 shows that in *D. melanogaster*, polymorphism declines toward the centromeres, whereas in *D. mauritiana*, levels of variability remain almost flat throughout

Table 2. Mean nucleotide diversity (π) and mean Tajima's D per chromosomal arm in *D. mauritiana* compared with the *D. melanogaster* population from Portugal

	Mean π <i>D. mauritiana</i> ^a	Mean π <i>D. melanogaster</i> ^a	Tajima's D <i>D. mauritiana</i> ^a	Tajima's D <i>D. melanogaster</i> ^a
X	0.0059	0.0039	–1.94	–1.76
2L	0.0092	0.0077	–1.70	–1.21
2R	0.0087	0.0060	–1.71	–1.41
3L	0.0095	0.0066	–1.67	–1.40
3R	0.0086	0.0059	–1.73	–1.50
4	0.0011	0.0009	–2.20	–2.42
Mean autosomes	0.0090	0.0066	–1.70	–1.38
% X of autosomes	65.7	60.1		
% X* 4/3 of autosomes	87.7	80.1		

Both data sets were repeat-masked, and means were calculated from nonoverlapping 10-kb windows. ^aThe *D. mauritiana* data set was analyzed using a minimum count of 3, a minimum coverage of 6, and a maximum coverage of 250; the *D. melanogaster* data set was analyzed using a minimum count of 2, a minimum coverage of 4, and a maximum coverage of 150. For calculation of Tajima's D , both data sets were subsampled to a 30-fold coverage and analyzed without correcting for sequencing errors and multiple sampling.

Figure 2. Nucleotide diversity (π) along the major chromosomal arms in *D. mauritiana* (red line) and *D. melanogaster* (gray line). The sliding window analysis was performed using 500-kb windows with a step size of 100 kb; chromosomal coordinates have been adjusted to *D. melanogaster*.

the entire chromosome. Moreover, not only is the level of variability reduced in low-recombining regions in *D. melanogaster*, but the allele frequency spectrum is affected as well. Tajima's D (Tajima 1989) is a frequently used summary statistic, which describes deviations of the allele frequency spectrum from the standard neutral model. We plotted Tajima's D along the *D. melanogaster* and *D. mauritiana* chromosomes and observed more negative Tajima's D values toward the centromere in *D. melanogaster* (Fig. 3). A similar trend was seen for Tajima's D of synonymous sites (Supplemental Fig. S1): Consistent with no reduced recombination rate, Tajima's D remains unaffected by proximity to the centromere for most of the *D. mauritiana* chromosomes. This shift toward more negative Tajima's D values in low-recombining regions of *D. melanogaster* is consistent with selection at linked sites affecting neutral variability, either due to recurrent sweeps of favorable mutations (hitchhiking) (Maynard Smith and Haigh 1974) or, possibly, due to background selection, caused by the removal of linked deleterious mutations (Charlesworth et al. 1993).

Because low recombination rates will decrease the efficacy of selection, we compared the ratio of nonsynonymous to synonymous polymorphisms along the chromosomes of both species (as in Presgraves 2005; Betancourt et al. 2009). In *D. melanogaster*, the number of nonsynonymous substitutions relative to synonymous ones increases with the decrease in recombination rate toward the centromere (Presgraves 2005). In *D. mauritiana*, however, almost no effect can be noticed (Fig. 4).

The effects of elevated recombination rates in *D. mauritiana* are further apparent from the patterns of codon usage (for details, see Supplemental Results).

Signatures of positive selection in *D. mauritiana*

The neutral theory predicts a correlation between polymorphism and divergence. The McDonald–Kreitman test builds on this prediction and compares the ratio of synonymous and nonsynonymous polymorphism to the ratio of synonymous and nonsynonymous divergence; under neutrality, these quantities will be equal (McDonald and Kreitman 1991). Using a polarized McDonald–Kreitman test, we surveyed polymorphism and divergence (from *D. melanogaster* and *Drosophila yakuba*) for 10,217 genes in *D. mauritiana*.

We found 43 genes ($\text{FDR} \leq 0.05$) that deviated significantly from the neutral expectation in the polarized test of *D. mauritiana* with *D. melanogaster* as reference and *D. yakuba* as outgroup (Supplemental Table S4). A detailed list of significant genes, including those identified by unpolarized versions of the McDonald–Kreitman test, are shown in Supplemental Tables S5–S7.

While several of these genes overlapped with previous studies (for further details, see Supplemental Results), we made three particularly interesting observations in *D. mauritiana*.

First, we find strong evidence for positive selection for a gene that has been proposed to cause morphological divergence (number of sex comb teeth) between the two sister species

Figure 3. Tajima's D along the major chromosomal arms in *D. mauritiana* (red line) and *D. melanogaster* (gray line). The sliding window analysis was performed using 500-kb windows with a step size of 100 kb; chromosomal coordinates have been adjusted to *D. melanogaster*.

D. simulans and *D. mauritiana*. Graze et al. (2007) identified *CD98hc* by tissue-specific gene expression differences between both species but did not characterize its molecular evolution. Our analysis suggests that not only regulatory changes but also structural variation (Hoekstra and Coyne 2007) contribute to morphological divergence.

Second, we did not detect an accelerated rate of evolution for olfactory and gustatory receptor (*Or* and *Gr*) or Accessory gland protein (*Acp*) genes, some of which have been found to evolve rapidly in other *Drosophila* species (Begun and Lindfors 2005; Guo and Kim 2007). While gene families are more likely to be excluded from de novo assemblies, we think that species-specific selection patterns are the better explanation (for further details, see the Supplemental Discussion).

Third, the overrepresentation of nucleoporin genes among the selected genes suggests that genomic conflict may be one major driver of adaptive evolution in *D. mauritiana*. The most significant Gene Ontology (GO) term for genes significant at FDR 0.05 (43 genes) in the polarized MK test is "SMAD protein import into the nucleus" (Supplemental Table S8; for the GO enrichment analysis of the unpolarized McDonald–Kreitman tests, see Supplemental Tables S9, S10). This pointed to nucleoporin genes, some of which have been previously described to be rapidly evolving and to cause hybrid incompatibility (Presgraves 2003; Presgraves and Stephan 2007; Tang and Presgraves 2009).

Further manual inspection of the top candidates from the polarized test identified three nucleoporin genes. The gene asso-

ciated with the most significant MK test, CG8771, is a homolog of the human nucleoporin gene *Nup188*, which is involved in controlling membrane protein traffic and maintenance of nuclear membrane homeostasis (Theerthagiri et al. 2010). The yeast homolog *Nup188* plays a role in structural organization of the nuclear pore (Nehrbass et al. 1996; Miller et al. 2000). As expected for a member of a nucleopore complex, CG8771 interacts with several other nucleopore proteins (as indicated in the protein–protein interaction database STRING 9.0) (Jensen et al. 2009). Interestingly, two of the interacting partners, *Nup107* and CG11943, a homolog of the human *NUP205* gene, were also found among the top 43 candidates. One of them, CG11943, has been previously described as a rapidly evolving gene in a comparison of *D. simulans* and *D. melanogaster* (Jagadeeshan and Singh 2005). Both *Nup107*, part of the *Nup107–160* complex (Vasu and Forbes 2001), and CG11943, a member of the *Nup53–93* complex (Chen and Xu 2010), appear not only to interact with CG8771 but also with each other (Jensen et al. 2009) (but see Theerthagiri et al. [2010] for evidence against an interaction between human *NUP188* and *NUP205*). Since several Nups have not yet been identified as nucleoporins in the *D. melanogaster* annotation r5.32 and are thus missing in the GO databases (e.g., CG8771), the GO term analysis does not adequately address whether or not Nups are overrepresented among our candidate genes. We thus tested further for an overrepresentation of nucleoporins among our candidate genes by assuming that about 30 nucleoporins exist in the *D. mauritiana* genome (Wente and Rout 2010) and find that Nups are

Figure 4. Ratio of nonsynonymous and synonymous nucleotide diversity (π_N/π_S) along the major chromosomal arms in *D. mauritiana* (red line) and *D. melanogaster* (gray line). The sliding window analysis was performed using 50 genes per window and a step size of one gene. For *D. mauritiana*, genes from all four gene sets were included and matched with the orthologous gene in *D. melanogaster*. Genes with a π_N/π_S ratio >3 were excluded from both data sets.

highly significantly overrepresented (two-tailed $P < 0.0001$, χ^2 -test with Yates correction).

Given this overrepresentation of Nups, we searched for further evidence of positive selection operating on additional Nups by relaxing our search criteria. *Nup154* is significant in the unpolarized test with *D. melanogaster* at FDR 0.1 and at FDR 0.001 with *D. yakuba* as outgroup. *Nup154* is a conserved nucleoporin essential for viability (Kiger et al. 1999) and crucial for normal oogenesis and spermatogenesis (Gigliotti et al. 1998; Colozza et al. 2011), and interacts with CG8771 and CG11943 (Jensen et al. 2009), homologs of the human *NUP188* and *NUP205* genes, respectively. *Nup160*, a hybrid lethality gene between *D. simulans* and *D. melanogaster* (Tang and Presgraves 2009), is significant at FDR 0.01 (FDR 0.001) in the unpolarized test with *D. melanogaster* (*D. yakuba*) as reference, and *Nup133* at FDR 0.1 in the polarized test (Supplemental Tables S5–S7).

Polarized tests based on *D. melanogaster* as reference are not suited to determine whether positive selection predates the split of *D. simulans* and *D. mauritiana* or is still ongoing after the split of the two species. Given the strong evidence for positive selection operating on Nups, we reasoned that we should have enough power to identify ongoing positive selection after the species split and repeated the polarized tests using *D. simulans* as reference and *D. melanogaster* as outgroup. We analyzed the three candidate Nups CG8771, CG11943, and *Nup107*, their interaction partners as listed in the STRING interaction database v. 9.0 (Jensen et al. 2009), as well as *Nup160*, and found strong evidence for ongoing positive selection after the split of *D. mauritiana* and *D. simulans* for several

Nups (Table 3). Analyzing Pool-seq data from African *D. simulans* (V Nolte and C Schlötterer, unpubl.) with *D. mauritiana* as reference, we also find evidence for ongoing positive selection in *D. simulans*. The X-linked gene, CG11943, a homolog of the human *NUP205* gene, shows one of the strongest signatures of recent positive selection in both species. Overall, the evidence for positive selection was stronger in *D. simulans* than in *D. mauritiana*: Most nucleoporins with signatures of ongoing rapid evolution in *D. mauritiana* show an even more significant test result in *D. simulans*, and two nucleoporins (*Nup133* and *Nup153*) appear to evolve rapidly in *D. simulans* only, but not in *D. mauritiana*.

Given the striking evidence for rapid evolution of Nups, which is possibly driven by intragenomic conflict, we turned our attention to RNAi genes, which are also thought to evolve rapidly due to genomic conflict (Obbard et al. 2006, 2009a,b). Only 16 out of 23 RNAi genes studied by Obbard et al. (2006, 2009a) and Kolaczowski et al. (2011) were included in the initial *D. mauritiana* annotation. Hence, we manually curated the annotation of the seven missing RNAi genes (*AGO2*, *armi*, *krimp*, *AGO3*, *mael*, *rhi*, and *squ*). Consistent with positive selection, we found two RNAi genes (*aub* and *AGO2*) to show a significant polarized MK test with *D. melanogaster* as reference and *D. yakuba* as outgroup (Supplemental Table S11). Three other genes (*armi*, *Fmr1*, and *Dcr-2*) were only significant ($P < 0.05$) when no correction for multiple testing was applied. Two genes in the *D. mauritiana* (*aub* and *dcr-2*) and one gene in the African *D. simulans* data set (*armi*) showed significant evidence for ongoing positive selection after the species split (Supplemental Table S12).

Table 3. *P*-values of polarized McDonald–Kreitman tests at candidate nucleoporin genes and some of their interacting partners using *D. mauritiana* and an African *D. simulans* sample with *D. simulans* and *D. mauritiana*, respectively, as closely related reference, and *D. melanogaster* or *D. yakuba* as outgroup

Polymorphism data	<i>D. mauritiana</i>		African <i>D. simulans</i>	
Reference species data	<i>D. melanogaster</i>	<i>D. simulans</i>	<i>D. melanogaster</i>	<i>D. mauritiana</i>
Outgroup species	<i>D. yakuba</i>	<i>D. melanogaster</i>	<i>D. yakuba</i>	<i>D. melanogaster</i>
CG8771	0.0000***	0.0339	0.0000***	0.0436
CG11943	0.0004**	0.0000***	0.0000***	0.0000***
Nup107	0.0000***	0.0202	0.0021**	0.0026**
Nup133	0.0016**	1.0000	0.0002***	0.0321
Nup153	0.0208*	0.2114	0.0350	0.0114*
Nup75	0.1042	0.0551	0.0306	0.0718
Nup154	0.0031**	0.0016*	0.0065*	0.0000***
CG6540	0.3261	0.2774	0.5820	0.2063
Nup62	0.1937	0.5211	0.2418	1.0000
Nup44A	1.0000	1.0000	1.0000	1.0000
Nup98	1.0000	0.3904	0.5204	0.0980
Nup96	0.0060*	0.5410	0.0014**	1.0000
Nup160	0.0005**	0.0557	0.0001***	0.0000***

Asterisks denote genes remaining significant after correcting for multiple testing.

(*) FDR *q*-value <0.05.

(**) FDR *q*-value <0.01.

(***) FDR *q*-value <0.001.

A polymorphism trough around two loci involved in genomic conflict

Classic selective sweeps, in which the favorable allele starts at a very low frequency and increases until it (almost) reaches fixation, cause a characteristic imprint on the polymorphism pattern in the genome (Smith and Haigh 1974; Kaplan et al. 1989). The variability in the genomic region flanking the target of selection is strongly reduced and increases gradually with distance from the selected site. The shape of such a trough depends on various parameters, such as the initial frequency of the selected allele, the selection coefficient, and the recombination rate. Figure 5 shows the partitioning of variation along the *D. mauritiana* X chromosome. Two very pronounced troughs in variability can be recognized that could not be attributed to alignment artifacts (see Supplemental Results). In both regions with reduced variability, we noticed a high differentiation (F_{ST}) (data not shown) from African *D. simulans*, but no increase in sequence divergence (D_{xy}) (Fig. 1).

The first region encompasses ~600 kb with a threefold reduction in variability relative to the average X-chromosomal diversity (mean π for the region = 0.002 vs. mean X-linked π = 0.0059; two-tailed Wilcoxon rank-sum test, nonoverlapping 10-kb windows, $P < 2.2 \times 10^{-16}$). In the central position of the trough, the variability is even further reduced (mean π = 0.001, coordinates 8.75–9.05 Mb). In addition, we observed a pronounced reduction in Tajima's *D* values compared with the remainder of the X chromosome (two-tailed Wilcoxon rank-sum test, nonoverlapping 10-kb windows, $P < 2.2 \times 10^{-16}$) (Fig. 3).

The width of the trough suggests an exceptionally strong selective sweep, since it is located in a genomic region of normal to high recombination (True et al. 1996), and no common inversion polymorphism has been described in *D. mauritiana* (for review, see Aulard et al. 2004). A close inspection of the function of the about 37 genes in the genomic region of reduced variability did not show any gene for which adaptive evolution was previously suggested. The only gene that could tentatively be associated with positive

selection is *Ser7*, since it seems to be involved in immune response (Irving et al. 2001; Hill-Burns and Clark 2009). We thus turned our attention to other possible causes of selective sweeps. In addition to beneficial alleles that provide some fitness benefit to the organism, alleles involved in genomic conflict can also have very strong selective advantages (Presgraves et al. 2009) and thus the potential to drive selective sweeps. We note that a pair of genes causing sex-ratio distortion in *D. simulans* is located within the region with the most extreme reduction in variability (Fig. 5; Supplemental Fig. S2): Alleles of the paralogous genes *Mother of Dox* (*MDox*) and *Dox* function as drivers in a well-characterized sex-ratio meiotic drive system in *D. simulans* (Tao et al. 2007a; Kingan et al. 2010). The estimated selection coefficient *s* ranges from 0.12 to 0.39, depending on the parameter values used (Supplemental Table S13), which could be consistent with strong selection during genomic conflict (Curtisinger 1984).

The second trough in variability on the X chromosome extends over an even larger region but shows a less pronounced reduction in variability. A genomic region of ~1000 kb between coordinates 16–17 Mb of the *D. mauritiana* reference genome shows an approximately twofold reduction in variability (mean π = 0.0030 vs. mean X-linked π = 0.0059, two-tailed Wilcoxon rank-sum test, nonoverlapping 10-kb windows, $P < 2.2 \times 10^{-16}$) (Fig. 5). Similar to the first sweep region, Tajima's *D* is also lower than in the remainder of the X chromosome (two-tailed Wilcoxon rank-sum test of Tajima's *D* values for 10-kb windows in each sweep region vs. 10-kb windows in the remainder of the X chromosome, $P < 2.2 \times 10^{-16}$) (Fig. 3). Estimates for the selection coefficient range from 0.04 to 0.46 (Supplemental Table S13).

The region of reduced variability contains more than 60 genes with many of them having no known function. The strongest reduction in variability is observed at the proximal border of the sweep window and harbors the haplolethal 16F gene cluster described in *D. melanogaster* (Prado et al. 1999).

Surprisingly, at the center of the window of reduced variability, we find another gene with a well-documented role in speciation and, possibly, genomic conflict, *Odyseus* (*OdsH*) (Ting et al. 1998; Bayes and Malik 2009). In addition, the sweep around the *OdsH* gene extends to the region in which the *enhancer of Dox* (*E[Dox]*) has been located, a not yet precisely mapped factor proximal to the gene *forked* that enhances the sex-ratio distorting effect of *Dox* (Fig. 5; Tao et al. 2007a).

Discussion

Quality of draft genomes based on paired-end Illumina sequencing

Here, we have built a high-quality draft genome of a *Drosophila* species using only short paired-end reads. Using a conservative, *D. melanogaster*-centric annotation, we recovered a similar number of genes as a previous genome project did for *D. simulans* and

Figure 5. Nucleotide diversity (π) along the *D. mauritiana* X chromosome. The location of genes potentially causing the two selective sweeps are indicated: (large red diamond) *MDox/Dox*; (large blue diamond) *OdsH*; (small red diamond) *E(Dox)*. Nucleotide diversity (π) is plotted in nonoverlapping 10-kb windows.

D. sechellia (Clark et al. 2007). The analyses in this study demonstrate that a draft genome facilitates addressing several important evolutionary questions. Nevertheless, we caution that it has some shortcomings. First, we are not able to provide a correct annotation of transposable elements in the *D. mauritiana* reference genome, since repetitive structures cannot be reliably assembled with short reads (Phillippy et al. 2008). Furthermore, heterochromatic sequences are poorly represented. Since the traces of genomic conflict are so apparent in *D. mauritiana*, and heterochromatic sequences may be important players in genomic conflict and speciation (Brideau et al. 2006; Cattani and Presgraves 2009; Ferree and Barbash 2009; Meiklejohn et al. 2011), we caution that the genomic signatures of ongoing genomic conflict are probably incomplete without the corresponding heterochromatic regions. Finally, gene families composed of closely related paralogs tend to be collapsed into a single copy during de novo assembly (e.g., the *Hsp70* gene cluster).

Nevertheless, our assembly recovered a large fraction of gene families that are frequently identified as targets of positive selection (Supplemental Table S14). Accessory gland protein (*Acp*) and seminal fluid protein (*Sfp*) genes, which belong to recently duplicated gene families, frequently evolve under positive selection (for review, see Ram and Wolfner 2007), but we did not find genes in these categories among the top candidates for adaptively evolving genes in the polarized McDonald–Kreitman test. Similarly, olfactory (*Or*) and gustatory receptor (*Gr*) genes are frequently involved into ecological adaptation and speciation (Guo and Kim 2007; McBride 2007; Tunstall et al. 2007; Gardiner et al. 2008, 2009), but we also failed to identify such genes among the top candidates. Since our annotation recovered a large fraction of the adaptively evolving genes described in *D. melanogaster*, we consider it unlikely that the absence of a molecular signature of adaptation in *D. mauritiana* is an annotation artifact. Rather, we speculate that the selective forces driving an adaptive response of these genes in other *Drosophila* species are less prominent in *D. mauritiana*.

Nucleoporins as a preferential target for positive selection

Despite the fact that the function and composition of nuclear pore complexes are highly conserved, recent work showed that some of their components, the nucleoporins, evolve rapidly, and

two of them cause hybrid lethality in *Drosophila* (Presgraves et al. 2003; Presgraves and Stephan 2007; Tang and Presgraves 2009). Presgraves and Stephan (2007) suggest three forms of genetic conflict that could drive the rapid evolution of nucleoporins: (1) host–parasite conflict due to viruses that need to enter via the gatekeeper nuclear pore complexes, which function to exclude invading viruses; (2) intragenomic conflict due to centromeric drive, since some Nups are associated with kinetochores; (3) intragenomic conflict due to other forms of segregation distortion, since nuclear pore complexes may potentially suppress them. Our analyses cannot distinguish between these hypotheses, but they provide additional evidence that many Nups evolve unusually rapidly due to positive selection. Furthermore, we show that the rapid evolution is not restricted to some time during the divergence between *D. melanogaster* and *D. simulans* (Presgraves et al. 2003; Presgraves and Stephan 2007; Tang and Presgraves 2009), but that positive selection is an ongoing process that continues after the split between *D. mauritiana* and *D. simulans*.

Our evolutionary analyses provide some insights into how nucleoporin genes may be involved in genomic conflict. CG11943, a homolog of the human *NUP205* gene, shows strong evidence of ongoing selection in *D. mauritiana* and *D. simulans*. Presgraves et al. (2003) and Tang and Presgraves (2009) previously identified *Nup96* and *Nup160* as one cause of hybrid lethality between *D. simulans* and *D. melanogaster*, due to an interaction with an as-yet-unidentified X-linked factor. Given that CG11943 is located on the X chromosome, we speculate that it may be an alternative interaction partner of *Nup160* and/or *Nup96* instead of (or in addition to) their suggested *Nup153*.

While the McDonald–Kreitman test with *D. simulans* as a reference indicates that the high rate of sequence evolution is ongoing in *D. mauritiana*, there was no clear signature of a selective sweep at nucleoporin genes in the polymorphism data (Fig. 6). Since this observation is consistent with the analyses of Presgraves and Stephan (2007) and Tang and Presgraves (2009), we hypothesize that positively selected mutations in interacting proteins may lead to complex sweep dynamics, which could retard the spread of a beneficial mutation. As a consequence, beneficial mutations at Nups may result in a signature that resembles more a soft sweep (Hermisson and Pennings 2005) rather than a hard sweep.

Figure 6. Nucleotide diversity (π) and Tajima's D at selected nucleoporin genes in *D. mauritiana*. (A) Nucleotide diversity (π) at selected nucleoporin genes in comparison to the average chromosome-wide nucleotide diversity (π) of the corresponding chromosome (dashed line). (B) Tajima's D at selected nucleoporin genes in comparison to the average chromosome-wide Tajima's D of the corresponding chromosome (dashed line).

Two recent selective sweeps potentially associated with genomic conflict

One of our most striking findings is that the *D. mauritiana* genome harbors two large regions—0.6 and 1 Mb—of profoundly reduced diversity, suggesting that two exceptionally strong selective sweeps have occurred in this species. In comparison, genomic signatures of previously described sweeps in *Drosophila* are much narrower with estimated selection coefficients being about an order of magnitude lower: The broadest valley of reduced variability described to date is caused by the insertion of a transposable element next to the gene *Cyp6g1*, which confers resistance to DDT—the sweep extends over ~100 kb and has an associated selection coefficient of 0.022 (Schlenke and Begun 2004).

The gene that could most parsimoniously be assumed to drive one of the sweeps on the *D. mauritiana* X chromosome is *OdsH*. This gene was initially described as a “speciation gene” that causes hybrid male sterility between *D. mauritiana* and *D. simulans* and shows a strongly accelerated rate of evolution in *D. mauritiana* (Ting et al. 1998). Later it was recognized that the *OdsH* gene prod-

uct interacts with Y-linked heterochromatin in hybrids between *D. mauritiana* and *D. simulans* but not in pure species, suggesting that it could be involved in genomic conflict (Bayes and Malik 2009; Meiklejohn et al. 2011). Consistent with previous reports, we observe a much higher number of nonsynonymous fixations along the *D. mauritiana* than the *D. simulans* lineage, but in both species neither the homeodomain nor the entire gene showed evidence of positive selection in a polarized McDonald–Kreitman test.

The second selective sweep might be caused by the Winters meiotic drive system, which is well-characterized in *D. simulans* (Tao et al. 2007a,b; Kingan et al. 2010). This system consists of at least three components: the drivers *MDox* and *Dox*, the autosomal dominant suppressor *Nmy*, and the enhancer of *Dox* [*E(Dox)*]. The sequences of these genes are highly similar, partially derived from one another by tandem duplication and retrotransposition, and contain tandem repeat structures (Tao et al. 2007a,b). As a result of the sequence similarity and repetitive structure of these loci, reliable sequence analysis of this region is difficult, even with targeted PCR approaches (Tao et al. 2007a,b), and essentially impossible with genome-wide short read sequencing.

Our *D. mauritiana* strains did not show obvious signs of sex-ratio distortion, but theoretical models predict that ongoing intragenomic conflict results in rapid cycles during which competing alleles rise and fall in frequency (Charlesworth and Hartl 1978; Carvalho and Vaz 1999; Hall 2004). Driver alleles, such as sex-ratio distorters, will increase in frequency until a suppressor allele ar-

rives, which spreads, and the genomic conflict ultimately disappears (i.e., the population reaches a balanced sex ratio). While an almost complete sweep of a strongly distorting allele appears unlikely, theoretical models have described situations under which such a pattern is predicted (Charlesworth and Hartl 1978; Carvalho and Vaz 1999; Hall 2004).

In some *D. simulans* populations, the *Dox* gene shows evidence of a partial selective sweep (Kingan et al. 2010), but it is difficult to distinguish highly localized selective sweeps from random fluctuations in variability due to the bottleneck associated with the out of Africa expansion (Jensen et al. 2005). We further scrutinized the genomic region around *Dox* using Pool-seq data from African *D. simulans* (V Nolte and C Schlötterer, unpubl.) and did not note any pronounced trough in variability around the *Dox* region, suggesting that, at least in the African population sample, no evidence for a selective sweep comparable to the one in *D. mauritiana* could be detected (Supplemental Fig. S3).

The *Dox* system was initially discovered in *D. simulans* (Dermitzakis et al. 2000), and the driver loci *MDox* and *Dox* have not yet been functionally analyzed in *D. mauritiana*. While an

analysis of the *D. mauritiana* alleles present at *MDox*, *Dox*, and *Nmy* is not possible from the Pool-seq data (see above), Tao et al. (2007b) inferred from sequence comparison that a functional suppressor allele at *Nmy* is present in *D. mauritiana*, suggesting the existence of a functional distorter (Tao et al. 2007a,b). No selective sweep could be detected in the *Nmy* gene region (data not shown), suggesting that no new allele at *Nmy* has swept through *D. mauritiana*.

Given the dynamic evolution of repetitive structures in the Winters sex-ratio genes, it is possible that in *D. mauritiana*, a new driver allele at *Dox* evolved and caused the pronounced sweep signature, while in *D. simulans*, the signature of an older sweep has already been erased. Alternatively, we could speculate that it may be easier for an allele to sweep in *D. mauritiana* since *D. mauritiana* has almost no population structure (Nunes et al. 2010), while the cosmopolitan species *D. simulans* shows a higher level of population differentiation (Hamblin and Veuille 1999).

Theoretical studies predict that a beneficial allele will spread much faster in panmictic populations, whereas population subdivision and low migration rates lead to a delay in the fixation of a beneficial mutation (Barton 2000; Santiago and Caballero 2005; Kim and Maruki 2011).

The importance of population structure for the detection of meiotic drive dynamics has also been highlighted in recent theoretical work. Hall (2004) suggested that the hitherto absence of documented cycling behavior in natural *Drosophila* populations may be the result of migration between subdivided populations with different drive parameters. Instead, in isolated populations that share the same drive dynamics due to panmixia large fluctuations in driver and suppressor can be seen. Since *D. mauritiana* shows no population differentiation on Mauritius, we think that this could explain the difference with the other *Drosophila* species.

It is apparent that more work is needed to characterize the driver, responder, and suppressor alleles in both species to shed further light onto the differences in evolutionary signatures observed between the two species.

Methods

D. mauritiana strains and Illumina sequencing

We used the *D. mauritiana* isofemale strain MS17 (<http://kyotofly.kit.ac.jp/cgi-bin/ehime/index.cgi>, stock number E-18912) to generate a *D. mauritiana* reference genome. Pool-seq data were obtained from 107 *D. mauritiana* lines collected at different time points and locations in Mauritius (Supplemental Table S2). Illumina libraries were generated following the instructions of the Illumina Paired-End Sample Preparation Kit and sequenced on a GAIIx.

De novo assembly and annotation of a *D. mauritiana* reference genome

To generate the *D. mauritiana* reference genome sequence, we initially performed a de novo assembly of Illumina reads using the software CLC Assembly Cell v. 3.1.0 beta2 (CLC Bio). In the second phase of the assembly procedure, we anchored de novo contigs on the reference genome of *D. melanogaster* r. 5.22 using the nucmer module in the MUMmer package v. 3.0 (Kurtz et al. 2004). The *D. mauritiana* chromosomes were built by overlapping or concatenating contigs. The longest isoform of each *D. melanogaster* protein from FlyBase release 5.32 served as template for annotation. Each protein sequence was aligned to the *D. mauritiana*

reference genome using exonerate v. 2.0 (Slater and Birney 2005). We generated four sets of gene annotations, using varying degrees of filtering criteria that are described in detail in Supplemental Methods.

Divergence estimates and codon usage analysis

We performed multiple alignments of the *D. mauritiana*, a *D. simulans*, the *D. sechellia* r.1.3, and the *D. melanogaster* r. 5.32 genome sequences using MAUVE (Darling et al. 2010) and calculated pairwise divergence *Dxy* between them using the PoPoolation package (Kofler et al. 2011). We used CALcal v. 1.4 (Puigbo et al. 2008) to determine the Codon Adaptation Index (CAI), originally developed by Sharp and Li (1987).

Reference mapping and variability estimates in *D. mauritiana* Pool-seq data

Paired-end reads of the pooled *D. mauritiana* sample were aligned to the MS17 draft genome (or *D. melanogaster* genome) using bwa v. 0.5.8 (Li and Durbin 2009). Alignments were filtered for a minimum mapping quality of 20 and for properly paired reads using SAMtools v. 0.1.9 (<http://samtools.sourceforge.net/>). Minimum requirements for coverage and allele count used in SNP calling are detailed in Supplemental Methods. Analyses of π and Tajima's *D* were performed with the PoPoolation package (Kofler et al. 2011). To test for recurrent positive selection in the *D. mauritiana* lineage, we performed McDonald–Kreitman tests. Multiple alignments of the coding sequence of each *D. mauritiana* gene with the orthologs of *D. melanogaster* r. 5.32 and *D. yakuba* r. 1.3 were generated using PRANK v. 100701 (Loytynoja and Goldman 2005). We combined the interspecific with the intraspecific alignments using custom Perl scripts and performed McDonald–Kreitman tests using the MK.pl script obtained from <http://www.dpgp.org/aholloway/Software.html> (Holloway et al. 2007). We calculated false discovery rates (FDR) using the LBE package (Dalmasso et al. 2005) and performed an analysis of Gene Ontology enrichment with GOrilla (<http://cbl-gorilla.cs.technion.ac.il/>) (Eden et al. 2009). Details of all analyses are provided in Supplemental Methods.

Statistical tests

Statistical tests were performed using R version 2.11.1 (The R Core Team 2010) unless stated otherwise.

Data access

All Illumina short reads used in this study are available from the NCBI Sequence Read Archive (SRA) (<http://www.ncbi.nlm.nih.gov/sra>) under the following accession numbers: the single *D. mauritiana* reference strain MS17 under SRA058420, the *D. mauritiana* Pool-seq data under SRA058664, the *D. simulans* reference strain Kib32 under SRA059282, and the African *D. simulans* Pool-seq data under SRA059292. The *D. mauritiana* strain MS17 reference genome and annotation are available at http://www.popoolation.at/mauritiana_genome/index.html. A BAM file containing *D. mauritiana* Pool-seq data is available at http://www.popoolation.at/mauritiana_genome/index.html. A searchable, user-friendly version of the *D. mauritiana* Pool-seq and the African *D. simulans* Pool-seq data is available at http://www.popoolation.at/pgt/dmau_browse.html and http://www.popoolation.at/pgt/dsim_browse.html.

Acknowledgments

We thank all members of the Institute of Population Genetics for discussion and support throughout the entire project. We are grateful to A.J. Betancourt for drawing our attention to the *MDox/Dox* locus and to N. de Maio for mathematical support. A.J. Betancourt, B.F. McAllister, C. Willett, R. Tobler, and three anonymous reviewers provided helpful comments on earlier versions of the manuscript. We are grateful to the Ehime-Fly *Drosophila* Stock center for providing us with fly stocks. This work was supported by the Austrian Science Funds (FWF) grant P19467 to C.S.

Author contributions: V.N. and C.S. conceived and designed the experiments; V.N. performed the experiments; V.N. analyzed the data; R.V.P. and R.K. contributed analysis tools; and V.N. and C.S. wrote the manuscript.

References

- Aulard S, Monti L, Chaminade N, Lemeunier F. 2004. Mitotic and polytene chromosomes: Comparisons between *Drosophila melanogaster* and *Drosophila simulans*. *Genetica* **120**: 137–150.
- Ballard JW. 2000. When one is not enough: Introgression of mitochondrial DNA in *Drosophila*. *Mol Biol Evol* **17**: 1126–1130.
- Barbash DA. 2007. *Nup96*-dependent hybrid lethality occurs in a subset of species from the *simulans* clade of *Drosophila*. *Genetics* **176**: 543–552.
- Barton NH. 2000. Genetic hitchhiking. *Philos Trans R Soc Lond B Biol Sci* **355**: 1553–1562.
- Bastide H, Cazemajor M, Ogereau D, Derome N, Hospital F, Montchamp-Moreau C. 2011. Rapid rise and fall of selfish sex-ratio X chromosomes in *Drosophila simulans*: Spatio-temporal analysis of phenotypic and molecular data. *Mol Biol Evol* **28**: 2461–2470.
- Bayes JJ, Malik HS. 2009. Altered heterochromatin binding by a hybrid sterility protein in *Drosophila* sibling species. *Science* **326**: 1538–1541.
- Begun DJ, Aquadro CF. 1992. Levels of naturally occurring DNA polymorphism correlate with recombination rates in *D. melanogaster*. *Nature* **356**: 519–520.
- Begun DJ, Lindfors HA. 2005. Rapid evolution of genomic *Acp* complement in the *melanogaster* subgroup of *Drosophila*. *Mol Biol Evol* **22**: 2010–2021.
- Begun DJ, Holloway AK, Stevens K, Hillier LW, Poh YP, Hahn MW, Nista PM, Jones CD, Kern AD, Dewey CN, et al. 2007. Population genomics: Whole-genome analysis of polymorphism and divergence in *Drosophila simulans*. *PLoS Biol* **5**: e310. doi: 10.1371/journal.pbio.0050310.
- Betancourt AJ, Welch JJ, Charlesworth B. 2009. Reduced effectiveness of selection caused by a lack of recombination. *Curr Biol* **19**: 655–660.
- Brideau NJ, Flores HA, Wang J, Maheshwari S, Wang X, Barbash DA. 2006. Two Dobzhansky-Muller genes interact to cause hybrid lethality in *Drosophila*. *Science* **314**: 1292–1295.
- Carvalho AB, Vaz SC. 1999. Are *Drosophila* SR drive chromosomes always balanced? *Heredity (Edinb)* **83**: 221–228.
- Cattani MV, Presgraves DC. 2009. Genetics and lineage-specific evolution of a lethal hybrid incompatibility between *Drosophila mauritiana* and its sibling species. *Genetics* **181**: 1545–1555.
- Charlesworth B, Hartl DL. 1978. Population dynamics of the segregation distorter polymorphism of *Drosophila melanogaster*. *Genetics* **89**: 171–192.
- Charlesworth B, Coyne JA, Barton NH. 1987. The relative rates of evolution of sex chromosomes and autosomes. *Am Nat* **130**: 113–146.
- Charlesworth B, Morgan MT, Charlesworth D. 1993. The effect of deleterious mutations on neutral molecular variation. *Genetics* **134**: 1289–1303.
- Chen X, Xu L. 2010. Specific nucleoporin requirement for Smad nuclear translocation. *Mol Cell Biol* **30**: 4022–4034.
- Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis M, Gelbart W, Iyer VN, et al. 2007. Evolution of genes and genomes on the *Drosophila* phylogeny. *Nature* **450**: 203–218.
- Colozza G, Montembault E, Quenerch' du E, Riparbelli MG, D'Avino PP, Callaini G. 2011. *Drosophila* nucleoporin *Nup154* controls cell viability, proliferation and nuclear accumulation of Mad transcription factor. *Tissue Cell* **43**: 254–261.
- Curtis JW. 1984. Components of selection in X chromosome lines of *Drosophila melanogaster*: Sex ratio modification by meiotic drive and viability selection. *Genetics* **108**: 941–952.
- Curtis JW, Feldman MW. 1980. Experimental and theoretical analysis of the “sex-ratio” polymorphism in *Drosophila pseudoobscura*. *Genetics* **94**: 445–466.
- Dalmaso C, Broet P, Moreau T. 2005. A simple procedure for estimating the false discovery rate. *Bioinformatics* **21**: 660–668.
- Darling AE, Mau B, Perna NT. 2010. progressiveMauve: Multiple genome alignment with gene gain, loss, and rearrangement. *PLoS ONE* **5**: e11147. doi: 10.1371/journal.pone.0011147.
- Dermitzakis ET, Masly JP, Waldrip HM, Clark AG. 2000. Non-Mendelian segregation of sex chromosomes in heterospecific *Drosophila* males. *Genetics* **154**: 687–694.
- Derome N, Metayer K, Montchamp-Moreau C, Veuille M. 2004. Signature of selective sweep associated with the evolution of sex-ratio drive in *Drosophila simulans*. *Genetics* **166**: 1357–1366.
- Derome N, Baudry E, Ogereau D, Veuille M, Montchamp-Moreau C. 2008. Selective sweeps in a 2-locus model for sex-ratio meiotic drive in *Drosophila simulans*. *Mol Biol Evol* **25**: 409–416.
- Eden E, Navon R, Steinfeld I, Lipson D, Yakhini Z. 2009. *GOrrilla*: A tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* **10**: 48. doi: 10.1186/1471-2105-10-48.
- Ferree PM, Barbash DA. 2009. Species-specific heterochromatin prevents mitotic chromosome segregation to cause hybrid lethality in *Drosophila*. *PLoS Biol* **7**: e1000234. doi: 10.1371/journal.pbio.1000234.
- Gardiner A, Barker D, Butlin RK, Jordan WC, Ritchie MG. 2008. *Drosophila* chemoreceptor gene evolution: Selection, specialization and genome size. *Mol Ecol* **17**: 1648–1657.
- Gardiner A, Butlin RK, Jordan WC, Ritchie MG. 2009. Sites of evolutionary divergence differ between olfactory and gustatory receptors of *Drosophila*. *Biol Lett* **5**: 244–247.
- Garrigan D, Kingan SB, Geneva AJ, Andolfatto P, Clark AG, Thornton KR, Presgraves DC. 2012. Genome sequencing reveals complex speciation in the *Drosophila simulans* clade. *Genome Res* **22**: 1499–1511.
- Gershenson S. 1928. A new sex-ratio abnormality in *Drosophila obscura*. *Genetics* **13**: 488–504.
- Gigliotti S, Callaini G, Andone S, Riparbelli MG, Pernas-Alonso R, Hoffmann G, Graziani F, Malva C. 1998. *Nup154*, a new *Drosophila* gene essential for male and female gametogenesis is related to the *Nup155* vertebrate nucleoporin gene. *J Cell Biol* **142**: 1195–1207.
- Graze RM, Barmina O, Tufts D, Naderi E, Harmon KL, Persianinova M, Nuzhdin SV. 2007. New candidate genes for sex-comb divergence between *Drosophila mauritiana* and *Drosophila simulans*. *Genetics* **176**: 2561–2576.
- Guo S, Kim J. 2007. Molecular evolution of *Drosophila* odorant receptor genes. *Mol Biol Evol* **24**: 1198–1207.
- Hall DW. 2004. Meiotic drive and sex chromosome cycling. *Evolution* **58**: 925–931.
- Hamblin MT, Veuille M. 1999. Population structure among African and derived populations of *Drosophila simulans*: Evidence for ancient subdivision and recent admixture. *Genetics* **153**: 305–317.
- Hamilton WD. 1967. Extraordinary sex ratios. *Science* **156**: 477–488.
- Hermisson J, Pennings PS. 2005. Soft sweeps: Molecular population genetics of adaptation from standing genetic variation. *Genetics* **169**: 2335–2352.
- Hey J, Kliman RM. 1993. Population genetics and phylogenetics of DNA sequence variation at multiple loci within the *Drosophila melanogaster* species complex. *Mol Biol Evol* **10**: 804–822.
- Hill-Burns EM, Clark AG. 2009. X-linked variation in immune response in *Drosophila melanogaster*. *Genetics* **183**: 1477–1491.
- Hoekstra HE, Coyne JA. 2007. The locus of evolution: evo devo and the genetics of adaptation. *Evolution* **61**: 995–1016.
- Holloway AK, Lawniczak MK, Mezey JG, Begun DJ, Jones CD. 2007. Adaptive gene expression divergence inferred from population genomics. *PLoS Genet* **3**: 2007–2013.
- Hudson RR. 1994. How can the low levels of DNA sequence variation in regions of the *Drosophila* genome with low recombination rates be explained? *Proc Natl Acad Sci* **91**: 6815–6818.
- Irving P, Troxler L, Heuer TS, Belvin M, Kopczynski C, Reichhart JM, Hoffmann JA, Hetru C. 2001. A genome-wide analysis of immune responses in *Drosophila*. *Proc Natl Acad Sci* **98**: 15119–15124.
- Jagadeeshan S, Singh RS. 2005. Rapidly evolving genes of *Drosophila*: Differing levels of selective pressure in testis, ovary, and head tissues between sibling species. *Mol Biol Evol* **22**: 1793–1801.
- Jensen JD, Kim Y, Dumont VB, Aquadro CF, Bustamante CD. 2005. Distinguishing between selective sweeps and demography using DNA polymorphism data. *Genetics* **170**: 1401–1410.
- Jensen LJ, Kuhn M, Stark M, Chaffron S, Creevey C, Muller J, Doerks T, Julien P, Roth A, Simonovic M, et al. 2009. STRING 8—a global view on proteins and their functional interactions in 630 organisms. *Nucleic Acids Res* **37**: D412–D416.
- Kaplan NL, Hudson RR, Langley CH. 1989. The “hitchhiking effect” revisited. *Genetics* **123**: 887–899.
- Kiger AA, Gigliotti S, Fuller MT. 1999. Developmental genetics of the essential *Drosophila* nucleoporin *Nup154*: Allelic differences due to an outward-directed promoter in the P-element 3' end. *Genetics* **153**: 799–812.

- Kim Y, Maruki T. 2011. Hitchhiking effect of a beneficial mutation spreading in a subdivided population. *Genetics* **189**: 213–226.
- Kingan SB, Garrigan D, Hartl DL. 2010. Recurrent selection on the Winters sex-ratio genes in *Drosophila simulans*. *Genetics* **184**: 253–265.
- Kofler R, Orozco-terWengel P, De Maio N, Pandey RV, Nolte V, Futschik A, Kosiol C, Schlotterer C. 2011. PoPoolation: A toolbox for population genetic analysis of next generation sequencing data from pooled individuals. *PLoS ONE* **6**: e15925. doi: 10.1371/journal.pone.0015925.
- Kolaczowski B, Hupalo DN, Kern AD. 2011. Recurrent adaptation in RNA interference genes across the *Drosophila* phylogeny. *Mol Biol Evol* **28**: 1033–1042.
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol* **5**: R12. doi: 10.1186/gb-2004-5-2-r12.
- Langley CH, Stevens K, Cardeno C, Lee YC, Schrider DR, Pool JE, Langley SA, Suarez C, Corbett-Detig RB, Kolaczowski B, et al. 2012. Genomic variation in natural populations of *Drosophila melanogaster*. *Genetics* **192**: 533–598.
- Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**: 1754–1760.
- Loytynoja A, Goldman N. 2005. An algorithm for progressive multiple alignment of sequences with insertions. *Proc Natl Acad Sci* **102**: 10557–10562.
- Lyttle TW. 1977. Experimental population genetics of meiotic drive systems I. Pseudo-Y chromosomal drive as a means of eliminating cage populations of *Drosophila melanogaster*. *Genetics* **86**: 413–445.
- Maheshwari S, Barbash DA. 2012. Cis-by-trans regulatory divergence causes the asymmetric lethal effects of an ancestral hybrid incompatibility gene. *PLoS Genet* **8**: e1002597. doi: 10.1371/journal.pgen.1002597.
- Maynard Smith J, Haigh J. 1974. The hitch-hiking effect of a favorable gene. *Genet Res* **23**: 23–35.
- McBride CS. 2007. Rapid evolution of smell and taste receptor genes during host specialization in *Drosophila sechellia*. *Proc Natl Acad Sci* **104**: 4996–5001.
- McDonald JH, Kreitman M. 1991. Adaptive protein evolution at the *Adh* locus in *Drosophila*. *Nature* **351**: 652–654.
- Meiklejohn CD, Landeen EL, Cook JM, Kingan SB, Presgraves DC. 2011. Sex chromosome-specific regulation in the *Drosophila* male germline but little evidence for chromosomal dosage compensation or meiotic inactivation. *PLoS Biol* **9**: e1001126. doi: 10.1371/journal.pbio.1001126.
- Mercot H, Atlan A, Jacques M, Montchamp-Moreau C. 1995. Sex-ratio distortion in *Drosophila simulans*: Co-occurrence of a meiotic drive and a suppressor of drive. *J Evol Biol* **8**: 283–300.
- Miller BR, Powers M, Park M, Fischer W, Forbes DJ. 2000. Identification of a new vertebrate nucleoporin, *Nup188*, with the use of a novel organelle trap assay. *Mol Biol Cell* **11**: 3381–3396.
- Moriyama EN, Powell JR. 1996. Intraspecific nuclear DNA variation in *Drosophila*. *Mol Biol Evol* **13**: 261–277.
- Morton RA, Choudhary M, Cariou ML, Singh RS. 2004. A reanalysis of protein polymorphism in *Drosophila melanogaster*, *D. simulans*, *D. sechellia* and *D. mauritiana*: Effects of population size and selection. *Genetica* **120**: 101–114.
- Nehrass U, Rout MP, Maguire S, Blobel G, Wozniak RW. 1996. The yeast nucleoporin *Nup188p* interacts genetically and physically with the core structures of the nuclear pore complex. *J Cell Biol* **133**: 1153–1162.
- Nunes MD, Wengel PO, Kreissl M, Schlotterer C. 2010. Multiple hybridization events between *Drosophila simulans* and *Drosophila mauritiana* are supported by mtDNA introgression. *Mol Ecol* **19**: 4695–4707.
- Obbard DJ, Jiggins FM, Halligan DL, Little TJ. 2006. Natural selection drives extremely rapid evolution in antiviral RNAi genes. *Curr Biol* **16**: 580–585.
- Obbard DJ, Gordon KH, Buck AH, Jiggins FM. 2009a. The evolution of RNAi as a defence against viruses and transposable elements. *Philos Trans R Soc Lond B Biol Sci* **364**: 99–115.
- Obbard DJ, Welch JJ, Kim KW, Jiggins FM. 2009b. Quantifying adaptive evolution in the *Drosophila* immune system. *PLoS Genet* **5**: e1000698. doi: 10.1371/journal.pgen.1000698.
- Orr HA, Betancourt AJ. 2001. Haldane's sieve and adaptation from the standing genetic variation. *Genetics* **157**: 875–884.
- Perez DE, Wu CI. 1995. Further characterization of the *Odysseus* locus of hybrid sterility in *Drosophila*: One gene is not enough. *Genetics* **140**: 201–206.
- Perez DE, Wu CI, Johnson NA, Wu ML. 1993. Genetics of reproductive isolation in the *Drosophila simulans* clade: DNA marker-assisted mapping and characterization of a hybrid-male sterility gene, *Odysseus* (*Ods*). *Genetics* **134**: 261–275.
- Phadnis N, Orr HA. 2009. A single gene causes both male sterility and segregation distortion in *Drosophila* hybrids. *Science* **323**: 376–379.
- Phillippy AM, Schatz MC, Pop M. 2008. Genome assembly forensics: Finding the elusive mis-assembly. *Genome Biol* **9**: R55. doi: 10.1186/gb-2008-9-3-r55.
- Prado A, Canal I, Ferrus A. 1999. The haplolethal region at the 16F gene cluster of *Drosophila melanogaster*: Structure and function. *Genetics* **151**: 163–175.
- Presgraves DC. 2003. A fine-scale genetic analysis of hybrid incompatibilities in *Drosophila*. *Genetics* **163**: 955–972.
- Presgraves DC. 2005. Recombination enhances protein adaptation in *Drosophila melanogaster*. *Curr Biol* **15**: 1651–1656.
- Presgraves DC. 2007. Does genetic conflict drive rapid molecular evolution of nuclear transport genes in *Drosophila*? *Bioessays* **29**: 386–391.
- Presgraves DC, Stephan W. 2007. Pervasive adaptive evolution among interactors of the *Drosophila* hybrid inviability gene, *Nup96*. *Mol Biol Evol* **24**: 306–314.
- Presgraves DC, Balagopalan L, Abmayr SM, Orr HA. 2003. Adaptive evolution drives divergence of a hybrid inviability gene between two species of *Drosophila*. *Nature* **423**: 715–719.
- Presgraves DC, Gerard PR, Cherukuri A, Lyttle TW. 2009. Large-scale selective sweep among Segregation Distorter chromosomes in African populations of *Drosophila melanogaster*. *PLoS Genet* **5**: e1000463. doi: 10.1371/journal.pgen.1000463.
- Puigbo P, Bravo IG, Garcia-Vallve S. 2008. CAIcal: A combined set of tools to assess codon usage adaptation. *Biol Direct* **3**: 38. doi: 10.1186/1745-6150-3-38.
- The R Core Team. 2010. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna.
- Ram KR, Wolfner MF. 2007. Seminal influences: *Drosophila* Acps and the molecular interplay between males and females during reproduction. *Integr Comp Biol* **47**: 427–445.
- Sandler L, Hiraizumi Y, Sandler I. 1959. Meiotic drive in natural populations of *Drosophila melanogaster*. I. The cytogenetic basis of segregation-distortion. *Genetics* **44**: 233–250.
- Santiago E, Caballero A. 2005. Variation after a selective sweep in a subdivided population. *Genetics* **169**: 475–483.
- Schlenke TA, Begun DJ. 2004. Strong selective sweep associated with a transposon insertion in *Drosophila simulans*. *Proc Natl Acad Sci* **101**: 1626–1631.
- Sharp PM, Li WH. 1987. The codon Adaptation Index—a measure of directional synonymous codon usage bias, and its potential applications. *Nucleic Acids Res* **15**: 1281–1295.
- Singh ND, Larracuente AM, Clark AG. 2008. Contrasting the efficacy of selection on the X and autosomes in *Drosophila*. *Mol Biol Evol* **25**: 454–467.
- Slater GS, Birney E. 2005. Automated generation of heuristics for biological sequence comparison. *BMC Bioinformatics* **6**: 31. doi: 10.1186/1471-2105-6-31.
- Smith JM, Haigh J. 1974. The hitch-hiking effect of a favourable gene. *Genet Res* **23**: 23–35.
- Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**: 585–595.
- Tang S, Presgraves DC. 2009. Evolution of the *Drosophila* nuclear pore complex results in multiple hybrid incompatibilities. *Science* **323**: 779–782.
- Tao Y, Hartl DL, Laurie CC. 2001. Sex-ratio segregation distortion associated with reproductive isolation in *Drosophila*. *Proc Natl Acad Sci* **98**: 13183–13188.
- Tao Y, Ararape L, Kingan SB, Ke Y, Xiao H, Hartl DL. 2007a. A sex-ratio meiotic drive system in *Drosophila simulans*. II: An X-linked distorter. *PLoS Biol* **5**: e293. doi: 10.1371/journal.pbio.0050293.
- Tao Y, Masly JP, Ararape L, Ke Y, Hartl DL. 2007b. A sex-ratio meiotic drive system in *Drosophila simulans*. I: An autosomal suppressor. *PLoS Biol* **5**: e292. doi: 10.1371/journal.pbio.0050292.
- Theerthagiri G, Eisenhardt N, Schwarz H, Antonin W. 2010. The nucleoporin *Nup188* controls passage of membrane proteins across the nuclear pore complex. *J Cell Biol* **189**: 1129–1142.
- Ting CT, Tsaor SC, Wu ML, Wu CI. 1998. A rapidly evolving homeobox at the site of a hybrid sterility gene. *Science* **282**: 1501–1504.
- True JR, Mercer JM, Laurie CC. 1996. Differences in crossover frequency and distribution among three sibling species of *Drosophila*. *Genetics* **142**: 507–523.
- Tunstall NE, Sirey T, Newcomb RD, Warr CG. 2007. Selective pressures on *Drosophila* chemosensory receptor genes. *J Mol Evol* **64**: 628–636.
- Vasu SK, Forbes DJ. 2001. Nuclear pores and nuclear assembly. *Curr Opin Cell Biol* **13**: 363–375.
- Wallace B. 1948. Studies on sex-ratio in *Drosophila pseudoobscura*. I. Selection and sex-ratio. *Evolution* **2**: 189–217.
- Wente SR, Rout MP. 2010. The nuclear pore complex and nuclear transport. *Cold Spring Harb Perspect Biol* **2**: a000562. doi: 10.1101/cshperspect.a000562.
- Werren JH. 2011. Selfish genetic elements, genetic conflict, and evolutionary innovation. *Proc Natl Acad Sci (Suppl 2)* **108**: 10863–10870.

Received February 29, 2012; accepted in revised form September 24, 2012.

Genome-wide patterns of natural variation reveal strong selective sweeps and ongoing genomic conflict in *Drosophila mauritiana*

Viola Nolte, Ram Vinay Pandey, Robert Kofler, et al.

Genome Res. 2013 23: 99-110 originally published online October 10, 2012

Access the most recent version at doi:[10.1101/gr.139873.112](https://doi.org/10.1101/gr.139873.112)

Supplemental Material <http://genome.cshlp.org/content/suppl/2012/11/08/gr.139873.112.DC1>

Related Content **Genome sequencing reveals complex speciation in the *Drosophila simulans* clade**
Daniel Garrigan, Sarah B. Kingan, Anthony J. Geneva, et al.
Genome Res. August , 2012 22: 1499-1511

References This article cites 107 articles, 20 of which can be accessed free at:
<http://genome.cshlp.org/content/23/1/99.full.html#ref-list-1>

Articles cited in:
<http://genome.cshlp.org/content/23/1/99.full.html#related-urls>

Open Access Freely available online through the *Genome Research* Open Access option.

Creative Commons License This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 3.0 Unported License), as described at <http://creativecommons.org/licenses/by-nc/3.0/>.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Affordable, Accurate
Sequencing.

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>