REVIEW

Molecular Dissection of Quantitative Traits: Progress and Prospects

Andrew H. Paterson¹

Department of Soil and Crop Science, Texas A&M University, College Station, Texas 77843-2474

The "molecular dissection" of quantitative traits began before the demonstration that DNA is the hereditary molecule. Opponents of the Mendelian genetic (particulate) theory argued that many traits showed "blending inheritance," with progeny intermediate between parents. This precipitated one of the great debates in the history of genetics, which was eventually resolved by the realization that blending inheritance might be accounted for by the independent transmission of many different Mendelian factors, together with the modifying effects of environment.

By early in the twentieth century, associations of genetic markers with differences in quantitative phenotypes had already been noted. For example, Sax (1923) noted association of differences in bean seed weight with seed-coat pigmentation, concluding that "*Factor differences for seed weight are . . . linked with . . . factors which determine the color of the pigment. Size differences may be effected by the independent action of . . . factors in different linkage groups. These factors, when combined, have a cumulative effect. The size factors in different chromosomes may not be equal in their effect.*"

The principles outlined by these pioneering studies were widely employed, both in model systems and domesticated species. However, until recently most maps contained too few genetic markers to permit either identification of suites of quantitative trait loci (QTLs) accounting for the bulk of variation in a phenotype, or precise mapping of QTLs to specific chromosomal locations. As stated by Thoday (1961), "*The main practical limitation of the technique seems to be the availability of suitable markers, and the time that can be given to the considerable work involved.*"

A growing body of molecular tools, first from protein polymorphisms (Markert and Moller 1959), and, more recently, from DNA polymorphisms, has contributed to the development of

"complete" genetic maps, encompassing all regions of all chromosomes in a plant or animal species. Complete genetic maps permit comprehensive analysis of the genome of an organism, revealing the locations of QTLs influencing virtually any characteristic that can be measured.

The development of DNA-based genetic mapping technology was propelled by the search for genes responsible for human genetic diseases and other traits (Botstein et al. 1980). QTLs influencing medically important phenotypes such as high blood pressure (Rapp et al. 1989) and hypertension (Jacob et al. 1991) are being identified in mammalian models. Recent results have illustrated the usefulness of genome mapping in dissecting complex behavioral characteristics (Plomin et al. 1994), such as avoidance, exploration (Neiderheiser et al. 1992), and substance abuse (Crabbe et al. 1994; Quock et al. 1994) in rodents, and reading disability in humans (Cardon et al. 1994). In one recent case, the effectiveness of a human disease agent (malaria) at parasitizing its vector (mosquito) has been dissected into several QTLs (Severson et al. 1995). Finally, many complex traits mapped in domestic animals, such as obesity (Andersson et al. 1994; Pellemounter et al. 1995), lactation (Georges et al. 1995), and others (cf. Cockett et al. 1994; Georges et al. 1994) may be relevant to human phenotypes. An ever-clearer picture of the correspondence between the chromosomes of humans and other mammals helps in the application of such comparative information (cf. O'Brien et al. 1993).

Plant breeding has provided a fertile field for molecular dissection of quantitative traits. Charles Darwin recognized that plant breeding programs were very useful for the study of heredity. The realization that crop improvement could benefit from DNA marker-assisted selection (cf. Paterson et al. 1991b) has led to rapid growth in molecular mapping of agriculturally important crops. Genetic mapping of traits associated with many aspects of crop productivity is published or

¹E-MAIL ahp2343@bioch.tamu.edu; FAX (409) 845-0456.

PATERSON

in progress, including basic growth patterns such as plant height (cf. Koester et al. 1993; Lin et al. 1995; M. Pereira, M. Lee, and P. Rayapati, unpubl.), tillering, rhizomatousness (Paterson et al. 1995a), flowering time (cf. Koester et al. 1993; Kowalski et al. 1994; Li et al. 1995a; Lin et al. 1995), and other morphological variants (Kennard et al. 1994); yield components such as the size, number, and harvestability of seed (Stuber et al. 1987, 1992; Abler et al. 1991; Fatokun et al. 1992; Doebley et al. 1994; Schon et al. 1994; Paterson et al. 1995a,b), biomass, and/or growth rates (DeVicente and Tanksley 1993; Bradshaw and Stettler 1995); quality parameters such as composition of fruit or seed (Paterson et al. 1988, 1990, 1991a; Weller et al. 1988; DeVicente and Tanksley 1993; Teutonico and Osborn 1994), shape of tubers (Van Eck et al. 1994), or specific gravity of wood (Groover et al. 1994); and the impact of adverse factors such as diseases (Bubeck et al. 1993; Leonards-Schippers et al. 1994; Wang et al. 1994; Jung et al. 1995; Li et al. 1995b), insects (Nienhuis et al. 1987; Bonierbale et al. 1994) and abiotic factors (Martin et al. 1989; Reiter et al. 1991), and the evolution of novel organs (Doebley et al. 1990). This partial list illustrates the breadth of topics in crop improvement in which QTL mapping is affording new insights.

In addition to medicine and agriculture, QTLs playing critical roles in evolution have been characterized. Bradshaw et al. (1995) recently mapped QTLs controlling floral traits that determine pollinator preference in two *Mimulus* species. These species are sympatric (grow in the same places) and can produce fertile hybrid progeny—but do not do so in nature, because one species is strictly bumblebee pollinated, and the other is strictly hummingbird pollinated. A few QTLs with large effects determine the floral characteristics (flower color, nectar volume and concentration, and stamen and pistil length) that influence pollinator preference—these mutations (QTLs) may represent the initial steps in reproductive isolation of these two species.

Analysis and Interpretation of QTL Mapping Experiments

Algorithms have been developed for QTL mapping in a wide range of pedigrees and experimental designs, including F_2 , backcross, recombinant inbred, and many other designs (cf. Weller 1986; Lander and Botstein 1989; Knapp et al. 1991; Luo and Kearsey 1991, 1992; Carbonell et al. 1992;

Darvasi and Weller 1992; Haley and Knott 1992; Knott and Haley 1992; Jansen 1992, 1993, 1994; Mackinnon and Georges 1992; Darvasi et al. 1993; Moreno-Gonzalez 1993; Rodolphe and Lefort 1993; Zeng 1993, 1994; Cardon and Fulker 1994; Eaves 1994; Haley et al. 1994; Jiang and Zeng 1995). All share the basic principle of testing correlation between marker genotypes and quantitative phenotypes. As a result of the availability of complete genetic maps, current analytical methods are able to use information from multiple markers that flank a QTL, in contrast to earlier methods that were limited to information from single markers at unknown distance and direction from the QTL. This permits more accurate estimates of location and phenotypic effect, of individual QTLs. Although many procedures to date have been parametric, approaches to handling nonparametric traits are beginning to emerge (Kruglyak and Lander 1995).

Perhaps the single most important consideration in analysis and interpretation of QTL data is the threshold employed for inferring that a QTL is statistically significant. Because QTL mapping involves many analyses of independent (unlinked) genetic markers throughout a genome, there are many opportunities for false-positive results. Stringent significance thresholds must be employed to avoid these. Nominal significance criteria of 99.8% or more for any single QTL are usually necessary to assure an “experiment-wide” confidence level of 95% for all QTLs reported across a genome. Appropriate criteria are often described in detail when an analytical approach is developed (cf. Lander and Botstein 1989). Alternatively, methods for empirical calculation of criteria appropriate to particular data sets have been described (Churchill and Doerge 1994; Rebai et al. 1994, 1995). As new opportunities for “comparative analysis” of previously published QTLs emerge (cf. Lin et al. 1995; Paterson et al. 1995b), it becomes ever more important that published QTLs satisfy statistical criteria that minimize the likelihood of false-positive results.

Genetic Basis of Quantitative Variation

QTL mapping is useful for investigating specific properties of individual genes contributing to quantitative traits. In contrast, classical quantitative genetics describes the aggregate behavior of suites of genes influencing a trait. The aggregate descriptions of quantitative genetics are very useful for guiding manipulation of plant and animal

QTL MAPPING: PROGRESS AND PROSPECTS

gene pools by breeders; however, an understanding of quantitative inheritance at the molecular level requires detailed descriptions of individual genes, and this is made possible by QTL mapping.

The reconciliation of results from QTL mapping with expectations from classical theory is progressing but remains incomplete. Basic properties of individual QTLs such as additivity and dominance are readily tested, and a wide range of different modes of gene action are evident for different QTLs (cf. Paterson et al. 1991a). Many basic genetic phenomena are readily reconciled with results from QTL mapping, such as transgression (progeny that exceeds parental genetic potential; Paterson et al. 1988; deVicente and Tanksley 1993) and heterosis (hybrid vigor; Stuber et al. 1992; Xiao et al. 1995). However, two particular areas continue to be controversial: the number of genes with pronounced effects on quantitative traits and the importance of interlocus interactions ("epistasis").

Gene Numbers and Phenotypic Effects

Geneticists have long debated the degree of complexity of quantitative traits (cf. Dove 1993). Many theories, ranging from "virtually infinite numbers of genes with tiny effects", to "few genes with large effects" have been proposed, championed, questioned, revised, rejected, and reincarnated. Geneticists realized that some assumptions invoked to simplify quantitative models, such as equality of gene effects and strict additivity of gene action, were unlikely to describe individual quantitative trait loci precisely. Such assumptions were tolerated so long as models were reasonably accurate in predicting the behavior of study populations. It was no particular surprise that QTL mapping showed such assumptions to be incorrect. However, it has remained controversial whether the results of QTL mapping experiments reflect the true complexity of quantitative inheritance or simply detect a subset of (relatively large) gene effects.

Figure 1 shows histograms of the fractional phenotypic variance explained by QTLs affecting several quantitative traits, mapped in sorghum and rice. Similar data have been published previously for tomato (Paterson et al. 1991a). In each case, a relatively small number of genes accounts for very large portions of phenotypic variance, with increasing numbers of genes accounting for progressively smaller portions of variance, until the significance threshold is reached. Curiously, the number of QTLs observed in "first-generation

Figure 1 Distribution of percent phenotypic variance explained by QTLs affecting several traits in sorghum and rice. The rice data includes 33 QTLs affecting height, flowering time, resistance to sheath blight, seed size, floret number, and spikelet sterility (Li et al. 1995a,b; Z. Li, S.R.M. Pinson, W.D. Park, A.H. Paterson, and J.W. Stansel, in prep.; Z. Li, S.R.M. Pinson, A.H. Paterson, W.D. Park, and J.W. Stansel, in prep.). The sorghum data include 37 QTLs affecting height, flowering time (Lin et al. 1995), rhizomatousness, tillering, regrowth after overwintering (Paterson et al. 1995a), and seed size (Paterson et al. 1995b).

mapping" (e.g., of a BC₁ or F₂ population; cf. Paterson et al. 1988) is often similar to the "effective number of factors" predicted by classical models. However, if genes explaining large portions of phenotypic variance are rendered homozygous, the significance threshold is reduced, and yet additional genes explaining even smaller portions of phenotypic variance are revealed (Paterson et al. 1990).

Although the "end" of this stepwise process of revealing genes of smaller and smaller effect is not clear, a large and growing body of data continues to suggest that relatively few genes account for the bulk of variation in many populations, with ever-larger numbers of genes contributing ever-smaller portions of variance (cf. Lande and Thompson 1990). This result will be supported further in a discussion of comparative analysis, with the finding that genes accounting for the bulk of variation in a trait appear to correspond in different reproductively isolated species.

Epistasis

Almost universally, the collective activities of mapped QTLs explain only a portion of the phe-

PATERSON

notypic difference between parents, even with nearly complete genome coverage by DNA markers. Classical evidence has strongly suggested the importance of epistasis, or nonlinear interactions between unlinked genetic loci, in quantitative inheritance (cf. Spickett and Thoday 1966; Falconer 1981; Mather and Jinks 1982; Pooni et al. 1987; Allard 1988).

Until recently, QTL mapping experiments have provided very little evidence in support of the importance of epistasis, with nonlinear interactions among DNA marker loci reaching statistical significance at approximately the frequency that would be expected to occur by chance (cf. Edwards et al. 1987; Paterson et al. 1988, 1991a). Hints of epistasis among QTLs have come from the demonstration of “genetic background effects” on quantitative traits in *Drosophila* (Spassky et al. 1965), rice (Kinoshita et al. 1982, Sato and Sakamoto 1983), and tomato (Tanksley and Hewitt 1988) and from the discovery of occasional loci reported to show interaction with multiple unlinked sites in a genome (cf. Paterson et al. 1988).

Modified experimental designs may reconcile QTL mapping results with the importance attributed to epistasis in classical studies. Doebley et al. (1995) developed genetic stocks differing by two QTLs suspected to interact epistatically, but otherwise uniform in genetic background, and found strong evidence for epistasis between the loci. Lark et al. (1995) used recombinant inbred lines to reduce the complexity of interactions and replicate phenotypic measurements, and found evidence of epistasis between QTLs in genetic control of several agronomic traits. Each of these results suggests that the absence of epistasis in previous QTL mapping studies may have been the result of insufficient replication and/or insufficient statistical resolution to detect interactions, in the presence of many QTLs with large main effects. Although the effects of some QTLs appear independent of interacting loci (cf. Paterson et al. 1990), in at least some cases it is becoming clear that “the whole” is, indeed, greater

than the sum of the parts. Improved documentation of epistasis may account for a portion of the “genetic difference between parents” that was unexplained previously by QTL mapping.

It is noteworthy that analysis of epistasis is even more subject to the problem of false-positive results than is analysis of individual QTLs (as discussed above). An important need in QTL research is a generally applicable set of criteria for inferring statistical significance of interactions between genetic loci, which controls “experiment-wise” error rates.

Comparative Analysis of QTLs

“Comparative mapping,” the alignment of the chromosomes of related taxa based on genetic mapping of common DNA markers, affords many benefits to genome analysis. Many DNA probes can be cross-utilized among different species in the same taxonomic family, increasing the number and map density of genetic markers available for many genera simultaneously. In one well-studied plant family, the Poaceae, extensive conservation of gene repertoire and order (cf. Hulbert et al. 1990; Ahn and Tanksley 1993; Ahn et al. 1993a,b; Kurata et al. 1994) has led to the suggestion that the cultivated cereals (diverse genera within the family Poaceae) might be treated as essentially a “single genetic system” (Bennetzen and Freeling 1993; Helentjaris 1993; Shields 1993).

Comparative mapping is also useful in molecular dissection of quantitative traits (Fig. 2). A close correspondence among QTLs affecting complex traits such as seed size, as well as traits of varying complexity in different taxa such as disarticulation (“shattering”) of the inflorescence and day-neutral flowering, has been shown for sorghum, sugarcane, maize, wheat, barley, and rice (Paterson et al. 1995b). Correspondence among an unexpectedly high proportion of genes affecting height and flowering of maize, sorghum, and other grasses has also been reported (Lin et al.

Figure 2 Comparative mapping of QTLs affecting seed size and shattering, on linkage group C of sorghum, and corresponding chromosomes of rice and maize (revised from Paterson et al. 1995b). Correspondence among QTLs for seed size is [(M) maize; (S) sorghum; (R) rice] (M c.1 // S lg C // R c.10); (M c.1 // R c.3); (M c.9 // R c.3); and correspondence between QTLs for shattering is: (M c.1 // M c.5 // S lg. C // R c.9). The sorghum linkage group C locus affecting shattering is “discrete,” accounting for virtually 100% of phenotypic variation (Paterson et al. 1995a).

OTL MAPPING: PROGRESS AND PROSPECTS

Figure 2 (See facing page for legend.)

PATERSON

1995; M. Pereira, M. Lee, and P. Rayapati, unpubl.). Correspondence of QTLs in different species of the plant genera *Lycopersicon* (Paterson et al. 1991a) and *Vigna* (Fatokun et al. 1992) had been reported previously; however, this correspondence spans rather short periods of genetic divergence, and the relative promiscuity of plant "species" makes it difficult to preclude the possibility of recent gene flow.

Correspondence of QTLs on duplicated chromosome segments within a particular species has also been suggested, indicating that chromosome duplication may contribute to polygenic inheritance. Specifically, pairs of loci affecting shattering of the maize inflorescence (see Fig. 2; Doebley et al. 1990; Paterson et al. 1995b), maize height and flowering, and sorghum height (Lin et al. 1995) fall on corresponding duplicated chromosome segments. Chromosome duplication in these taxa preceded domestication by millions of years. Domestication, in the past 10,000 years, most likely imparted strong selective advantages to reduced shattering, reduced height, and early flowering. Consequently, if duplicated genes retained common functions, it is intuitive that QTLs might be found on homologous (duplicated) chromosomal sites (Paterson et al. 1995b). Ongoing research in these and other polyploids is likely to reveal additional cases of putatively duplicated QTLs.

Implications of QTL Correspondence

The suggestion that mutations in corresponding genes may account for phenotypic variation in taxa that have been reproductively isolated for millions of years has many implications. Perhaps first and foremost, QTL analysis in one taxon may predict results in other taxa. Such predictive value would make QTL mapping results more broadly applicable than was previously envisioned, enabling research on facile systems to be extrapolated to more difficult ones.

Correspondence of QTLs across diverse taxa also provides strong empirical support for the use of model systems in research on complex phenotypes. For example, the ease of genetic analysis in rodents and domesticated mammals has permitted mapping of genes associated with diabetes, hypertension, obesity, alcohol/drug addiction, and other medically important phenotypes. The difficulties associated with mapping complex traits in humans may be partly overcome by the possibility of cloning QTLs from mouse or other

mammals that also account for phenotypic variation in humans. In a similar manner, crop plants that grow particular organs of extraordinary size, such as the enlarged root of turnip, inflorescence ("curd") of cauliflower, or fruit of tomato, might be used to isolate genes important to particular aspects of plant growth and development.

Finally, correspondence among QTLs in different taxa strongly supports the hypothesis that a relatively small number of mutations (genes) accounts for a large portion of phenotypic variation in many populations. If the possibilities for such mutations were infinite and all equally probable, such correspondence seems unlikely. In contrast, if a few genes play disproportionately large roles in genetic control of a trait, then mutations in (one of) these genes are more likely to have effects sufficiently large to drive the mutant allele to fixation.

It remains unclear whether the correspondence among QTLs that is found in interspecific crosses (cf. Paterson et al. 1991a; Fatokun et al. 1992; Lin et al. 1995; M. Pereira, M. Lee, and P. Rayapati, unpubl.) is paralleled by correspondence of QTLs in more closely related genotypes such as elite crop cultivars (cf. Beavis et al. 1991). This may indicate that crop gene pools are homogeneous at mutant alleles with large effects and fixed during the initial stage of crop domestication and that variation within elite gene pools of individual crops is a result of subsequent mutations at a larger number of loci with smaller effects.

Toward Cloning of QTLs

Map-based cloning has been used successfully to isolate mutant alleles imparting discrete phenotypes; however, QTLs continue to be refractory to cloning, for several reasons. Individual QTLs exert a relatively small effect on phenotype, which can be obscured by the effects of other genes or by nongenetic factors that are difficult to control. Breeding approaches can be used to create populations segregating for individual QTLs (cf. Paterson et al. 1990; Dorweiler et al. 1993) but are time and labor intensive. The resolution with which QTLs can be mapped is rather coarse; even using "second-generation" experimental designs based on progeny testing, a resolution of ~3 cM is the best achieved to date. In a genome such as that of human or many crop plants, 3 cM represents ~0.1% of the genome, containing perhaps 100 or

QTL MAPPING: PROGRESS AND PROSPECTS

more genes. Identification of these genes can be difficult (Gardiner and Mural 1995). Furthermore, complementation testing of so many genes, requiring transformation of the dominant allele into the recessive genotype, is an enormous undertaking, even in taxa that enjoy good transformation systems.

Several recent developments make cloning of QTLs more tractable. First, the technology for cloning and manipulating large segments of intact chromosomal DNA is becoming ever more efficient, facilitating "chromosome walks" (Steinmetz et al. 1982). A decade ago, chromosome walks were based on phage and cosmid vectors with maximal carrying capacity of 15,000–30,000 nucleotides of exogenous DNA. "Artificial chromosomes" carrying 10 times this amount of DNA can now be developed routinely and are publicly available for several organisms. Initially, artificial chromosomes containing exogenous DNA were propagated in yeast (Burke et al. 1987); however, bacterial systems (Shizuya et al. 1992) are quickly gaining acceptance. Improved vectors with new combinations of selectable markers and systems that facilitate transformation of megabase DNA segments (cf. Pachnis et al. 1990; Pavan et al. 1990; Eliceriri et al. 1991; Strauss and Jaenisch 1992) will accelerate the search for QTLs, especially in combination with site-specific recombination systems (cf. Sauer and Henderson 1988).

There is a large and growing number of approaches for identifying candidate genes that might be QTLs. Direct isolation of genes from megabase DNA clones is becoming more efficient (Lovett et al. 1991; Parimoo et al. 1991; Collins 1992; Morgan et al. 1992). Several PCR-based techniques are now available for identifying genes expressed specifically in particular tissues, developmental stages, or genotypes (Hara et al. 1991; Liang and Pardee 1992; Sun et al. 1992; Lopez-Fernandez and Mazo 1993; Liang et al. 1994). Moreover, random sequencing of cDNAs is adding to a rapidly growing repertoire of plant genes that are readily accessible (cf. Hofte et al. 1993; Kurata et al. 1994; Newman et al. 1994; Sasaki et al. 1994). By combining such information with genetic mapping to test which candidates are located in the target chromosomal region, isolation of candidate genes might be accelerated.

The identification of multiple independent mutant alleles at QTLs would permit testing of correlation between phenotype and mutation in

a particular transcript, as has been used in map-based cloning of genes in humans and other organisms. Using comparative information, genotypes of different taxa might be identified that show a common phenotype that can be explained by mutations mapping to corresponding chromosomal locations (cf. Paterson et al. 1995b; Lin et al. 1995). Sequencing of dominant and recessive alleles at "candidate genes" in each taxon might allow a mutation in a particular transcript to be correlated with the mutant phenotype, in much the same manner as has been employed to identify several disease genes in human.

The suggestion that QTLs might be cloned by identifying allelic mutations with discrete effects (Robertson 1985) has also come closer to realization. Comparative analysis of different taxa provides more opportunities to seek such discrete mutations, and several examples have been reported (Beavis et al. 1991; Lin et al. 1995; Paterson et al. 1995b; M. Pereira, M. Lee, and P. Rayapati, unpubl.). Identification of discrete mutations corresponding to QTLs offers a more rapid and efficient means to dissect quantitative traits than do breeding approaches (Paterson et al. 1990; Dorweiler et al. 1993).

Summary

QTL mapping is an increasingly useful approach to the study and manipulation of complex traits important in agriculture, evolution, and medicine. The molecular dissection of quantitative phenotypes, supplementing the principles of classical quantitative genetics, is accelerating progress in the manipulation of plant and animal genomes. A growing appreciation of the similarities among different organisms and the usefulness of comparative genetic information is making genome analysis more efficient, and providing new opportunities for using model systems to overcome the limitations of less-favorable systems.

The expanding repertoire of techniques and information available for studying heredity is removing obstacles to the cloning of QTLs. Although QTL mapping alone is limited to a resolution of 0.1%–1.0% of a genome, use of QTL mapping in conjunction with a search for mapped candidate genes, with emerging technologies for isolation of genes expressed under conditions likely to account for the quantitative phenotype, and with ever more efficient megabase DNA manipulation and characterization bodes

PATERSON

well for the prospect of isolating the genetic determinants of QTLs in the foreseeable future. In the words of Thoday (1961), "*An extensive attack on quantitative genetics made from this point of view as well as the biometric approach should be a great help in answering questions concerning the nature of polygenes . . .*"

ACKNOWLEDGMENTS

I thank M. Burow, C. Katsar, and T.-H. Lan for critical reading of the manuscript and numerous colleagues and staff for valuable discussions. Aspects of this research were funded by the International Consortium for Sugarcane Biotechnology, U. S. Department of Agriculture (Plant Genome and Biotechnology Risk Assessment Programs), Pioneer Hibred International, Texas Higher Education Coordinating Board, and Texas Agricultural Experiment Station.

REFERENCES

- Abler, B.S.B., M.D. Edwards, and C.W. Stuber. 1991. Isozymatic identification of quantitative trait loci in crosses of elite maize inbreds. *Crop Sci.* **31**: 267–274.
- Ahn, S. and S.D. Tanksley. 1993. Comparative linkage maps of the rice and maize genomes. *Proc. Nat. Acad. Sci.* **90**: 7980–7984.
- Ahn, S., J.A. Anderson, M.E. Sorrells, and S.D. Tanksley. 1993. Homeologous relationships of rice, wheat, and maize chromosomes. *Mol. & Gen. Genet.* **241**: 483–490.
- Allard, R.W. 1988. Genetic changes associated with the evolution of adaptedness in cultivated plants and their wild progenitors. *J. Hered.* **79**: 225–238.
- Altenbach et al.. 1995. USDA Plant Genome Research Program. *Ad. Agron.* (in press).
- Andersson, L., C.S. Haley, H. Ellegren, S.A. Knott, M. Johansson, K. Andersson, L. Andersson-Eklund, I. Edfors-Lilja, M. Fredholm, I. Hansson, J. Hakansson, and K. Lundstrom. 1994. Genetic mapping of quantitative trait loci for growth and fatness in pigs. *Science* **263**: 1771–1774.
- Beavis, W.D., D. Grant, M. Albertsen, and R. Fincher. 1991. Quantitative trait loci for plant height in four maize populations and their associations with qualitative loci. *Theor. Appl. Genet.* **83**: 141–145.
- Bennetzen, J.L. and M. Freeling. 1993. Grasses as a single genetic system: Genome composition, collinearity and compatibility. *Trends Genet.* **9**: 259–261.
- Bonierbale, M.W., R.L. Plaisted, O. Pineda, and S.D. Tanksley. 1994. QTL analysis of trichome-mediated insect resistance in potato. *Theor. Appl. Genet.* **87**: 973–987.
- Botstein, D., R.L. White, M. Skolnick, and R.W. Davis. 1980. Construction of a genetic linkage map in man using restriction fragment length polymorphism. *Am. J. Hum. Genet.* **32**: 314–331.
- Bovenhuis, H. and J.I. Weller. 1994. Mapping and analysis of dairy cattle quantitative trait loci by maximum likelihood methodology using milk protein genes as genetic markers. *Genetics* **137**: 267–280.
- Bradshaw, H.D. and R.F. Stettler. 1995. Molecular genetics of growth and development in Populus. IV. Mapping QTLs with large effects on growth, form, and phenology of traits. *Genetics* **139**: 963–973.
- Bradshaw, H.D., S.M. Wilbert, K.G. Otto, and D.W. Schemske. 1995. Genetic mapping of floral traits associated with reproductive isolation in monkeyflowers (*Mimulus*). *Nature* **376**: 762–765.
- Bubeck, D.M., M.M. Goodman, W.D. Beavis, and D. Grant. 1993. Quantitative trait loci controlling resistance to gray leaf spot in maize. *Crop Sci.* **33**: 838–847.
- Burke, D.T., G.F. Carle, and M.V. Olson. 1987. Cloning of large segments of exogenous DNA into yeast by means of artificial chromosome vectors. *Science* **236**: 806–812.
- Carbonell, E.A., T.M. Gerig, E. Balansard, and M.J. Asins. 1992. Interval mapping in the analysis of nonadditive quantitative trait loci. *Biometrics* **48**: 305–315.
- Cardon L.R. and D.W. Fulker. 1994. The power of interval mapping of quantitative trait loci, using selected sib pairs. *Am. J. Hum. Genet.* **55**: 825–833.
- Cardon L.R., S.D. Smith, D.W. Fulker, W.J. Kimberling, B.F. Pennington, and J.C. Defries. 1994. Quantitative trait locus for reading disability on chromosome 6. *Science* **266**: 276–279.
- Churchill, G.A. and R.W. Doerge. 1994. Empirical threshold values for quantitative trait mapping. *Genetics* **138**: 963–971.
- Cockett, N.E., S.P. Jackson, T.L. Shay, D. Nielsen, S.S. Moore, M.R. Steele, W. Barendse, R.D. Green, and M. Georges. 1994. Chromosomal localization of the callipyge gene in sheep (*Ovis aries*) using bovine DNA markers. *Proc. Natl. Acad. Sci.* **91**: 3019–3023.
- Collins, F.S. 1992. Positional cloning: Let's not call it reverse anymore. *Nature Genet.* **1**: 3–6.
- Crabbe, J.C., J.K. Belknap, and K.J. Buck. 1994. Genetic animal models of alcohol and drug abuse. *Science* **264**: 1715–1723.
- Darvasi, A., A. Weinreb, V. Minke, J.I. Weller, and M. Soller. 1993. Detecting marker-QTL linkage and estimating QTL gene effect and map location using a saturated linkage map. *Genetics* **134**: 943–951.

QTL MAPPING: PROGRESS AND PROSPECTS

- Darvasi, A. and J.I. Weller. 1992. On the use of the moments method of estimation to obtain approximate maximum likelihood estimates of linkage between a genetic marker and a quantitative locus. *Heredity* **68**: 43–46.
- DeVicente, M.C. and S.D. Tanksley. 1993. QTL analysis of transgressive segregation in an intraspecific tomato cross. *Genetics* **134**: 585–596.
- Doebley, J. and A. Stec. 1993. Inheritance of the morphological differences between maize and teosinte: Comparisons of results for two F₂ populations. *Genetics* **134**: 559–570.
- Doebley, J., A. Stec, J. Wendel, and M. Edwards. 1990. Genetic and morphological analysis of a maize-teosinte F₂ population: Implications for the origin of maize. *Proc. Natl. Acad. Sci.* **87**: 9888.
- Doebley, J., A. Bacigalupo, and A. Stec. 1994. Inheritance of kernel weight in two maize-teosinte hybrid populations: Implications for crop evolution. *J. Hered.* **85**: 191–195.
- Doebley, J., A. Stec, and C. Gustus. 1995. *teosinte branched 1* and the origin of maize: Evidence for epistasis and the evolution of dominance. *Genetics* **141**: 333–346.
- Dorweiler, J., A. Stec, J. Kermicle, and J. Doebley. 1993. *Teosinte glume architecture-1*: A genetic locus controlling a key step in maize evolution. *Science* **262**, 233.
- Dove, W.F. 1993. The gene, the polygene, and the genome. *Genetics* **134**: 999–1002.
- Eaves, L.J. 1994. Effect of genetic architecture on the power of human linkage studies to resolve the contribution of quantitative trait loci. *Heredity* **72**: 175–192.
- Edwards, M.D., C.W. Stuber, and J.F. Wendel. 1987. Molecular-marker-facilitated investigations of quantitative-trait loci in maize. I. Numbers, genomic distribution and types of gene action. *Genetics* **116**: 113–125.
- Eliceriri B., T. Labella, Y. Hagino, A. Srivastava, D. Schlessinger, G. Pilia, G. Palmieri, and M. D'Urso. 1991. Stable integration and expression in mouse cells of yeast artificial chromosomes harboring human genes. *Proc. Natl. Acad. Sci.* **88**: 2179–2183.
- Falconer, D.S. 1981. *Introduction to quantitative genetics*, 2nd ed. Longman Press, London, UK.
- Fatokun, C.A., D.I. Menacio-Hautea, D. Danesh, and N.D. Young. 1992. Evidence for orthologous seed weight genes in cowpea and mungbean, based upon RFLP mapping. *Genetics* **132**: 841–846.
- Fehr, W. 1984. "Genetic contributions to yield gains of five major crop plants," Special Publication no. 7, Crop Science Society of America, Madison, WI.
- Fulker, D.W. and L.R. Cardon. 1994. A sib-pair approach to interval mapping of quantitative trait loci. *Am. J. Hum. Genet.* **54**: 1092–1103.
- Gardiner, K. and R. Mural. 1995. Getting the message: Identifying transcribed sequences. *Trends Genet.* **11**: 77–79.
- Georges, M., A.B. Dietz, A. Mishra, D. Nielsen, L.S. Sargeant, A. Sorensen, M.R. Steele, X. Zhao, H. Leipold, J.E. Womack, and M. Lathrop. 1994. Microsatellite mapping of the gene causing Weaver disease in cattle will allow the study of an associated quantitative trait locus. *Proc. Natl. Acad. Sci.* **90**: 1058–1062.
- Georges, M., D. Nielsen, M. MacKinnon, A. Mishra, R. Okimoto, A.T. Pasquino, L.S. Sargeant, A. Sorensen, M.R. Steele, X. Zhao, J.E. Womack, and I. Hoeschele. 1995. Mapping quantitative trait loci controlling milk production in dairy cattle by exploiting progeny testing. *Genetics* **139**: 907–920.
- Groover, A., M. Devey, T. Fiddler, J. Lee, R. Megraw, T. Mitchell-Olds, B. Sherman, S. Vujcic, C. Williams, and D. Neale. 1994. Identification of quantitative trait loci influencing wood specific gravity in an outbred pedigree of loblolly pine. *Genetics* **138**: 1293–1300.
- Haley, C.S. and S.A. Knott. 1992. A simple regression method for mapping quantitative trait loci in line crosses using flanking markers. *Heredity* **69**: 315–324.
- Haley, C.S., S.A. Knott, and J.-M. Elsen. 1994. Mapping quantitative trait loci in crosses between outbred lines using least squares. *Genetics* **136**: 1195–1207.
- Hara, E., T. Kato, S. Nakada, S. Sekiya, and K. Oda. 1991. Subtractive cDNA cloning using oligo(dT)30 latex and PCR: Isolation of cDNA clones specific to undifferentiated human embryonal carcinoma cells. *Nucleic Acids Res.* **19**: 7097–7104.
- Helentjaris, T. 1993. Implications for conserved genomic structure among plant species. *Proc. Natl. Acad. Sci.* **90**: 8308–8309.
- Hofte, H., et al. 1993. An inventory of 1152 expressed sequence tags obtained by partial sequencing of cDNAs from *Arabidopsis thaliana*. *Plant J.* **4**: 1051–1061.
- Hulbert, S.H., T.E. Richter, J.D. Axtell, and J.L. Bennetzen. 1990. Genetic mapping and characterization of sorghum and related crops by means of maize DNA probes. *Proc. Natl. Acad. Sci.* **87**: 4251–4255.
- Jacob, H.J., K. Lindpainter, S.E. Lincoln, K. Kusumi, R.K. Bunker, Y.-P. Mao, D. Ganten, V.J. Dzau, and E.S. Lander. 1991. Genetic mapping of a gene causing hypertension in the stroke-prone spontaneously hypertensive rat. *Cell* **67**: 213–224.
- Jansen, R.C. 1992. A general mixture model for mapping

PATERSON

- quantitative trait loci by using molecular markers. *Theor. Appl. Genet.* **85**: 252–260.
- . 1993. Interval mapping of multiple quantitative trait loci. *Genetics* **135**: 205–211.
- . 1994. Controlling the type I and type II errors in mapping quantitative trait loci. *Genetics* **138**: 871–881.
- Jansen, R.C. and P. Stam. 1994. High resolution of quantitative traits into multiple loci via interval mapping. *Genetics* **136**: 1447–1455.
- Jiang, C. and Z.-B. Zeng. 1995. Multiple trait analysis of genetic mapping for quantitative trait loci. *Genetics* **140**: 1111–1127.
- Jung, M., T. Weldekidan, D. Schaff, A. Paterson, S. Tingey, and J. Hawk. 1995. Generation means analysis and genetic mapping of anthracnose stalk rot resistance in maize. *Theor. Appl. Genet.* (in press).
- Kennard, W.C., M.K. Slocum, S.S. Figdore, and T.C. Osborn. 1994. Genetic analysis of morphological variation in *Brassica oleracea* using molecular markers. *Theor. Appl. Genet.* **87**: 721–732.
- Kinoshita, T. and N. Shinbashi. 1982. Identification of dwarf genes and their character expression in the isogenic background. *Jpn. J. Breed.* **32**: 219–231.
- Knapp, S.J. and W.C. Bridges. 1990. Using molecular markers to estimate quantitative trait locus parameters, power, and genetic variances for unreplicated and replicated progeny. *Genetics* **126**: 769–777.
- Knapp, S.J., W.C. Bridges, and D. Birkes. 1990. Mapping quantitative trait loci using molecular marker linkage maps. *Theor. Appl. Genet.* **79**: 583–592.
- Knott, S.A. and C.S. Haley. 1992. Maximum likelihood mapping of quantitative trait loci using full-sib families. *Genetics* **132**: 1211–1222.
- Koester, R.P., P.H. Sisco, and C.W. Stuber. 1993. Identification of quantitative trait loci controlling days to flowering and plant height in two near isogenic lines of maize. *Crop Sci.* **33**: 1209–1216.
- Kowalski, S.D., T.-H. Lan, K.A. Feldmann, and A.H. Paterson. 1994. Comparative mapping of *Arabidopsis thaliana* and *Brassica oleracea* chromosomes reveals islands of conserved gene order. *Genetics* **138**: 499–510.
- Kruglyak, L. and E.S. Lander. 1995. A nonparametric approach for mapping quantitative trait loci. *Genetics* **139**: 1421–1428.
- Kurata, N., G. Moore, Y. Nagumara, T. Foote, M. Yano, Y. Minobe, and M. Gale. 1994. Conservation of genome structure between rice and wheat. *Bio/Technology* **12**: 276–278.
- Kurata, N., Y. Nagamura, K. Yamamoto, Y. Harushima, N. Sue, J. Wu, B.A. Antonio, A. Shomura, T. Shimizu, S.Y. Lin, T. Inoue, A. Fukuda, T. Shimano, Y. Kuboki, T. Toyama, Y. Miyamoto, T. Kirihaara, K. Hayasaka, A. Miyao, L. Monna, H.S. Zhong, Y. Tamura, Z.X. Wang, Y. Momma, Y. Umehara, M. Yano, T. Sasaki, and Y. Minobe. 1994. A 300 kilobase interval genetic map of rice including 883 expressed sequences. *Nature Genet.* **8**: 365–372.
- Lande, R. and R. Thompson. 1990. Efficiency of marker-assisted selection in the improvement of quantitative traits. *Genetics* **124**: 743–756.
- Lander, E.S. and D. Botstein. 1989. Mapping Mendelian factors underlying quantitative traits using RFLP linkage maps. *Genetics* **121**: 185–199; Corrigendum. *Genetics* **136**: 705.
- Lark, K.G., K. Chase, F. Adler, L.M. Mansur, and J.H. Orf. 1995. Interactions between quantitative trait loci in soybean in which trait variation at one locus is conditional upon a specific allele at another. *Proc. Natl. Acad. Sci.* **92**: 4656–4660.
- Leonards-Schippers, C., W. Gieffers, R. Schaefer-Pregl, E. Ritter, S.J. Knapp, F. Salamini, and C. Gebhardt. 1994. Quantitative resistance to *Phytophthora infestans* in potato: A case study for QTL mapping in an allogamous plant species. *Genetics* **137**: 68–77.
- Li, Z., S.R.M. Pinson, J.W. Stansel, and W.D. Park. 1995a. Identification of quantitative trait loci for heading date and plant height in cultivated rice. *Theor. Appl. Genet.* **91**: 374–381.
- Li, Z., S.R.M. Pinson, M.A. Marchetti, J.W. Stansel, and W.D. Park. 1995b. Characterization of quantitative trait loci in cultivated rice contributing to field resistance to sheath blight (*Rhizoctonia solani*). *Theor. Appl. Genet.* **91**: 374–381.
- Liang, P. and A.B. Pardee. 1992. Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* **257**: 967–971.
- Liang, P., L. Averboux, W. Zhu, and A.B. Pardee. 1994. Ras activation of genes: *Mob-1* as a model. *Proc. Natl. Acad. Sci.* **91**: 12515–12519.
- Lin, Y.-R., K.F. Schertz, and A.H. Paterson. 1995. Comparative analysis of QTLs affecting plant height and maturity across the Poaceae, in reference to an interspecific sorghum population. *Genetics* **141**: 391–411.
- Lopez-Fernandez, L.A. and J. Mazo. 1993. Construction of subtractive cDNA libraries from limited amounts of mRNA and multiple cycles of subtraction. *BioTechniques* **15**: 654–659.
- Lovett, M., J. Kere, and L.M. Hinton. 1991. Direct selection: A method for the isolation of cDNAs encoded by large genomic regions. *Proc. Natl. Acad. Sci.* **88**: 9628–9632.

QTL MAPPING: PROGRESS AND PROSPECTS

- Luo, Z.W. and M.J. Kearsey. 1991. Maximum likelihood estimation of linkage between a marker gene and a quantitative trait locus II. Application to backcross and doubled haploid populations. *Heredity* **66**: 117–124.
- . 1992. Interval mapping of quantitative trait loci in an F₂ population. *Heredity* **69**: 236–242.
- Mackinnon, M.J. and M. Georges. 1992. The effects of selection of linkage analysis for quantitative traits. *Genetics* **132**: 1177–1185.
- Mangin, B., B. Goffinet, and A. Rebai. 1994. Constructing confidence intervals for QTL location. *Genetics* **138**: 1301–1308.
- Markert, C.L. and F. Moller. 1959. Multiple forms of enzymes: Tissue, ontogenetic, and species-specific patterns. *Proc. Natl. Acad. Sci.* **45**: 753–763.
- Martin, B., J. Nienhuis, G. King, and A. Schaefer. 1989. Restriction fragment length polymorphisms associated with water use efficiency in tomato. *Science* **243**: 1725–1728.
- Mather, K.P. and J.L. Jinks. 1982. *Biometrical genetics*, 3rd ed. Chapman and Hall, London, UK.
- Moreno-Gonzalez, J. 1993. Efficiency of generations for estimating marker-associated QTL effects by multiple regression. *Genetics* **135**: 223–231.
- Morgan, J.G., G.M. Dolganov, S.E. Robbins, L.M. Hinton, and M. Lovett. 1992. *Nucleic Acids Res.* **20**: 5173–5179.
- Neiderheiser, J.M., R. Plomin, and G.E. McClearn. 1992. The use of CXB recombinant inbred mice to detect quantitative trait loci in behavior. *Physiol. Behav.* **52**: 429–439.
- Newman, T., F.J. deBruijn, P. Green, K. Keegstra, H. Kende, L. McIntosh, J. Ohlrogge, N. Raikhel, S. Somerville, M. Thomashow, E. Retzel, and C. Somerville. 1994. Genes galore: A summary of methods for accessing results from large-scale partial sequencing of anonymous *Arabidopsis* cDNA clones. *Plant Physiol.* **106**: 1241–1255.
- Nienhuis, J., T. Helentjaris, M. Slocum, B. Ruggero, and A. Schaefer. 1987. Restriction fragment length polymorphism analysis of loci associated with insect resistance in tomato. *Crop Sci.* **27**: 797–803.
- O'Brien, S.J., J.E. Womack, L.A. Lyons, K.J. Moore, N.A. Jenkins, and N.G. Copeland. 1993. Anchored reference loci for comparative genome mapping in mammals. *Nature Genet.* **3**: 103–112.
- Pachnis, V., L. Pevny, R. Rothstein, and F. Costantini. 1990. Transformation of yeast artificial chromosomes carrying human DNA from *Saccharomyces cerevisiae* into mammalian cells. *Proc. Natl. Acad. Sci.* **87**: 5109–5113.
- Parimoo, S., S.R. Patanjali, H. Shukla, D.D. Chaplin, and S.M. Weissman. 1991. cDNA selection: Efficient PCR approach for the selection of cDNAs encoded in large chromosomal DNA fragments. *Proc. Natl. Acad. Sci.* **88**: 9623–9627.
- Paterson, A.H., E.S. Lander, J.D. Hewitt, S. Peterson, S.E. Lincoln, and S.D. Tanksley. 1988. Resolution of quantitative traits into Mendelian factors by using a complete map of restriction fragment length polymorphisms. *Nature* **335**: 721–726.
- Paterson, A.H., J.W. Deverna, B. Lanini, and S.D. Tanksley. 1990. Fine mapping of quantitative trait loci using selected overlapping recombinant chromosomes, in an interspecies cross of tomato. *Genetics* **124**: 735–742.
- Paterson, A.H., S. Damon, J.D. Hewitt, D. Zamir, H.D. Rabinowitch, S.E. Lincoln, E.S. Lander, and S.D. Tanksley. 1991a. Mendelian factors underlying quantitative traits in tomato: Comparison across species, generations, and environments. *Genetics* **127**: 181–197.
- Paterson, A.H., S.D. Tanksley, and M.E. Sorrells. 1991b. DNA markers in crop improvement. In *Advances in agronomy* (ed. D.L. Sparks), pp. 39–90. Academic Press, New York.
- Paterson A.H., K.F. Schertz, Y.-R. Lin, S.-C. Liu, and Y.-L. Chang. 1995a. The weediness of wild plants: Molecular analysis of genes influencing dispersal and persistence of johnsongrass, *Sorghum halepense* (L.) Pers. *Proc. Natl. Acad. Sci.* **92**: 6127–6131.
- Paterson, A.H., Y.R. Lin, Z. Li, K.F. Schertz, J.F. Doebley, S.R.M. Pinson, S.C. Liu, J.W. Stansel, and J.E. Irvine. 1995b. Convergent domestication of cereal crops by independent mutations at corresponding genetic loci. *Science* **269**: 1714–1718.
- Pavan, W.J., P. Hieter, and R.H. Reeves. 1990. Modification and transfer into an embryonal carcinoma cell line of a 360-kilobase human-derived yeast artificial chromosome. *Mol. Cell. Biol.* **10**: 4163–4169.
- Pelleymounter, M.A., M.J. Cullen, M.B. Baker, R. Hecht, D. Winters, T. Boone, and F. Collins. 1995. Effects of the obese gene product on body weight regulation in ob/ob mice. *Science* **269**: 540–543.
- Pooni, H.S., D.J. Coombs, and P.S. Jinks. 1987. Detection of epistasis and linkage of interacting genes in the presence of reciprocal differences. *Heredity* **58**: 257–266.
- Plomin, R., M.J. Owen, and P. McGuffin. 1994. The genetic basis of complex human behaviors. *Science* **264**: 1733–1739.
- Quock, R.M., J.L. Mueller, L.K. Vaughn, and J.K. Bellnap. 1994. Nitrous oxide (N₂O) antinociception in BXD recombinant inbred (RI) mouse strains and identification of quantitative trait loci (QTL). *FASEB J.* **8**: A628.
- Rapp, J.P., S. Wang, and H. Dene. 1989. A genetic

PATERSON

- polymorphism in the renin gene of Dahl rats co-segregates with blood pressure. *Science* **243**: 542–544.
- Rebai, A., B. Goffinet, and B. Mangin. 1994. Approximate thresholds of interval mapping tests for QTL detection. *Genetics* **138**: 235–240.
- . 1995. Comparing power of different methods for QTL detection. *Biometrics* **51**: 87–99.
- Reiter, R.S., J.G. Coors, M.R. Sussman, and W.H. Gabelman. 1991. Genetic analysis of tolerance to low-phosphorus stress in maize using restriction fragment length polymorphisms. *Theor. Appl. Genet.* **82**: 561–568.
- Robertson, D.S. 1985. A possible technique for isolating genic DNA for quantitative traits in plants. *J. Theor. Biol.* **117**: 1–10.
- Rodolphe, F. and M. Lefort. 1993. A multi-marker model for detecting chromosomal segments displaying QTL activity. *Genetics* **134**: 1277–1288.
- Sari-Gorla, M., M.E. Pe, D.L. Mulcahy, and E. Ottaviano. 1992. Genetic dissection of pollen competitive ability in maize. *Heredity* **69**: 423–430.
- Sasaki, T., J. Song, Y. Koga-bon, E. Matsui, F. Fang, H. Higo, H. Nagasaki, M. Hori, M. Miya, E. Murayama-Kayano, et al. 1994. Toward cataloging all rice genes: Large-scale sequencing of randomly-chosen rice cDNAs from a callus cDNA library. *Plant J.* **6**: 615–624.
- Sato, S. and I. Sakamoto. 1983. Inheritance of heading time in isogenic line rice cultivar, Taichung 65 carrying earliness genes from a reciprocal translocation homozygote, T3-7. *Jpn. J. Breed.* **33**: 118–119.
- Sauer, B. and N. Henderson. 1988. Site-specific DNA recombination in mammalian cells by the cre recombinase of bacteriophage P1. *Proc. Natl. Acad. Sci.* **85**: 5166–5170.
- Sax, K. 1923. The association of size differences with seed-coat pattern and pigmentation in *Phaseolus vulgaris*. *Genetics* **8**: 552–560.
- Schon, C.C., A.E. Melchinger, J. Boppenmaier, E. Brunklaus-Jung, R.G. Herrmann, and J.F. Seitzer. 1994. RFLP mapping in maize: Quantitative trait loci affecting testcross performance of elite European flint lines. *Crop Sci.* **34**: 378–389.
- Severson, D.W., V. Thathy, A. Mori, Y. Zhang, and B.M. Christensen. 1995. Restriction fragment length polymorphism mapping of quantitative trait loci for malaria parasite susceptibility in the mosquito *Aedes aegypti*. *Genetics* **139**: 1711–1717.
- Shields, R. 1993. Pastoral synteny. *Nature* **365**: 297–298.
- Shizuya, H., B. Birren, U.-J. Kim, V. Mancino, T. Slepak, Y. Tachiiri, and M. Simon. 1992. Cloning and stable maintenance of 300-kilobase-pair fragments of human DNA in *Escherichia coli* using an F-factor-based vector. *Proc. Natl. Acad. Sci.* **89**: 8794–8797.
- Soller, M., T. Brody, and A. Genizi. 1979. The expected distribution of marker-linked quantitative effects in crosses between inbred lines. *Heredity* **43**: 179–190.
- Spassky, B., T. Dobzhansky, and W.W. Anderson. 1965. Genetics of natural populations. XXXVI. Epistatic interactions of the components of the genetic load in *Drosophila pseudoobscura*. *Genetics* **52**: 653–664.
- Spickett, S.G. and J.M. Thoday. 1966. Regular response to selection 3. Interaction between located polygenes. *Genet. Res.* **7**: 96–121.
- Steinmetz, M., M. Minard, S. Horvath, J. McNicholas, J. Srelinger, C. Wake, E. Long, B. Mach, and L. Hood. 1982. A molecular map of the immune response region from the major histocompatibility complex of the mouse. *Nature* **300**: 35–42.
- Strauss, W.M. and R. Jaenisch. 1992. Molecular complementation of a collagen mutation in mammalian cells using yeast artificial chromosomes. *EMBO J.* **11**: 417–422.
- Stuber, C.W., M.D. Edwards, and J.F. Wendel. 1987. Molecular-marker-facilitated investigations of quantitative-trait loci in maize. II. Factors influencing yield and its component traits. *Crop Sci.* **27**: 639–648.
- Stuber, C.W., S.E. Lincoln, D.W. Wolff, T. Helentjaris, and E.S. Lander. 1992. Identification of genetic factors contributing to heterosis in a hybrid from two elite inbred lines using molecular markers. *Genetics* **132**: 823–839.
- Sun, T., H.M. Goodman, and F.M. Ausubel. 1992. Cloning the *Arabidopsis* GA1 locus by genomic subtraction. *Plant Cell* **4**: 119–128.
- Suzuki, D.T., A.J.F. Griffiths, and R.C. Lewontin. 1986. *An introduction to genetic analysis*, 3rd ed. W.H. Freeman and Company, New York.
- Tanksley, S.D. and J.D. Hewitt. 1988. Use of molecular markers in breeding for soluble solids in tomato: A re-examination. *Theor. Appl. Genet.* **75**: 811–823.
- Templeton, A.R. 1979. The unit of selection in *Drosophila mercatorum*. II. Genetic evolution and the origin of coadapted genomes in parthenogenetic strains. *Genetics* **92**: 1265–1282.
- . 1980. The theory of speciation via the founder principle. *Genetics* **94**: 1011–1038.
- Teutonico, R.A. and T.C. Osborn. 1994. Mapping of RFLP and qualitative trait loci in *Brassica rapa* and comparison to the linkage maps of *B. napus*, *B. oleracea*, and *Arabidopsis thaliana*. *Theor. Appl. Genet.* **89**: 885–894.

QTL MAPPING: PROGRESS AND PROSPECTS

Thoday, J.M. 1961. Location of polygenes. *Nature* **191**: 368–370.

Van Eck, H.J., J.M.E. Jacobs, P. Stam, J. Ton, W.J. Stiekema, and E. Jacobsen. 1994. Multiple alleles for tuber shape in diploid potato detected by qualitative and quantitative genetic analysis using RFLPs. *Genetics* **137**: 303–309.

Wang, G., D.J. MacKill, J.M. Bonman, S.R. McCouch, M.C. Champoux, and R.J. Nelson. 1994. RFLP mapping of genes conferring complete and partial resistance to blast in a durably resistant rice cultivar. *Genetics* **136**: 1421–1434.

Weller, J.I. 1986. Maximum likelihood techniques for the mapping and analysis of quantitative trait loci with the aid of genetic markers. *Biometrics* **42**: 627–640.

Weller, J.I., M. Soller, and T. Brody. 1988. Linkage analysis of quantitative traits in an interspecific cross of tomato, *Lycopersicon esculentum* × *Lycopersicon pimpinellifolium*, by means of genetic markers. *Genetics* **118**: 329–340.

Wright, S. 1932. The roles of mutation, inbreeding, crossbreeding, and selection in evolution. *Proc. VI Intl. Congr. Genet.* **1**: 356–366.

———. 1951. The genetic structure of populations. *Ann. Eugenics* **15**: 323–354.

Xiao, J., J. Li, L. Yuan, and S.D. Tanksley. 1995. Dominance is the major genetic basis of heterosis in rice as revealed by QTL analysis using molecular markers. *Genetics* **140**: 745–754.

Zeng, Z.-B. 1993. Theoretical basis for separation of multiple linked gene effects in mapping quantitative trait loci. *Proc. Natl. Acad. Sci.* **90**: 10972–10976.

———. 1994. Precision mapping of quantitative trait loci. *Genetics* **136**: 1457–1468.

Received September 18, 1995; accepted in revised form October 26, 1995.

Molecular dissection of quantitative traits: progress and prospects.

A H Paterson

Genome Res. 1995 5: 321-333

Access the most recent version at doi:[10.1101/gr.5.4.321](https://doi.org/10.1101/gr.5.4.321)

References

This article cites 133 articles, 72 of which can be accessed free at:
<http://genome.cshlp.org/content/5/4/321.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

**Affordable, Accurate
Sequencing.**

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>