LETTERS

An Integrated Genetic Map of the Pearl Locus of Mouse Chromosome 13

Albert B. Seymour,^{1,2} Brenda L. Yanak,² Edward P. O'Brien,³
Michael E. Rusiniak,³ Edward K. Novak,³ Larry H. Pinto,⁴
Richard T. Swank,³ and Michael B. Gorin^{1,2,5}

¹Departments of Human Genetics, and ²Ophthalmology, University of Pittsburgh, Pittsburgh, Pennsylvania 15213; ³Molecular and Cellular Biology Department, Roswell Park Cancer Institute, Buffalo, New York 14263; ⁴Neurobiology Department, Northwestern University, Evanston, Illinois 60201

We have used a *Mus domesticus/spretus* congenic animal and two interspecific backcross panels to map genetically 30 sequence-tagged sites (STSs) and 13 genes to the vicinity of the pearl locus on mouse chromosome 13. The STSs defining the mapped region are from DI3Mit9 to DI3Mit37, spanning 10.6 cM. Genes mapped to this region include Versican (*Cspg2*), GTPase activating protein (*Rasa*), dihydrofolate reductase (*Dhfr*), arylsulfatase (*As-1*), thrombin receptor (*Cf2r*), hexosaminidase b (*Hexb*), 3-hydroxy-3-methylglutaryl coenzyme A reductase (*Hmgcr*), microtubule associated protein 5/1b (*Mtap5*), phosphodiesterase (*Pde*), phosphatidylinositol 3' kinase (*Pik3rl*), rat integrin α -subunit (*Itgal*), collagen receptor α 2-subunit (*Itga2*), and 5-hydroxytryptamine 1a receptor (*Htr1a*). This high resolution genetic map of the pearl region of chromosome 13 establishes the order of multiple markers, including genes whose human homologs are located within a limited region of human chromosome 5, with respect to the phenotypic anchor marker pearl.

The mouse pearl mutation is a recessive, hypopigmentation mutation that affects retinal development and function, and causes inherited platelet storage pool deficiencies (Balkema et al. 1983; Linden and Pinto 1985; O'Brien et al. 1995). Pearl has been studied as a model of human inherited congenital stationary night blindness and the inherited human Hermansky-Pudlak Syndrome. Originally identified in 1954 in a C3H strain, it has subsequently been transferred into a C57Bl/6J line, C57Bl/6J PIN *pe/pe* (Avner et al. 1988). The pearl locus was first localized to chromosome 13 by linkage to *Arylsulfatase B* and *Lth1* (Elliott et al. 1985). Because of its heritable visual phenotype, pearl served as an anchor locus for other genes and markers that have been mapped to chromosome 13 (Holcombe et al. 1991; O'Brien et al. 1995; Xu et al. 1996).

In an effort to further localize the pearl locus, a congenic strain, B6/*spretus* *pe*^{PIN}N12F6, was

developed to introduce a highly polymorphic region suitable for the mapping of new markers to the pearl region (Rikke et al. 1993). The selective breeding of a region of *Mus spretus* containing the pearl locus into a C57Bl/6J background allowed for the binning of genes and other markers to within or outside a 10.0-cM interval that contains the pearl locus.

Interspecific backcross panels have been developed to allow high resolution recombinant mapping (Copeland and Jenkins 1991). Markers that are in the pearl region have been localized partly by using a 96-mouse interspecific backcross panel of C57Bl/6J and *M. spretus* from Jackson Laboratories. A 528-mouse backcross panel segregating pearl, muted, and satin phenotypes mapped pearl to within a 1.4-cM region (O'Brien et al. 1995). This panel has subsequently been propagated to 1250 mice and the pearl region is now localized to within 0.5 cM.

We have integrated mapping data from these sources as part of our efforts to define the critical pearl region with respect to the closest flanking markers. This integrated map establishes the or-

⁵Corresponding author.

E-MAIL Gorin@vision.eei.upmc.edu; FAX (412) 647-5880.

MAP OF THE PEARL LOCUS ON MOUSE CHROMOSOME 13

der of multiple genes and microsatellite markers on chromosome 13.

RESULTS

The congenic B6/*spretus* pe^{+PIN} N12F6 mouse strain contains a 10-cM genomic region of *M. spretus* DNA surrounding the pearl locus with a small flanking region of C3H genomic DNA from the original pearl strain. The remainder of its genome is derived from C57Bl/6J. Forty microsatellites and sequence-tagged sites (STSs) with nine expressed genes were typed in *M. spretus* C57Bl/6J and C3H strains to determine the boundaries of the *spretus* genomic region containing the pearl gene. STS markers PL2, PL3, D13SH1, D13Gor2, D13Gor3, D13Gor4, D13Mit27, D13Mit28, D13Mit29, D13Mit105, D13Mit106, D13Mit107, D13Mit160, D13Mit161, D13Mit169, D13Mit258, D13Mit104, D13Mit69, D13Mit145, D13Mit191, D13Mit36, D5S39 (human), and mouse genes *Dhfr*, *Hexb*, *As-1*, *Hmgcr*, *Rasa*, and *Mtap5* were binned within the 10-cM interval. *Htr1a* and *Ctla3* were binned telomeric to the 10-cM interval.

Southern hybridization with the probe for *Srd5 alpha* revealed a 1.7-kb informative allele in the congenic that matched the C3H parental strain. We were unable to distinguish whether the position of *Srd5 alpha*, based on the Southern blot data, was centromeric or telomeric to the *M. spretus* region. We have placed it in the proximal C3H region based on the 1993 and 1994 consensus maps (Justice and Stephenson 1993, 1994) and a previous study that localized *Srd5 alpha* centromeric with respect to *As-1* and *Rasa* (Jenkins et al. 1991), which we have binned into the *M. spretus* region.

The strain of origin of most alleles could be determined by directly comparing the products of the appropriate parental DNAs. D13Mit36, however, identified an allele in the congenic B6/*spretus* pe^{+PIN} N12F6 mouse that could either be attributed to *M. spretus* or C57Bl/6J. Examination of the locus in the original C57Bl/6J PIN pe/pe mouse showed that the D13Mit36 allele corresponded to the C3H allele rather than to the C57Bl/6J allele. We concluded that the observed allele for D13Mit36 in the congenic was specific for *M. spretus*. In contrast, D13Mit37, which is telomeric to D13Mit36, identified a C3H allele in both the congenic and C57Bl/6J PIN pe/pe . Taken together, this indicates that the boundary of the *M. spretus* region in this con-

genic strain must lie between D13Mit36 and D13Mit37.

We mapped *Dhfr*, *Pik3r1*, and several microsatellites onto the Jackson [(C57Bl/6J $\times$ *spretus*) F_1 $\times$ *M. spretus*] (BSS) interspecific backcross panel. These data are shown in Figure 1. The marker positions were calculated using Map manager v. 2.6. Marker order was determined by minimizing the number of multiple recombination events. *Dhfr* showed a band shift when analyzed by single-strand conformation polymorphism (SSCP); C57Bl/6J revealed a higher mobility band than *M. spretus*. *Pik3r1* revealed a T-C polymorphism at nucleotide 2833 in the untranslated region between C57Bl/6J and *M. spretus*, respectively. This single-base polymorphism was not consistently informative by SSCP analysis, therefore a PCR-based sequence analysis was performed to determine its segregation throughout the backcross panels. The resolution of this panel is 7.9 cM at the 95% confidence level.

We further localized the pearl region with the interspecific backcross panel described by O'Brien et al. (1995). This panel, originally de-

Figure 1 Genetic map of mouse chromosome 13 configured by Jackson Laboratories, showing the positions and relative distances of STS markers and two genes in the pearl region. Highlighted markers were mapped in our laboratory. Numbers at left indicate distances in centimorgans. Information regarding these markers can be obtained from Jackson Laboratories.

MAP OF THE PEARL LOCUS ON MOUSE CHROMOSOME 13

1994) to 1.8 cM telomeric to the set of markers nonrecombinant with the pearl locus. D13Mit28 and D13Mit29 are nonrecombinant with each other in this panel and D13Mit105 and D13Mit106 are 0.1 cM and 0.2 cM centromeric to D13Mit28/29, respectively. D13Mit104 has been placed telomeric to D13Mit105 by 0.4 cM and telomeric to D13Mit106 by 0.6 cM. The positions of D13Mit145, D13Mit144, and D13Mit109 have been localized to 0.3, 0.6, and 1.2 cM, respectively, telomeric to D13Mit161. In the Jackson backcross panel we identified a recombination between D13Mit160 and the pearl locus. A similar recombination event has not been observed in the 1250-mouse interspecific backcross panel. Because of the higher resolution of this panel, we have placed D13Mit160 centromeric but closer to D13Mit258, D13Mit104, D13Mit161, D13Mit169, and D13Mit108.

Many of the genes that we placed on this map were potential candidate genes for the pearl mutation, specifically *Cspg2*, *Rasa*, *Itga2*, and *Pik3r1*. The mapping results have eliminated these, as well as the other mapped genes, as candidate genes based on recombination events from the pearl locus.

All of the genes that we have placed in the region of the pearl locus on mouse chromosome 13 have been associated with a limited region of human chromosome 5. The orders of these genes and their respective genetic distances on the human genome have not been established. Although it is likely that this region of mouse chromosome 13 is syntenic with human 5q13, further human genome physical mapping efforts will be necessary to establish if small rearrangements distinguish these regions of the two genomes.

The human spinal muscular atrophy (SMA) locus is telomeric to the syntenic pearl region of mouse chromosome 13 based on the human chromosome 5 mapping data. MAP1B, the human homolog of *Mtap5*, which shows linkage to the human SMA locus is 1.6 cM distal to the pearl region. It has been reported that this region of the genome is very unstable with regards to genomic rearrangements (Selig et al. 1995). One pearl lineage has been reported to have a spontaneous reversion rate of 0.5% (Pinto et al. 1985), suggesting that, at least for one pearl allele, the disease phenotype could be caused by a DNA insertion. Elucidation of the actual pearl gene and the characterization of the different pearl mutations may clarify this potential genomic instability with respect to this gene.

The high resolution mapping and the abundance of markers in this region allow the initiation of physical mapping techniques to identify the gene responsible for the pearl phenotype as well as providing a set of markers that can be used in the mapping of other genes to this region of mouse chromosome 13.

METHODS

The interspecific backcross panel of PWK and SaMuPe has been described previously (O'Brien et al. 1995). The pearl (pearl $PIN^{pe/pe}$) and congenic B6/*spretus* pe^{+PIN} N12F6 have been described previously (Avner et al. 1988; Rikke et al. 1993). The C3H, C57Bl/6J mice and the 96-mouse backcross panel of BSS, were purchased from Jackson Laboratories.

Genomic DNA was isolated from mouse spleen and kidney according to standard protocols (Sambrook et al. 1989; Taylor and Rowe 1989). For Southern blotting, 10 μ g of mouse genomic DNA was digested with *Eco*RI, *Pst*I, *Bam*HI, *Bgl*II, or *Hind*III according to manufacturer's conditions. The digests were electrophoresed on 1.0% agarose (FMC), stained with EtBr, and transferred to Hybond N+ filters (Amersham). The DNA was immobilized at 80°C for 2 hr. The hybridizations were performed as described previously (Xu et al. 1996). Markers were determined to be within the pearl-containing *M. spretus* region of the congenic animal if the hybridization patterns of the congenic animal were identical to those of *M. spretus* and distinct from the C57Bl/6J background.

Candidate genes that have been mapped previously to mouse chromosome 13 or localized to a presumed syntenic region of human chromosome 5 were used to determine their position with respect to the pearl locus. The probes are described in Table 1. All probes were excised from the vector with the appropriate enzyme, separated by gel electrophoresis on 1.0% agarose, excised, and purified with GeneClean (Bio 101, Inc.). The DNA Probes were labeled by the random hexamer method with [α - 32 P]dCTP (Feinberg and Vogelstein 1983; NEN/Dupont).

D13Mit series primer sets were purchased from Research Genetics. The forward primer for each set was end-labeled with T4 kinase (GIBCO) and [γ - 32 P]ATP (NEN/Dupont). The PCR conditions were 94°C for 2 min followed by 35 cycles of 94°C for 30 sec, 55°C for 1 min, and 72°C for 1 min with 1 unit of *Taq* polymerase in 25 mM TAPS (pH 9.3), 1.5 mM $MgCl_2$, 50 mM KCl, 1 mM DTT, and 0.2 mM each of dATP, dCTP, dGTP, and dTTP in a final volume of 20 μ l. Products were electrophoresed on a 6.0% polyacrylamide, 7 M urea denaturing sequencing gel, dried, and analyzed by autoradiography.

The D13Gor primers were described previously (Xu et al. 1996). The *Pik3r1* primers are specific for nucleotides 2752–2935 in the 3' untranslated region of phosphatidylinositol 3' kinase regulatory subunit resulting in a product of 184 bp. The primer sequences are as follows. *Pik3r1F* 5'-TTCCTCACCTTCAAGCCACCCAAG-3'; *Pik3r1R* 5'-AGGTTAGAAACGTCTGGTCATCCAAC-3'. The *Dhfr* primers were OD3PF 5'-ACAGGAAGATGCTTTCAAGTTC-3' and OD3PR 5'-GAGGTCTCATGGGAGGGG-3'. *Thrombin receptor* (*Cf2r*), D13SH1, PL2, and PL3 primer sequences

Table 1. Genes Mapped to the Pearl Region of Mouse Chromosome 13

Gene	Probe name	Enzyme ^a		Informative alleles ^d		Reference
		congenic ^b	PWK ^c	congenic	PWK	
Versican	Cspg2	N.D. ^e	<i>KpnI</i>	N.D.	14.5, 9.4	Zimmermann and Ruoslahti (1988)
GTPase-activating protein	Rasa	<i>EcoRI</i>	<i>HindIII</i>	19.1	4.8	Hsieh et al. (1989)
Arylsulfatase B	As1	<i>BamHI</i>	<i>EcoRI</i>	12.3	22, 9.0, 7.5	Schuchman et al. (1990)
Hexosaminidase B	HexB	<i>BamHI</i>	<i>PstI</i>	16.4, 5.4	4.5	Bapat et al. (1988)
3-hydroxy-3-methylglutaryl coenzyme A reductase	Hmgcr	N.D.	<i>KpnI</i>	N.D.	6.4	Sundaresan et al. (1989)
Microtubule-associated protein 5/1B	Mtap5	<i>EcoRI</i>	<i>XbaI</i>	4.5	9.6	Garner et al. (1990)
Cyclic nucleotide phosphodiesterase	Pde	N.D.	<i>HindIII</i>	N.D.	12.0	Milatovich et al. (1994)
Collagen receptor $\alpha 2$ subunit	Itga2	N.D.	<i>BglII</i>	N.D.	6.0	Takada and Hemler (1989)
Rat integrin $\alpha 1$ -subunit	Itga1	N.D.	<i>PstI</i>	N.D.	9.6	Ignatius et al. (1990)
5-hydroxytryptamine (serotonin) 1a receptor	Htr1a	<i>EcoRI</i>	<i>BglII</i>	1.2	4.9	Kobilka et al. (1987)
Steroid 5 α -reductase	Srd5alpha	<i>BglII</i>	N.D.	1.7	N.D.	Jenkins et al. (1991)
Cytotoxic T lymphocyte-associated sequence 3	Ctla3	<i>EcoRI</i>	N.D.	24.5, 13.5	N.D.	Gershenfeld and Weismann (1986)

^aEnzyme used to digest genomic DNA to identify polymorphism.^bGenomic DNA from the congenic strain was compared to that of C57 and Spretus as described in the text.^cGenomic DNA from PWK was compared to SaMuPe to observe crossovers in the backcross panel.^dAllele sizes in kilobases (kb).^e(N.D.) Not determined.

have been described previously (Casavant and Hardies 1993; Rikke et al. 1993; Poirier et al. 1996). The PCR conditions for *Dhfr* were identical to those for the D13Mit series. The PCR conditions for *Pik3r1* required an annealing temperature of 60°C.

SSCP analysis for *Dhfr* was done per standard protocol with a few modifications (Dracopoli et al. 1994). Samples were denatured in 98% formamide/0.025% bromophenol blue/0.025% xylene cyanol and 0.65 mM NaOH at 95°C for 5 min and electrophoresed on a 0.5 × MDE gel (Hydrolink) overnight at 4.0°C. The gel was dried and analyzed by autoradiography.

DNA sequencing of PCR products was performed using a cycle sequencing kit (Epicentre). Primers were end-labeled as described above. The products were separated on a 6.0% 7 M urea denaturing sequencing gel, dried, and analyzed by autoradiography.

ACKNOWLEDGMENTS

We thank Drs. Lucy Rowe and Mary Barter at Jackson

Laboratory for assistance in the mapping on the Jackson backcross panel, and we are grateful to Dr. Stephen C. Hardies at The University of Texas Health Science Center for providing *Dhfr* primers (OD3PF, OD3PR) and critically reading this manuscript. This work was supported by National Institutes of Health (NIH) grant EY9192, National Eye Institute Core grant EYO8098, and Eye and Ear Foundation of Pittsburgh (M.B.G.), NIH grant EY01221 (L.H.P.) and National Heart, Lung, and Blood Institute grant HL31698 (R.T.S.).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

REFERENCES

- Avner, P., L. Amar, L. Dandolo, and J.L. Cuenet. 1988. Genetic analysis of the mouse using interspecific crosses. *Trends Genet. Rev.* **4**: 18–23.

MAP OF THE PEARL LOCUS ON MOUSE CHROMOSOME 13

- Balkema, G.W., N.J. Mangini, and L.H. Pinto. 1983. Discrete visual defects in pearl mutant mice. *Science* **219**: 1085–1087.
- Bapat, B., M. Ethier, K. Neote, D. Mahuran, and R.A. Gravel. 1988. Cloning and sequence analysis of a cDNA encoding the beta-subunit of mouse beta-hexosaminidase. *FEBS Lett.* **237**: 191–195.
- Casavant, N.C. and S.C. Hardies. 1993. Targeted cloning of a subfamily of LINE-1 elements by subfamily-specific LINE-1-PCR. *Mamm. Genome* **4**: 193–201.
- Copeland, N.G. and N.A. Jenkins. 1991. Development and applications of a molecular genetic linkage map of the mouse genome. *Trends Genet.* **7**: 113–118.
- Drapopoli, N.C., J.L. Haines, B.R. Korf, D.T. Moir, C.C. Morton, C.E. Seidman, J.G. Seidman, and D.R. Smith, eds. 1994. Searching candidate genes for mutations. In *Current protocols in human genetics*. Chap. 7, pp. 7.4.1–7.4.6. John Wiley/Greene, New York, NY.
- Elliott, R.W., W.L. Daniel, B.A. Taylor, and E.K. Novak. 1985. Linkage of loci affecting a murine liver protein and arylsulfatase B to chromosome 13. *J. Heredity* **76**: 243–246.
- Feinberg, A.P. and B. Vogelstein. 1983. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* **132**: 6–13.
- Garner, C.C., A. Garner, G. Huber, C. Kazak, and C. Matus. 1990. Molecular cloning of microtubule-associated protein 1 (MAP1A) and microtubule-associated protein 5 (MAP1B): Identification of distinct genes and their differential expression in developing brain. *J. Neurochem.* **55**: 146–154.
- Gershenfeld, H.K. and I.L. Weissman. 1986. Cloning of a cDNA for a T cell-specific serine protease from a cytotoxic T lymphocyte. *Science* **232**: 854–858.
- Holcombe, R.F., D.A. Stephenson, A. Zweidler, R.M. Stewart, V.M. Chapman, and J.G. Seidman. 1991. Linkage of loci associated with two pigment mutations on mouse chromosome 13. *Genet. Res. Camb.* **58**: 41–50.
- Hoyle, J., I.G. Yulug, S.E. Egan, and E.M.C. Fisher. 1994. The gene that encodes the phosphatidylinositol-3 kinase regulatory subunit (p85 alpha) maps to chromosome 13 in the mouse. *Genomics* **24**: 400–402.
- Hsieh, C.-L., U.S. Vogel, R.A.F. Dixon, and U. Francke. 1989. Chromosome localization and cDNA sequence of murine and human genes for ras p21 GTPase activating protein (GAP). *Somat. Cell & Mol. Genet.* **15**: 579–590.
- Ignatius, M.J., T.H. Large, M. Houde, J.W. Tawil, A. Barton, F. Esch, S. Carbonetto, and L.F. Reichardt. 1990. Molecular cloning of the rat integrin α_1 -subunit: A receptor for laminin and collagen. *J. Cell Biol.* **111**: 709–720.
- Jenkins, E.P., C.-L. Hsieh, A. Milatovich, K. Normington, D.M. Berman, U. Francke, and D.W. Russell. 1991. Characterization and chromosomal mapping of a human steroid 5 alpha-reductase gene and pseudogene and mapping of the mouse homolog. *Genomics* **11**: 1102–1112.
- Justice, M.J. and D.A. Stephenson. 1994. Mouse chromosome 13. *Mamm. Genome* **5**: S196–S206.
- . 1993. Mouse chromosome 13. *Mamm. Genome* **4**: S192–S202.
- Kobilka, B.K., T. Friele, S. Collins, T. Yang-Feng, T.S. Kobilka, U. Francke, R.J. Lefkowitz, and M.G. Caron. 1987. An intronless gene encoding a potential member of the family of receptors coupled to guanine nucleotide regulatory proteins. *Nature* **329**: 75–79.
- Linden, R. and L.H. Pinto. 1985. Developmental genetics of the retina: Evidence that the pearl mutation in the mouse affects the time course of natural cell death in the ganglion cell layer. *Exp. Brain Res.* **60**: 79–86.
- Milatovich, A., G. Bolger, T. Michaeli, and U. Francke. 1994. Chromosome localizations of genes for five cAMP-specific phosphodiesterases in man and mouse. *Somat. Cell & Mol. Genet.* **20**: 75–86.
- O'Brien, E., E. Novak, L. Zhen, K. Manly, D. Stephenson, and R. Swank. 1995. Molecular markers near two mouse chromosome 13 genes, muted and pearl, which cause platelet storage pool deficiency. *Mamm. Genome* **6**: 19–24.
- Pinto, L.H., M.A. Williams, H. Suzuki, N. Mangini, G.W. Balkema Jr., and J.W. Vanable Jr. 1985. Visual defects in mouse hypopigmentation mutants. *Neurosci. Res.* **S2**: S239–S251.
- Poirier, C., E.P. O'Brien, A.L. Brunialti, J.C. Chambard, R.T. Swank, and J.L. Guenet. 1996. The gene encoding thrombin receptor (Cf2r) maps to mouse chromosome 13. *Mamm. Genome* **7**: 322.
- Rikke, B.A., L.H. Pinto, M.B. Gorin, and S.C. Hardies. 1993. *M. spretus*-specific LINE-1 DNA probes applied to the cloning of the murine pearl locus. *Genomics* **15**: 291–296.
- Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: A laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Schuchman, E.H., C.E. Jackson, and R.J. Desnick. 1990. Human arylsulfatase B: MOPAC cloning, nucleotide sequence of a full-length cDNA, and regions of amino acid identity with arylsulfatases A and C. *Genomics* **6**: 149–158.
- Selig, S., S. Bruno, J.M. Scharf, C.H. Wang, E. Vitale, T.C. Gilliam, and L.M. Kunkel. 1995. Expressed cadherin pseudogenes are localized to the critical region of the

SEYMOUR ET AL.

spinal muscular atrophy gene. *Proc. Natl. Acad. Sci.* **92**: 3702–3706.

Sundaresan, S., T.L. Yang-Feng, and U. Francke. 1989. Genes for HMG-CoA reductase and serotonin 1a receptor are on mouse chromosome 13. *Somat. Cell & Mol. Genet.* **15**: 465–469.

Takada, Y. and M.E. Hemler. 1989. The primary structure of the VLA-2/collagen receptor alpha 2 subunit (platelet GPIa): Homology to other integrins and the presence of a possible collagen-binding domain. *J. Cell Biol.* **109**: 397–407.

Taylor, B.A. and L.B. Rowe. 1984. Genes for serum amyloid A proteins map to chromosome 7 in the mouse. *Mol. Gen. Genet.* **195**: 491–499.

Xu, H.P., B.L. Yanak, M.H. Wigler, and M.B. Gorin. 1996. New polymorphic markers in the vicinity of the pearl locus on mouse chromosome 13. *Mamm. Genome* **7**: 16–19.

Zimmermann, D.R. and E. Ruoslahti. 1988. Multiple domains of the large fibroblast proteoglycan, versican. *EMBO J.* **8**: 2975–2981.

Received February 26, 1996; accepted in revised form April 16, 1996.

An integrated genetic map of the pearl locus of mouse chromosome 13.

A B Seymour, B L Yanak, E P O'Brien, et al.

Genome Res. 1996 6: 538-544

Access the most recent version at doi:[10.1101/gr.6.6.538](https://doi.org/10.1101/gr.6.6.538)

References

This article cites 30 articles, 5 of which can be accessed free at:
<http://genome.cshlp.org/content/6/6/538.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Affordable, Accurate
Sequencing.

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>