Research

A Prokaryotic Gene Cluster Involved in Synthesis of Lysine through the Amino Adipate Pathway: A Key to the Evolution of Amino Acid Biosynthesis

Hiromi Nishida,¹ Makoto Nishiyama,^{2,4} Nobuyuki Kobashi,² Takehide Kosuge,³ Takayuki Hoshino,³ and Hisakazu Yamane²

¹Institute of Molecular and Cellular Biosciences, The University of Tokyo, Tokyo 113-0032, Japan; ²Biotechnology Research Center, The University of Tokyo, Tokyo 113-8657, Japan; ³Institute of Applied Biochemistry, University of Tsukuba, Tsukuba, Ibaraki 305-8572, Japan

In previous studies we determined the nucleotide sequence of the gene cluster containing *lys20*, *hacA* (*lys4A*), *hacB* (*lys4B*), *orfE*, *orfF*, *rimK*, *argC*, and *argB* of *Thermus thermophilus*, an extremely thermophilic bacterium. In this study, we characterized the role of each gene in the cluster by gene disruption and examined auxotrophy in the disruptants. All disruptants except for the *orfE* disruption showed a lysine auxotrophic phenotype. This was surprising because this cluster consists of genes coding for unrelated proteins based on their names, which had been tentatively designated by homology analysis. Although the newly found pathway contains α -aminoadipic acid as a lysine biosynthetic intermediate, this pathway is not the same as the eukaryotic one. When each of the gene products was phylogenetically analyzed, we found that genes evolutionarily-related to the lysine biosynthetic genes in *T. thermophilus* were all present in a hyperthermophilic and anaerobic archaeon, *Pyrococcus horikoshii*, and formed a gene cluster in a manner similar to that in *T. thermophilus*. Furthermore, this gene cluster was analogous in part to the present leucine and arginine biosyntheses pathways. This lysine biosynthesis cluster is assumed to be one of the origins of lysine biosynthesis and could therefore become a key to the evolution of amino acid biosynthesis.

Lysine had been believed to be synthesized from aspartic acid through the diaminopimelic acid (DAP) pathway in all prokaryotic organisms. An extremely thermophilic Gram-negative bacterium, *Thermus thermophilus*, was shown to synthesize lysine not via DAP, but from 2-oxoglutaric acid and acetyl-CoA through the α -aminoadipic acid (AAA) pathway, (Fig. 1) which is known to be common in fungi (Kosuge and Hoshino 1998; Kobashi et al. 1999). Because higher eukaryotes other than animals synthesize lysine through DAP, the AAA pathway is a characteristic that distinguishes fungi from higher eukaryotes (Vogel 1964, 1965; Broquist 1971).

In general, prokaryotes biosynthesize not only lysine but also methionine, threonine, and isoleucine from aspartic acid. *T. thermophilus* also possesses an aspartate kinase (Kobashi et al. 1999) that catalyzes the first reaction in the DAP pathway. However, this enzyme is only used for synthesis of methionine, threonine, and probably isoleucine. In *T. thermophilus*, lysine is synthesized through the AAA pathway (Kobashi et al. 1999; Kosuge and Hoshino 1998). This was the first discovery of lysine biosynthesis via AAA in bacte-

ria. Here we report that the gene cluster for lysine biosynthesis through the AAA pathway is also found in a hyperthermophilic and anaerobic archaeon, *Pyrococcus horikoshii*, whose genome has been sequenced completely (Kawarabayashi et al. 1998), and discuss the evolution of amino acid biosynthesis.

RESULTS AND DISCUSSION

Lysine Biosynthetic Cluster in *T. thermophilus*

We characterized the role of each gene in the cluster by gene disruption (Kosuge and Hoshino 1997, 1998) and examined auxotrophy of the disruptants (Fig. 2). Because the genes in the cluster are closely arranged in the same orientation, the cluster seems to be an operon. Gene disruption by inserting a kanamycin-resistance gene could cause reduced expression of a distal gene, which might be required for lysine biosynthesis in an operon. At present, it is unclear whether or not the genes are transcribed in a single mRNA and have independent promoters. In any case, to weaken the possible polar effect, the kanamycin-resistance gene cassette and the promoter were inserted in the same orientation as the gene cluster, which may allow expression of the genes located downstream via a read-

⁴Corresponding author.
E-MAIL umanis@mail.ecc.u-tokyo.ac.jp; FAX 81-3-5841-8030.

Figure 1 Comparison of biosynthetic pathways of lysine, leucine, and arginine, and a related part in the citric acid cycle. Red arrows indicate the putative AAA pathway of *T. thermophilus* and *P. horikoshii*.

Figure 2 Map of Km-resistance gene insertion point. Each of the pUCLU and pUCLD derivatives carrying the Km-resistance gene cassette at indicated positions was used for the disruption of the chromosomal copy of *T. thermophilus*.

through transcript originated from the inserted promoter. All of the disruptants, except for those with *orfE* disruption, showed a lysine auxotrophic phenotype (Table 1). This shows that all of the genes except for *orfE* are involved in lysine biosynthesis. This was surprising because this cluster consists of genes coding for apparently unrelated proteins, as judged by their names, which were tentatively assigned by homology analysis. Considering the functions of the homologs of the gene products and auxotrophy of the disruptants, *lys20* was shown to encode homocitrate synthase, and *hacA* and *hacB* to code for large and small subunits of homoaconitase, respectively (Kosuge and Hoshino 1998; Kobashi et al. 1999). Based on auxotrophy of the disruptants and the homology of the gene products, we assume that the ribosomal protein S6 modification enzyme (RimK), *N*-acetylglutamate kinase (ArgB), and *N*-acetylglutamate 5-semialdehyde dehydrogenase (ArgC) are involved in modification of AAA, phosphotransfer to the AAA derivative, and its reduction to yield an aldehyde compound, respectively (see below). This clearly indicates that the AAA pathway for lysine biosynthesis is used in prokaryotes as well as in eukaryotes, although some modification must be present. We also found the cluster of lysine biosynthetic genes in *P.*

horikoshii but in no other organism in the international DNA/protein databases (EMBL, GenBank, and DDBJ).

Phylogenetic Analysis of Components in the AAA Pathway of *T. thermophilus*

The above results let us reconsider amino acid biosynthesis. We here phylogenetically analyze each protein encoded by the *lys20*, *hacA*, *hacB*, *rimK*, *argC*, and *argB* genes. The alignment data are available on <http://iam.u-tokyo.ac.jp/misyst/lysclust.html>.

Homocitrate Synthase of *T. thermophilus*

Homocitrate synthase catalyzes the reaction to form homocitric acid with 2-oxoglutaric acid and acetyl-CoA. This is the first reaction in the AAA pathway (Fig. 1). Homocitrate synthase of *T. thermophilus* (Lys20) is grouped with those of the yeasts *Saccharomyces cerevisiae* and *Yarrowia lipolytica* (Fig. 3A). In addition, this group is closely related to a protein, PH1727, of *P. horikoshii* and a protein, LeuA1, of a hyperthermophilic bacterium *Aquifex aeolicus* (Deckert et al. 1998). The lineage of homocitrate synthase is closely related to that of 2-isopropylmalate synthase (LeuA). LeuA is involved in the branched pathway toward leucine and catalyzes the reaction similar to that of homocitrate synthase, in which 2-isopropylmalic acid is used in place of 2-oxoglutaric acid as one of the substrates. The phylogenetic tree shows that homocitrate synthase and 2-isopropylmalate synthase share a common ancestor.

Homoaconitase of *T. thermophilus*

Homoaconitase (homoaconitate hydratase) catalyzes conversion of *cis*-homoaconitic acid to either homocitric acid or homoisocitric acid, or vice versa. The large subunit of homoaconitase (HacA/Lys4A) of *T. thermophilus* is grouped with a protein, PH1726, of *P. horikoshii* (Fig. 3B). This group is connected to the lineage of 3-isopropylmalate dehydratase (LeuC) homologs

Table 1. Effect of Gene Disruption on Growth of *T. thermophilus*

Disrupted gene	Phenotype	Growth in minimal medium containing α -amino adipate
<i>hacB</i>	lysine auxotroph	+
<i>orfE</i>	wild type	—
<i>orfF</i>	lysine auxotroph	—
<i>rimK</i>	lysine auxotroph	—
<i>pseudo-argC</i>	lysine auxotroph	—
<i>argB</i>	lysine auxotroph	—

Figure 3 Comparison of TCA cycle, AAA lysine biosynthesis, and leucine biosynthesis related proteins. (A) Phylogenetic relationships among the amino acid sequence encoded in *lys20* gene of *T. thermophilus* and high-scoring amino acid sequences from the result of homology search using PSI-BLAST (Altschul et al. 1997). A total of 332 amino acid sites were considered without gap regions in alignment. The bar indicates 10% difference of amino acid sequence. The DAD (DDBJ amino acid sequence database) accession nos. used in this phylogenetic analysis are D85684-1-1, D63999-62-1, Z46907-2-1, AE000772-5-1, U32779-6-1, AE00117-11-1, X71612-5-1, Z75208-75-1, U92974-14-1, U67561-1-1, AE000909-1-1, AE000922-3-1, U67500-4-1, AP000007-42-1, Z74230-1-1, Z74179-1-1, Z49114-1-1, AE000685-7-1, AE001091-11-1, U67579-2-1, AE000851-8-1, and AE001038-8-1. (B) Phylogenetic relationships among the amino acid sequence encoded in the *hacA* gene of *T. thermophilus* and the related sequences. A total of 279 amino acid sites were considered without gap regions in alignment. The DAD accession nos. used in this phylogenetic analysis are AP000007-41-1, AE000954-24-1, X99624-1-1, U46154-1-1, J05224-1-1, U87939-1-1, U80040-1-1, U17709-1-1, AE000756-5-1, AE000901-10-1, U67499-13-1, U67543-1-1, AE000967-2-1, AE000922-4-1, AE000714-13-1, U92974-16-1, Z99263-32-1, AL021287-2-1, AL031124-34-1, X84647-1-1, U32779-8-1, AE00117-9-1, Z99262-3-1, D67033-1-1, X53090-1-1, M33166-1-1, and D63833-1-1.

from *A. aeolicus* and three archaea, *Archaeoglobus fulgidus* (Klenk et al. 1997), *Methanobacterium thermoautotrophicum* (Smith et al. 1997) and *Methanococcus jannaschii* (Bult et al. 1996). Similar results were obtained

with the small subunit, HacB/Lys4B, of homoaconitase (data not shown).

Each of these three archaea has two 3-isopropylmalate dehydratase homologs, both of which are also related to HacA of *T. thermophilus* and PH1726 of *P. horikoshii*. This may reflect the past existence of a common origin of homoaconitase and 3-isopropylmalate dehydratase in those microorganisms.

T. thermophilus homoaconitase consists of large and small subunits, HacA and HacB, like 3-isopropylmalate dehydratase, LeuC and LeuD, whereas fungal homoaconitase is a single, long polypeptide containing sequences for both large and small subunits. The environment of fungi has been quite different from the extremely thermophilic environment of *Thermus* and *Pyrococcus*. In addition, the phylogenetic tree (Fig. 3B) shows that *T. thermophilus* homoaconitase and its homolog of *P. horikoshii* share an ancestor with eukaryotic homoaconitases but that they possibly separated from the eukaryotic enzymes in the early stages of evolution. All the observations suggest that the homoaconitase gene was not transferred horizontally from prokaryotes to fungi.

The phylogenetic tree in Figure 3B contains aconitase in addition to homoaconitase and 3-isopropylmalate dehydratase. Aconitase (aconitate hydratase) reversibly catalyzes the conversion of *cis*-aconitic acid to either citric acid or isocitric acid in the citric acid cycle and therefore shares structural characteristics with homoaconitase and 3-isopropylmalate dehydratase. In this respect, recently Irvin and Bhattacharjee (1998) also reported that fungal homoaconitase shared a common ancestor with aconitase and 3-isopropylmalate dehydratase.

T. thermophilus RimK-Like Protein

T. thermophilus contains a *rimK*-like gene within the gene cluster for lysine biosynthesis. The RimK protein of *Escherichia coli* was isolated and characterized as a protein that adds glutamic acid residues to the carboxyl terminus of ribosomal protein S6 of the organism (Kang et al. 1989). The protein family including *E. coli* RimK is defined as a number of proteins with ATP-dependent carboxylate-amine ligase activity, where acylphosphate intermediates are involved in the catalytic mechanism (Galperin and Koonin 1997). The phylogenetic tree (Fig. 4) shows that there are two major groups: biotin carboxylase, and RimK. The *T. thermophilus* RimK homolog is again most closely related to *P. horikoshii* PH1721. Other archaea, such as *A. fulgidus*, *M. thermoautotrophicum*, and *M. jannaschii*, have at least two proteins, each of which is classified to the biotin carboxylase or RimK groups. Interestingly, *M. jannaschii* has two homologs in the RimK subgroup: One is grouped with those of *T. thermophilus* and *P. horikoshii*, and the other is grouped with those including *E. coli* RimK. Considering that RimK-like protein of *T. thermophilus* is involved in lysine biosynthesis, the RimK homolog that is grouped with those of *T. thermophilus* and *P. horikoshii* might also have a similar catalytic activity required for lysine biosynthesis.

Figure 4 Phylogenetic relationships among the amino acid sequence encoded in the *rimK* gene of *T. thermophilus* and the related sequences. A total of 164 amino acid sites were considered without gap regions in alignment. The bar indicates 10% difference of amino acid sequence. The DAD accession nos. used in this phylogenetic analysis are L14612-2-1, AE000404-10-1, AF007100-1-1, AF068249-1-1, L38260-1-1, L14862-1-1, U59234-2-1, AE000942-12-1, U67563-10-1, AE001090-1-1, AE001286-7-1, AP000007-36-1, U32828-12-1, X15859-2-1, U39679-12-1, AE000016-2-1, U67510-6-1, U67542-8-1, and AE000094-2-1.

T. thermophilus ArgC and ArgB

ArgC catalyzes the conversion of *N*-acetylglutamate 5-phosphate to *N*-acetylglutamate 5-semialdehyde. This is the third reaction in the arginine biosynthetic pathway. By homology search using PSI-BLAST (Altschul et al. 1997) for proteins homologous to that encoded by the *argC* gene in the lysine biosynthetic gene cluster of *T. thermophilus*, we found another gene having the name *argC* in the *T. thermophilus* genome (Baetens et al. 1998). This shows that *T. thermophilus* has at least two *argC* homologous genes: One is used in arginine biosynthesis, and the other is used in lysine biosynthesis. Hereafter, we call the protein working in arginine biosynthesis as true-ArgC, and tentatively call the protein involved in lysine biosynthesis as pseudo-ArgC.

In the phylogenetic tree (Fig. 5A), *T. thermophilus* true-ArgC is grouped with an ArgC homolog of *A. aeolicus* and those of archaea, while pseudo-ArgC is most closely related to an ArgC homolog, PH1720, of *P. horikoshii*. *P. horikoshii* has no other *argC* homolog in its complete genome sequence database. This suggests that ArgC of *P. horikoshii* is involved in both arginine and lysine biosyntheses. In any case, it is evident that *T. thermophilus* (and probably *P. horikoshii*) lysine biosynthesis is related to the present arginine biosynthetic pathway.

The phylogenetic tree for ArgB of *T. thermophilus* is

shown in Figure 5B. The tree shows that ArgB of *T. thermophilus* is also closely related to its homolog PH1718 in *P. horikoshii*. The *T. thermophilus* and *P. horikoshii* lineage is far from both prokaryotic and eukaryotic ArgBs which catalyze conversion of *N*-acetylglutamate to *N*-acetylglutamate 5-phosphate in the arginine biosynthetic pathway. *P. horikoshii* has a single ArgB homolog as is the case with ArgC.

In the arginine biosynthetic pathway, *N*-acetylglutamate is converted to *N*-acetylglutamate 5-semialdehyde via *N*-acetylglutamate 5-phosphate, which is a series of the reactions catalyzed by ArgB and ArgC. On the other hand, in the lysine biosynthesis pathway of eukaryotes, 2-amino adipic acid is converted to 2-amino adipic acid 6-semialdehyde via an adenosylated derivative (Fig. 1) by a multifunctional enzyme, amino adipate re-

Figure 5 Comparison of arginine biosynthesis and *Thermus* lysine biosynthesis related proteins. (A) Phylogenetic relationships among the amino acid sequence encoded in the *argC* gene of *T. thermophilus* and the related sequences. A total of 267 amino acid sites were considered without gap regions in alignment. The bar indicates 10% difference of amino acid sequence. The DAD accession nos. used in this phylogenetic analysis are D64004-19-1, U48355-1-1, X52834-1-1, X99978-2-1, AL031541-2-1, M83659-1-1, Z85982-15-1, AF049897-1-1, M21446-1-1, AE000861-7-1, U67552-4-1, AE000961-28-1, AE000761-6-1, Y10525-2-1, AP000007-35-1, Z69727-9-1, X98880-1-1, U18813-19-1, and L27746-1-1. (B) Phylogenetic relationships among the amino acid sequence encoded in the *argB* gene of *T. thermophilus* and the related sequences. A total of 157 amino acid sites were considered without gap regions in alignment. The DAD accession nos. used in this phylogenetic analysis are AE001016-12-1, AE000806-5-1, U67464-11-1, AE000771-3-1, M94625-2-1, D90905-74-1, X86157-3-1, L06036-3-1, Z26919-3-1, X98880-1-1, U18813-19-1, L27746-1-1, Z69727-9-1, and AP000007-33-1.

ductase (Morris and Jinks-Robertson 1991; Miller and Bhattacharjee 1996). No significant similarity in amino acid sequence is observed between aminoadipate reductase and ArgB/C. Because the lysine biosynthetic pathway in *T. thermophilus* is suggested to proceed in a mechanism similar to that in arginine biosynthesis, we assume that a 2-amino adipic acid derivative possibly modified by a RimK homolog is converted to a 6-semialdehyde derivative by pseudo-ArgC and ArgB in the lysine biosynthetic gene cluster. Thus, the lysine biosynthetic pathway in *T. thermophilus* may not be completely the same as that in the eukaryotes.

Horizontal Gene Transfer Between *Thermus* and *Pyrococcus*?

As described above, all the enzymes encoded in the lysine biosynthetic gene cluster in *T. thermophilus* are closely related to the proteins in *P. horikoshii*. Surprisingly, the corresponding genes in *P. horikoshii* are grouped in a manner similar to *T. thermophilus* (Fig. 6). Here, we mention the possibility of horizontal gene-cluster transfer between *T. thermophilus* and *P. horikoshii*, each of which live in a hyper-thermogenic environment, because only *T. thermophilus* and *P. horikoshii* have similar lysine biosynthetic gene clusters among the 19 organisms examined. All the phylogenetic trees show that the gene cluster for lysine biosynthesis in *T. thermophilus* and *P. horikoshii* are closely related to each other.

However, the trees also indicate that the branching point is deep, probably close to the early divergence of bacteria and archaea. Considering that the earliest organisms had lived under hot and anaerobic conditions, we assume that this gene cluster preserves one of the origins of lysine biosynthesis and that the lysine biosynthetic gene cluster has been independently conserved in both organisms. Nelson et al. (1999) reported that 24% of the predicted coding sequences in the *Thermotoga maritima* genome were most similar to proteins in archaeal species, and almost half were similar to *P. horikoshii*. However, *T. maritima* does not have the gene cluster for AAA lysine

Figure 6 Comparison of *T. thermophilus* gene cluster for lysine biosynthesis and the corresponding region in *P. horikoshii*. The percentages show amino acid sequence identity between the homologous genes.

biosynthesis but has the *lysA* gene for DAP lysine biosynthesis.

Amino Acid Biosynthesis of *P. horikoshii*

Although *P. horikoshii* does not have a gene manipulation system, its amino acid biosynthesis can be deduced by analysis of phylogenetic trees and by use of genetic evidence from *T. thermophilus*, whose lysine gene organization is similar. Diaminopimelic acid decarboxylase (LysA) catalyzes *meso*-2,6-diaminopimelic acid to lysine. This is the last reaction in the biosynthesis from aspartic acid to lysine (Fig. 1). The homology search using PSI-BLAST reveals that *P. horikoshii* has no LysA homolog. This is consistent with the idea that *P. horikoshii* synthesizes lysine not through DAP, but via the AAA pathway like *T. thermophilus*. When the gene clusters are compared between *T. thermophilus* and *P. horikoshii*, both organisms possess a single additional gene, *orfE* and PH1722, in the cluster (Fig. 6). We do not know the function of OrfE. However, the amino acid sequence of OrfE is homologous to that of OrfF, which is required for lysine biosynthesis in *T. thermophilus*. Interestingly, both OrfE and OrfF are composed of repetitive short segments of 20–30 amino acids, all of which contained a C-P/E-x-C-G motif. We speculate that gene duplication of *orfF* may have occurred. PH1722 has high amino acid sequence similarity to 3-isopropylmalate dehydrogenase (LeuB) and isocitrate dehydrogenase (data not shown). By analogy of the reactions mediated by these enzymes, we assume that PH1722 possesses homoisocitrate dehydrogenase activity. *P. horikoshii* possesses no other homologs of PH1722, PH1724, PH1726, and PH1727 in its genome. This suggests that these four proteins might be involved in leucine biosynthesis as LeuB, LeuD, LeuC, and LeuA, respectively, as well as lysine biosynthesis, as *P. horikoshii* was reported to show only Trp auxotrophy (Gonzalez et al. 1998). It should be noted that these enzymes are structurally similar to enzymes consisting of a part of the citric acid cycle (Fig. 1). Consid-

ering that the earliest organisms are believed to have lived under hot, anaerobic conditions, it is reasonable to postulate that amino acid biosynthesis developed earlier than the citric acid cycle, which functions under aerobic conditions. It is possible that the aconitase of the tricarboxylic acid (TCA) cycle evolved from homoaconitase or 3-isopropylmalate dehydratase.

In addition to the four genes PH1722, PH1724, PH1726, and PH1727, *P. horikoshii* has single *argC* and *argB* homologs in its genome. Furthermore, the archaeon has *argD* (PH1716) and *argE* (PH1715) homologs, which may have the activities of aminotransferase and deacylase just downstream of the putative *argB* gene PH1718. By analogy to arginine biosynthesis, we assume that the semialdehyde-type AAA derivative formed by PH1720 and PH1718 could be converted to lysine by the ArgD and ArgE homologs in *P. horikoshii*. The organism contains only a single copy of *argC*, *argB*, *argD*, and *argE* homologs in its genome. We therefore speculate that the products of these genes may possess the activity required for arginine biosynthesis.

Although *P. horikoshii* and *T. thermophilus* have very similar clusters of lysine biosynthetic genes, it is unlikely that these organisms have identical synthetic pathways, because *T. thermophilus* has at least two *argC* homologs while *P. horikoshii* has a single *argC* homolog. This suggests that lysine and arginine biosyntheses are separated in *T. thermophilus*. Consistent with this, the *argB* disruptant of *T. thermophilus* requires only lysine for its growth, but not arginine (Table 1). In this study, it is suggested that *P. horikoshii* might have developed unique amino acid biosynthetic systems in which several amino acids are synthesized by a limited number of enzymes with broad substrate specificity. The presence of lysine biosynthesis through AAA is suggested for an extremely thermophilic anaerobic archaeon, *Thermoproteus neutrophilus*, based on acetate assimilation patterns (Schäfer et al. 1989). It is therefore interesting to elucidate the lysine biosynthetic pathway of *T. neutrophilus*.

Evolution of Amino Acid Biosynthesis

P. horikoshii grows optimally under anaerobic conditions at a temperature of nearly 100°C. Considering that the earliest organisms had lived under hot and anaerobic conditions, *Pyrococcus* may preserve prototypic proteins. The enzyme recruitment model for the evolution of metabolic pathways suggests the enzymes which are somewhat loose in their substrate specificity could initially function in multiple pathways and evolve late into two or more specific enzymes by gene duplication (Jensen 1976). In fact, several enzymes from *Pyrococcus* are reported to have broad substrate specificity (Kengen et al. 1993; Bauer et al. 1996; Fischer et al. 1996; Mai and Adams 1996; Durbecq et al. 1997; Glasemacher et al. 1997). In this study we propose that *P. horikoshii*, like *T. thermophilus*, synthesizes lysine through the AAA pathway. We further suggest that unlike *T. thermophilus*, at least four of these lysine genes are also involved in leucine synthesis and four other genes in that cluster are involved in arginine synthesis. This system suggested for *P. horikoshii* could be one of the earliest amino acid biosynthetic systems, in which enzymes recognized at least two different substrates.

Interestingly, the other archaea, *A. fulgidus*, *M. thermoautotrophicum*, and *M. jannaschii*, each have two or three homologs of *lys20*, *hacA*, and *hacB*. The phylogenetic trees (Fig. 2A,B) show that each gene was duplicated or triplicated at an early stage of evolution, probably before the species diverged. At present, we do not know the homologs' physiological substrates. However, it may be possible to speculate that some of them are still in a stage of evolution seeking for enhancing substrate specificity. *Pyrococcus* has lived on the earth for a long time under hot and anaerobic conditions, which may allow the organism to have some prototypic enzymes. In this study we propose a hypothesis that the present biosynthetic pathways for lysine, leucine, and arginine could have developed by ancestor enzymes with broad substrate specificity. We believe that further studies on the amino acid biosynthetic systems in *P. horikoshii* will help elucidate the evolution of amino acid biosynthesis.

METHODS

T. thermophilus was transformed as described (Koyama et al. 1986). Disruption of each gene was performed according to the method of Kosuge and Hoshino (1997, 1998). Each disruption was confirmed by Southern hybridization. The detailed data are available on request.

We performed a homology search using PSI-BLAST (Altschul et al. 1997) with the given parameter values on the DNA data bank of Japan (DDBJ). This program compares a given query amino acid sequence against all other proteins in the databases to identify related sequences. Here the query amino acid sequences were AAA lysine biosynthesis-related proteins from *T. thermophilus*. Each of the multiple align-

ments was created using the CLUSTAL W program (Thompson et al. 1994) on DDBJ among a query sequence and high-scoring sequences of the PSI-BLAST result. The phylogenetic tree from p-distance estimation by the neighbor-joining method (Saitou and Nei 1987) with 1000 bootstrap analyses (Felsenstein 1985) was constructed based on the multiple alignment using MEGA version 1.01 (Kumar et al. 1993).

ACKNOWLEDGMENTS

We thank Dr. Tetsuo Hamamoto for providing helpful comments for completing the draft.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

NOTE

We renamed *lys20*, *hacA* (*lys4A*), *hacB* (*lys4B*), *orfE*, *orfF*, *rimK*, *argC*, and *argB* in the cluster of *T. thermophilus*, as *lysS*, *lysT*, *lysU*, *lysV*, *lysW*, *lysX*, *lysY*, and *lysZ*, respectively, because the cluster is actually involved in lysine biosynthesis.

REFERENCES

- Altschul, S.F., T.L. Madden, A.A. Schäffer, J. Zhang, Z. Zhang, W. Miller, and D.J. Lipman. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acid Res.* **25**: 3389–3402.
- Baetens, M., C. Legrain, A. Boyen, and M. Glansdorff. 1998. Genes and enzymes of the acetyl cycle of arginine biosynthesis in the extreme thermophilic bacterium *Thermus thermophilus* HB27. *Microbiology* **144**: 479–492.
- Bauer, M.W., E.J. Bylina, R.V. Swanson, and R.M. Kelly. 1996. Comparison of β -glucosidase and β -mannosidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. Purification, characterization, gene cloning, and sequence analysis. *J. Biol. Chem.* **271**: 23749–23755.
- Broquist, H.P. 1971. Lysine biosynthesis (yeast). *Methods Enzymol.* **17**: 112–129.
- Bult, C.J., O. White, G.J. Olsen, L. Zhou, R.D. Fleischmann, G.G. Sutton, J.A. Blake, L.M. FitzGerald, R.A. Clayton, J.D. Gocayne et al. 1996. Complete genome sequence of the methanogenic archaeon, *Methanococcus jannaschii*. *Science* **273**: 1058–1073.
- Deckert, G., P.V. Warren, T. Gaasterland, W.G. Young, A.L. Lenox, D.E. Graham, R. Overbeek, M.A. Snead, M. Keller, M. Aujay et al. 1998. The complete genome of the hyperthermophilic bacterium *Aquifex aeolicus*. *Nature* **392**: 353–358.
- Durbecq, V., C. Legrain, M. Roovers, A. Piérard, and N. Glansdorff. 1997. The carbamate kinase-like carbamoyl phosphate synthetase of the hyperthermophilic archaeon *Pyrococcus furiosus*: A missing link in the evolution of carbamoyl phosphate biosynthesis. *Proc. Natl. Acad. Sci.* **94**: 12803–12808.
- Felsenstein, J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* **39**: 783–791.
- Fischer, L., R. Bromann, S.W. Kengen, W.M. de Vos, and F. Wagner. 1996. Catalytic potency of β -glucosidase from the extremophile *Pyrococcus furiosus* in glucoconjugate synthesis. *Biotechnology* **14**: 88–91.
- Galperin, M.Y. and E.V.A. Koonin. 1997. Diverse superfamily of enzymes with ATP-dependent carboxylate-amine/thiol ligase activity. *Protein Sci.* **12**: 2639–2643.
- Glasemacher, J., A.K. Bock, R. Schmid, and P. Schönheit. 1997. Purification and properties of acetyl-CoA synthase (ADP-forming), an archaeal enzyme of acetate formation and ATP synthesis, from the hyperthermophile *Pyrococcus furiosus*. *Eur. J. Biochem.* **244**: 561–567.
- Gonzalez, J.M., Y. Masuchi, F.T. Robb, J.W. Ammerman, D.L. Maeder, M. Yanagibayashi, J. Tamaoka, and C. Kato. 1998. *Pyrococcus horikoshii* sp. nov., a hyperthermophilic archaeon

- isolated from a hydrothermal vent at the Okinawa Trough. *Extremophiles* **2**: 123–130.
- Irvin, S.D. and J.K. Bhattacharjee. 1998. A unique fungal lysine biosynthesis enzyme shares a common ancestor with tricarboxylic acid cycle and leucine biosynthetic enzymes found in diverse organisms. *J. Mol. Evol.* **46**: 401–408.
- Jensen, R.A. 1976. Enzyme recruitment in evolution of new function. *Annu. Rev. Microbiol.* **30**: 409–425.
- Kang, W.K., T. Icho, S. Isono, M. Kitakawa, and K. Isono. 1989. Characterization of the gene *rimK* responsible for the addition of glutamic acid residues to the C-terminus of ribosomal protein S6 in *Escherichia coli* K12. *Mol. & Gen. Genet.* **217**: 281–288.
- Kawarabayashi, Y., M. Sawada, H. Horikawa, Y. Haikawa, Y. Hino, S. Yamamoto, M. Sekine, S. Baba, H. Kosugi, A. Hosoyama et al. 1998. Complete sequence and gene organization of the genome of a hyper-thermophilic archaeobacterium, *Pyrococcus horikoshii* OT3. *DNA Res.* **5**: 55–76.
- Kengen, S.W., E.J. Luesink, A.J. Stams, and A.J. Zehnder. 1993. Purification and characterization of extremely thermostable β -glucosidase from the hyperthermophilic archaeon *Pyrococcus furiosus*. *Eur. J. Biochem.* **213**: 305–312.
- Klenk, H.-P., R.A. Clayton, J.-F. Tomb, O. White, K.E. Nelson, K.A. Ketchum, R.J. Dodson, M. Gwinn, E.K. Hickey, J.D. Peterson et al. 1997. The complete genome sequence of the hyperthermophilic, sulphate-reducing archaeon *Archaeoglobus fulgidus*. *Nature* **390**: 364–370.
- Kobashi, N., M. Nishiyama, and M. Tanokura. 1999. Aspartate kinase-independent lysine synthesis in an extremely thermophilic bacterium, *Thermus thermophilus*: Lysine is synthesized via α -aminoadipic acid not via diaminopimelic acid. *J. Bacteriol.* **181**: 1713–1718.
- Kosuge, T. and T. Hoshino. 1997. Molecular cloning and sequence analysis of the *lysR* gene from the extremely thermophilic eubacterium, *Thermus thermophilus* HB27. *FEMS Microbiol. Lett.* **157**: 73–79.
- . 1998. Lysine is synthesized through the α -aminoadipate pathway in *Thermus thermophilus*. *FEMS Microbiol. Lett.* **169**: 361–367.
- Koyama, Y., T. Hoshino, N. Tomizuka, and K. Furukawa. 1986. Genetic transformation of the extreme thermophile *Thermus thermophilus* and of other *Thermus* spp. *J. Bacteriol.* **166**: 338–340.
- Kumar, S., K. Tamura, and M. Nei. 1993. MEGA: Molecular Evolutionary Genetics Analysis, version 1.01. Pennsylvania State University, University Park, PA.
- Mai, X. and M.W. Adams. 1996. Purification and characterization of two reversible and ATP-dependent acetyl coenzyme A synthases from the hyperthermophilic archaeon *Pyrococcus furiosus*. *J. Bacteriol.* **178**: 5897–5903.
- Miller, K.G. and J.K. Bhattacharjee. 1996. The *LYS5* gene of *Saccharomyces cerevisiae*. *Gene* **172**: 167–168.
- Morris, M.E. and S. Jinks-Robertson. 1991. Nucleotide sequence of the *LYS2* gene of *Saccharomyces cerevisiae*: Homology to *Bacillus brevis* tyrocidine synthetase 1. *Gene* **98**: 141–145.
- Nelson, K.E., R.A. Clayton, S.R. Gill, M.L. Gwinn, R.J. Dodson, D.H. Haft, E.K. Hickey, J.D. Peterson, W.C. Nelson, K.A. Ketchum et al. 1999. Evidence for lateral gene transfer between Archaea and Bacteria from genome sequence of *Thermotoga maritima*. *Nature* **399**: 323–329.
- Saitou, N. and M. Nei. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406–425.
- Schäfer, S., T. Paalme, R. Vilu, and G. Fuchs. 1989. ^{13}C -NMR study of acetate assimilation in *Thermoproteus neutrophilus*. *Eur. J. Biochem.* **186**: 695–700.
- Smith, D.R., L.A. Doucette-Stamm, C. Deloughery, H. Lee, J. Dubois, T. Aldredge, R. Bashirzadeh, D. Blakely, R. Cook, K. Gilbert et al. 1997. Complete genome sequence of *Methanobacterium thermoautotrophicum* ΔH : Functional analysis and comparative genomics. *J. Bacteriol.* **179**: 7135–7155.
- Thompson, J.D., D.G. Higgins, and T.J. Gilson. 1994. CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acid Res.* **22**: 4553–4559.
- Vogel, H.J. 1964. Distribution of lysine pathways among fungi: Evolutionary implications. *Am. Nat.* **98**: 446–455.
- . 1965. Lysine biosynthesis and evolution. In *Evolving genes and proteins* (ed. V. Bryson and H.J. Vogel), pp. 25–40. Academic Press, New York, NY.

Received July 13, 1999; accepted in revised form October 14, 1999.

A Prokaryotic Gene Cluster Involved in Synthesis of Lysine through the Amino Adipate Pathway: A Key to the Evolution of Amino Acid Biosynthesis

Hiromi Nishida, Makoto Nishiyama, Nobuyuki Kobashi, et al.

Genome Res. 1999 9: 1175-1183

Access the most recent version at doi:[10.1101/gr.9.12.1175](https://doi.org/10.1101/gr.9.12.1175)

References

This article cites 31 articles, 7 of which can be accessed free at:
<http://genome.cshlp.org/content/9/12/1175.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Affordable, Accurate
Sequencing.

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>