Methods

High-Throughput Isolation of *Caenorhabditis elegans* Deletion Mutants

Leo X. Liu,^{1,4} Jill M. Spoerke, Evan L. Mulligan, Jing Chen,² Brian Reardon, Bethany Westlund, Lin Sun,³ Ken Abel, Barbara Armstrong, Gary Hardiman, Judith King, Lisa McCague, Michael Basson, Ralph Clover, and Carl D. Johnson

Axys Pharmaceuticals, NemaPharm Group, South San Francisco, California 94080 and La Jolla, California USA

The nematode *Caenorhabditis elegans* is the first animal whose genome is completely sequenced, providing a rich source of gene information relevant to metazoan biology and human disease. This abundant sequence information permits a broad-based gene inactivation approach in *C. elegans*, in which chemically mutagenized nematode populations are screened by PCR for deletion mutations in a specific targeted gene. By handling mutagenized worm growth, genomic DNA templates, PCR screens, and mutant recovery all in 96-well microtiter plates, we have scaled up this approach to isolate deletion mutations in >100 genes to date. Four chemical mutagens, including ethyl methane sulfonate, ethylnitrosourea, diepoxystyrene, and ultraviolet-activated trimethylpsoralen, induced detectable deletions at comparable frequencies. The deletions averaged ~1400 bp in size when using a ~3 kb screening window. The vast majority of detected deletions removed portions of one or more exons, likely resulting in loss of gene function. This approach requires only the knowledge of a target gene sequence and a suitable mutagen, and thus provides a scalable systematic approach to gene inactivation for any organism that can be handled in high density arrays.

The rapid progress in genome sequencing projects has propelled a shift in genome analysis from structural genomics to functional genomics, that is, the genome-driven systematic study of gene function (Hieter and Boguski 1997). In this postgenomic era, model organisms in which large-scale functional analyses and rapid genetic experiments are possible—chiefly the yeast *Saccharomyces cerevisiae*, the fruit fly *Drosophila melanogaster*, and the nematode *Caenorhabditis elegans*—are increasingly useful in understanding human disease pathways (Miklos and Rubin 1996; Oliver 1996; Ahringer 1997). With the recent completion of the *C. elegans* genomic sequence (The *C. elegans* Sequencing Consortium 1998), the first complete animal genome is now available. Computational analysis of partial *C. elegans* genome data has revealed that many positionally cloned human disease genes have *C. elegans* orthologs (genes encoding proteins with similar multidomain architecture and predicted function) (Mushegian et al. 1997, 1998). The *C. elegans* genome also contains a significant proportion of apparently nematode-specific protein families that may be relevant for nematode biology and parasitism (Sonnhammer and Durbin 1997; Blaxter 1998).

A rapid method to ascertain gene function by targeted gene inactivation in *C. elegans* would be highly desirable, but homologous recombination as in the mouse has not yet proven feasible (Plasterk 1995). Microinjection of target-specific RNA, for reasons not completely understood, is a remarkably effective means of transcriptional interruption in this small organism (Fire et al. 1998). However, the extent of RNA-mediated inactivation is difficult to assess in many cases, and such inactivation obviously does not produce a germ-line lesion necessary for genetic crosses, suppressor screens, and other longer-term genetic manipulations. One large-scale approach to germ-line inactivation is to induce random mutations in the animal population, followed by screening for mutations in a target gene of known sequence. The most well-developed method of this so-called target-selected gene inactivation in *C. elegans* has used random transposon Tc1 insertions to generate a collection of mutants, followed by PCR screening for the presence of Tc1 in a gene of choice (Zwaal et al. 1993). However, because Tc1 insertion alone does not usually result in gene inactivation, it is necessary to subsequently screen individual Tc1 alleles for animals in which the transposon and flanking DNA have been deleted through transposon excision, a relatively infrequent event (Plasterk 1995). An alternative approach is to use chemical mutagens to directly induce deletions in a population, and then to screen by PCR for deleted segments within a selected target region (Yandell et al. 1994). Jansen et al.

Present addresses: ¹Cambria Biosciences, Waltham, Massachusetts 02454 USA; ²Therion Biologics, Cambridge, Massachusetts 02138 USA; ³Phylos, Lexington, Massachusetts 02421 USA.

⁴Corresponding author.

E-MAIL lliu@cambriabio.com; FAX (781) 736-3107.

(1997, 1999) have established the broader feasibility of this approach by isolating mutants of the *C. elegans* heterotrimeric G protein gene family.

In this study we describe our results using random chemical mutagenesis and PCR screening to rapidly isolate deletion mutations in a large number of genes encoding proteins with a broad range of functions. *C. elegans* is unique among model animal species in that it can be grown in liquid cultures and also can be stored frozen but viable at -80°C . We have taken advantage of these properties to devise a rapid and scalable procedure for gene disruption almost entirely on the basis of microtiter plate arrays of whole animals and genomic DNA. We used four different chemical agents to create mutagenized libraries, and found that all four mutagens induce detectable deletions. Almost all of the deletions were significant enough to result in loss of exons, frame shifts, and other molecular lesions likely to cause loss of gene function. We discuss the sensitivity, specificity, limitations, and broader utility of this approach to systematic gene inactivation in *C. elegans*.

RESULTS

Library Construction and Screening

To enhance the scalability and speed of mutagenized library construction and screening, we relied on microtiter plate-based culture and analysis wherever possible (Fig. 1). F_1 larvae of mutagenized P_0 animals were distributed directly into microtiter plate wells (~ 20 larvae representing 40 mutagenized genomes) and grown in liquid cultures to yield F_2 larvae. Approximately one-half of the worms were processed into a corresponding genomic DNA microtiter plate array, with the remainder frozen at -80°C as viable animals. The initial round of nested PCR screening for deletion mutations in a targeted gene was performed on microtiter plate pools each containing ~ 4000 mutagenized genomes. PCR primers were designed to flank ~ 3 kb of exon-rich regions of the selected ORF, in many cases specifically targeting putative functional domains. Amplicons smaller than the wild-type amplicon were taken to represent candidate deletions. Such amplicons were expected to be preferentially amplified, compen-

Figure 1 Schematic of mutant library construction and screening. F_1 larvae of chemically mutagenized P_0 animals are cultured in microtiter arrays to produce F_2 larvae, constituting the worm library. An aliquot of worms from each well is removed and processed into a corresponding genomic DNA array; simultaneously, the 96 samples from each plate are pooled and processed to produce genomic DNA plate pools. The DNA plate pools are screened by nested PCR in 96-well plates for deletion amplicons, which identify a positive worm library plate Y. After confirmation by quadruplicate PCR (not illustrated), the genomic DNA array for that specific library plate is similarly screened to identify the specific well. The corresponding well is then retrieved from the frozen mutant worm library and thawed. Individual live worms are allowed to have self-progeny in microtiter wells, and aliquots of each well are tested by PCR to determine which of the thawed worms contain the deletion (see Methods for details).

sating for the much lower representation of the mutant target template within the genomic DNA pool. Candidate-positive plate pools were resampled to eliminate false positives, which constituted about three-fourths of the first-round candidate deletions. After confirmation that a plate pool contained a deletion, the corresponding DNA array was then screened to identify a specific microtiter well. The corresponding well of the frozen worm library was thawed and individual animals were grown in microtiter wells; portions of these cultures were screened by PCR to identify heterozygous and homozygous mutant strains. A small number of thawed library wells failed to yield the expected deletion mutant, and the failure rate exceeded 50% when <100 viable animals were recovered from a well (data not shown).

To evaluate various chemicals that have been used to induce mutations in *C. elegans*, we created mutant libraries using different mutagens, including ethylmethane sulfonate (EMS), ethylnitrosourea (ENU), diepoxyoctane (DEO), and ultraviolet-activated trimethylpsoralen (UV-TMP). Each library consisted of forty-eight 96-well plate arrays and ~200,000 mutagenized genomes. For any given target, usually 4–8 such libraries (comprising 0.8–1.6 million mutagenized genomes) were screened simultaneously for potential deletions. At this scale, deletion mutants were isolated from almost all loci that were targeted, and in many instances more than one deletion was identified; in the latter cases usually only the largest deletion was thawed from the worm library. We were unable to detect deletions in a small number (<5) of targeted genes, and these failed loci were too few to determine whether they shared any structural characteristics. The average number of animals recovered from frozen wells differed for each mutagen as follows: DEO 191 ± 159 , EMS 359 ± 330 , ENU 343 ± 249 , and UV-TMP 199 ± 166 .

Deletion Mutations

To date we have isolated deletion mutations in >100 independent gene targets using this method, and a representative listing is shown in Table 1. Among these are genes encoding a broad range of protein functional groups, including kinases, transcription factors, membrane receptors, metabolic enzymes, proteases, hormones, and nematode homologs of oncogenes and tumor suppressor genes. However, the largest target group comprises genes encoding proteins of unknown function, reflecting our strategic choice of using the *C. elegans* model to elucidate the function of novel predicted proteins.

The average size of chemically induced deletions was ~1400 bp (range 700–2900 bp), as detected by a PCR-screening window of ~3 kb (Fig. 2). In several instances, smaller deletions (on the order of 400–500 bp)

were fortuitously detected on screening of a specific library plate. The average size of deletions isolated from different mutagenized libraries ranged from ~1320 bp for EMS up to 1580 bp for DEO, and a broad size range was observed for each agent (Fig. 2). DNA sequencing revealed that the majority were simple deletions, but about one-fifth of the sequenced deletions contained small (3–27 bp) insertions at the breakpoints. All of the latter deletions derived from libraries mutagenized by DEO, EMS, or ENU. Only one of our isolated deletions (*xfl-1*, *nr2025*), isolated from a UV-TMP library, contained a large inverted insertion, which is in contrast with the examples reported previously for *gpa* deletion mutants obtained by UV-TMP, in which two of three deletions contained large insertions (Jansen et al. 1997). In addition, in our series, both the proximal and distal deletion breakpoints in most cases fell within a region of two or more consecutive identical nucleotides, rendering the exact breakpoint ambiguous.

C. elegans genes are compact, with most introns ~50 bp and exons 80–250 bp in length (Blumenthal and Steward 1997). Thus, deletions of >500 bp within a targeted *C. elegans* ORF are likely to significantly compromise gene function. Within an ORF, the position of such disabling deletions can be as follows: (1) the deletion begins or ends in an exon, removing portions of one or more exons, and usually leads to a premature stop codon (Fig. 3a); (2) the deletion begins and ends in introns, but removes one or more exons encoding domains essential for protein function (Fig. 3b); or (3) the deletion removes the entire ORF (Fig. 3c). Less than 2% of the deletion mutations that we isolated removed only noncoding DNA (data not shown), a figure that was optimized by selecting target regions that excluded long stretches of noncoding DNA. Given the characteristics of these molecular lesions, it is likely that most of these intragenic deletions represent severe or null mutations. The resulting phenotypes range from apparently wild-type to dramatic and lethal defects. Of note, less than one-half of the isolated deletion mutants (Table 1) displayed an obvious phenotype (such as lethal, uncoordinated, or sterile), but further analysis is necessary and ongoing in many cases.

Although generally 4–8 libraries were screened for each target, the total number of mutant libraries generated during the period of this study was greater. Additional libraries were produced as template DNA became depleted in the course of multiple target screens, and all of these libraries were included for analysis. In addition, the cumulative number of libraries prepared with each chemical mutagen was different for each agent. To estimate the likelihood of isolating a deletion with different mutagens, we compared the number of deletions relative to the total number of primary library screens for each mutagen (Table 2). By this analy-

Table 1. Selected Deletion Mutations

Cosmid name	GenBank acc. no.	Gene or protein	Allele name	Mutagen	Deletion size (bp)	Deletion effect on ORF	Preliminary phenotype	Reference ^a
B0478.1	U57054	Jun kinase homolog	nr2035	DEO	1825	promoter-intron 7	no obvious phenotype	S. Kostas and A. Fire D. Fambrough
C02B8.4	U40421	Twist transcription factor homolog	nr2061	EMS	1267	Intron 1-exon 3	no obvious phenotype	
C03B1.12	U40952	Lysosomal-associated membrane protein	nr2045	ENU	1772	promoter-intron 3	no obvious phenotype	
C04G6.3	U55854	phospholipase D homolog	nr2015	EMS	545	exon 7-exon 8	no obvious phenotype	M. Koelle Westlund (1998)
C05B5.7	Z32679	RGS GAP protein homolog	nr2017	EMS	637	exon 1-intron 2	mild egg-laying defective	
C18E3.8	AF000265	<i>hop-1</i> presenilin	nr2003	UV-TMP	1196	promoter-intron 3	decreased brood size at 15° C	
C44H4.6	Z79598	GSK-3 β kinase homolog	nr2056	EMS	1411	intron 2-intron 6	no obvious phenotype	S. Pierce; Gregoire et al. (1998)
C47E8.5	Z75530	<i>daf-21</i> HSP90 homolog	nr2081	ENU	863	exon 1-exon 4	larval arrest	
F02E8.6	U53340	Niemann-Pick type C protein homolog	nr2022	EMS	1120	exon 3-exon 6	no obvious phenotype	
F10B5.4	Z48334	Tubby homolog	nr2044	DEO	1816	intron 1-beyond 3' terminus	no obvious phenotype	P. Sengupta A. Golden R. Ranganathan
F13B12.a	Z70683	insulin homolog	nr2091	EMS	1653	deletes entire ORF	no obvious phenotype	
F28H6.1	AL031621	<i>akt-2</i> kinase	nr2021	DEO	2208	promoter-exon 4	no obvious phenotype	
F35H12.3	U41540	<i>sel-12</i> presenilin	nr2011	ENU	1426	intron 2-intron 6	egg-laying defect	Westlund (1998)
F59G1.5	U53332	Frataxin homolog	nr2040	DEO	505	promoter-intron 2	no obvious phenotype	
K06A1.4	U23449	nuclear hormone receptor homolog	nr2034	ENU	1189	exon 3-intron 5	no obvious phenotype	
K06A5.7	AF039038	<i>cdc-25</i> homolog	nr2036	UV-TMP	2695	deletes entire ORF	sterile with reduced germline	Hobert et al. (1999)
K06C4.6	U64843	<i>mod-1</i> serotonin-gated ion channel	nr2043	EMS	1209	intron 2-exon 5	resistant to serotonin-stimulated locomotion	
K06E6.1	U55375	<i>lin-6</i> homeodomain protein	nr2073	ENU	1703	exon 4-beyond 3' terminus	expulsion defective, nose-up foraging defective, synDaf-c, axonal defects	
K10C3.6	Z83235	HNF4 transcription factor homolog	nr2041	EMS	893	intron 2-exon 4	no obvious phenotype	Sluder et al. (1999)
R11A5.4	Z83122	phosphoenolpyruvate kinase homolog	nr2043	EMS	1902	intron 2-exon 5	no obvious phenotype	
T01H8.1	Z80219	p90RSK kinase homolog	nr2031	EMS	758	intron 2-exon 2	no obvious phenotype	
T07A9.6	AF036706	<i>daf-18</i> PTEN homolog	nr2037	UV-TMP	990	intron 2-exon 3	dauer defective	Gil et al. (1999) D. Zarkower
T14B1.1	Z46829	<i>tip-1</i> novel protein	nr2084	ENU	1206	exon 1-beyond 3' terminus	no obvious phenotype	
T20H4.1	U00037	<i>osm-10</i> novel protein	nr2076	EMS	1122	intron 2-exon 5	osmolarity response defective	
Y18D10	AL008871	GSK-3 β kinase homolog	nr2047	ENU	1526	intron 2-beyond 3' terminus	maternal effect early embryonic lethal	D. Hoepfner; Shaham (1998)
Y47D3	AL031635	p70S6 kinase homolog	nr2074	ENU	850	intron 2-exon 3	no obvious phenotype	
Y48E1B.13	Z93393	<i>csp-1</i> caspase homolog	nr2018	UV-TMP	1220	exon 4-intron 5	no obvious phenotype	
ZK154.7	U70844	TNF-converting enzyme homolog	nr2030	EMS	787	exon 9-intron 14	no obvious phenotype	E. Kipreos P. Kuwabara
ZK520.4	Z50796	<i>cul-2</i> cell cycle cullin	nr2020	EMS	1050	exon 5-exon 5	embryonic arrest	
ZK632/6	Z22181	calnexin homolog	nr2010	UV-TMP	774	exon 1-exon 3	no obvious phenotype	
ZK675.1	Z46812	patched homolog	nr2029	EMS	960	exon 6-exon 6	sterile	

^aPersonal communications, unless otherwise noted.

Figure 2 Deletion size by mutagen, depicting those deletions selected with a PCR screening window of 2800–3400 bp. Not shown are a small number of larger (>3000 bp) deletions identified with PCR primers placed 4–5 kb apart in the target region, as well as smaller (<500 bp) deletions fortuitously detected on screening of specific library plates. Horizontal bar, average deletion size for each mutagen.

sis, UV-TMP appeared to generate a higher frequency of deletions compared with the other three chemical mutagens.

DISCUSSION

In this paper, we describe our experience with a rapid procedure for isolating deletion mutations in *C. elegans* that uses PCR to screen for deletions in a target gene within a randomly mutagenized population of animals. The growth of mutagenized animals, preparation of genomic DNA templates, PCR screens, and mutant worm recovery are all handled in scalable microtiter liquid plate well formats that maximize speed and utility. All of the mutagens we used resulted in significant deletion mutations in target genes almost certainly sufficient to produce loss of function.

In practice, this protocol yields a deletion in a selected target for roughly every 600,000 mutagenized genomes screened. However, many factors affect the apparent sensitivity of this procedure. Small deletions of <500 bp were not isolated in screening plate pools for two likely reasons. First, such amplicons would be difficult to distinguish from an abundant wild-type band on a gel, unless special and more time-consuming electrophoresis conditions were applied. Second, mutant templates with small deletions would be similar in size to the much more abundant (4000-fold in the primary screening pools) wild-type templates and less likely to be preferentially amplified. The fact that we discovered several smaller (<500 bp) deletions only on screening of plate arrays (100-fold less

complex than the plate pools) suggests that smaller deletions may be more abundant than larger deletions. Reducing the complexity of the DNA pools may improve the sensitivity of detecting these smaller deletions.

Larger deletions of >3 kb would also be useful because of their increased likelihood of severely affecting coding regions; furthermore, in certain cases the simultaneous disruption of two adjacent genes (e.g., evolutionarily duplicated genes) by one large deletion is desirable. The upper range of detectable deletions is limited by the size of the PCR deletion screening window, which, in this series, was 2800–3400 bp, and the processivity of the thermostable polymerase and other components of the amplification reaction. Modification of PCR reagents and conditions would allow a larger screening window, and we have recently isolated larger deletions using a target window of up to 5 kb (data not shown). Increasing the sensitivity of deletion detection would also increase the likelihood of obtaining more than one allele for each target in a single screen, which would be desirable for comparative genetic and phenotypic analysis.

This PCR-based approach is highly specific because of the use of nested pairs of screening primers derived from the target gene. We sequenced a number of false-positive deletion amplicons from the primary round of plate pool screening; all were derived from the target gene rather than an extraneous locus (data not shown). Although the genesis of these truncated amplicons is incompletely understood, we observed a comparable number of false-positive amplicons using unmutagenized genomic DNA as a control template (data not shown), and suspect that they may arise from polymerase slippage across gaps formed by secondary loops in the DNA template.

A wide variety of chemicals and irradiation methods have been used to induce mutations in *C. elegans* (Anderson 1995), and we used four in making our libraries for reverse genetics: EMS, ENU, DEO, and UV-TMP. EMS is the most widely used mutagen for *C. elegans* because of its potency in generating loss-of-function or reduction-of-function alleles in classical phenotypic screens. Many EMS-induced mutations are point mutations (of which the vast majority are G/C→A/T transitions) (Anderson 1995), which, of course, would not be detected in a screen for deletions. However, ~13% of EMS-induced mutations are reported to be deletions or other rearrangements, though most are small (Anderson 1995). ENU appears to generate a rate of mutations similar to EMS; studies of ENU-induced mutations in the *unc-93* gene revealed an increased frequency of A/T→G/C transitions compared with EMS, with a frequency of intragenic deletions of 13% (De Stasio et al. 1997). DEO has not been as broadly used as a *C. elegans* mutagen, but a study of

Figure 3 Schematics of representative deletions. (a) The deletion of *daf-18* PTEN removes a portion of exon 3, which encodes the distal portion of the tensin homology domain and the catalytic phosphatase domain known to be required for lipid phosphatase activity (Gil et al. 1999). (b) The *sel-12* deletion removes exons 3–6, which encode predicted transmembrane domains necessary for proper protein topology (Westlund 1998). (c) The *cdc-25* deletion removes the entire ORF.

mutations of the *unc-54* gene revealed that DEO generated mutations at a lower rate than EMS, and 27% of DEO-induced mutations were multilocus deletions and deficiencies, as defined by genetic criteria (Anderson and Brenner 1984). Treatment with trimethylpsoralen followed by ultraviolet activation, a procedure that leads to DNA cross-linking, appears to generate a lower frequency of mutations than EMS, as measured by a generation of *unc-22* and *pal-1* mutations, but a relatively high proportion of deletions (Yandell et al. 1994).

For the purposes of a PCR-based reverse genetics screening approach, it is important to identify a mutagen that generates the highest frequency of small deletions (on the order of 100–5000 bp) rather than point mutations that cannot be detected. Furthermore, highly potent inducers of large numbers of point mu-

tations would exacerbate the confounding background of extraneous mutations. In our experience, libraries prepared with each of the four mutagens yielded deletions at similar rates (Table 2). In addition, the average size of detected deletions induced by the different mutagens was not significantly different (Fig. 2). The frequency of deletions obtained from each of the four mutagens far exceeded the spontaneous rate of deletions of this size range, which is predicted to be <1 per 100 million genomes (P. Anderson, pers. comm.). About one-fifth of our chemically induced mutations contained short (≤ 7 bp) direct sequence repeats or longer (> 2 bp) insertions at the deletion termini, in excess of what would be expected from spontaneous deletions (Pulak and Anderson 1988; data not shown). Such insertions may reflect mechanisms of break repair with nonhomologous endjoining, in response to chemically induced breaks in double-stranded DNA (Friedberg et al. 1995). To measure the efficiency of each mutagen in

creating detectable deletions, we retrospectively determined the relative yield of deletions from each type of mutagen library (Table 2). By this indirect measure, UV-activated trimethylpsoralen appears to generate the highest frequency of detectable deletions, which confirms prior observations (Yandell et al. 1994). However, fewer animals, on average, were recovered from frozen UV-TMP-treated worm populations compared with EMS or ENU, suggesting that UV-TMP treatment may cause more generally deleterious effects. In our experience, γ irradiation or formaldehyde treatment, which reportedly causes more major rearrangements (Anderson 1995), did not produce significant numbers of deletion mutations in limited experiments (data not shown). Because deletions, rather than mutations per se (the majority of which are point mutations), are the goal of this reverse genetics screening approach, the identification of a specific deletogen chemical would be very helpful.

A disadvantage of random mutagenesis procedures is that substantial numbers of extraneous mutations are generated in the background of any selected target gene, and it can be difficult to titrate the mutagen dose so as to induce large numbers of mutations in the population while also minimizing the number of mutations in any one animal. A strong mutant phenotype may result from the deletion of interest or from mutation of a non-target gene. Similarly, a pleiotropic phenotype may be due to either multiple effects of a single mutation or to mutations in more than one gene. Thus, thorough outcrossing followed by transgenic res-

Table 2. Deletion Yield by Mutagen

Mutagen	No. of libraries	No. of 1° screens	No. of deletion mutants	Ratio of deletions/screens
DEO	3	171	12	0.070
EMS	9	479	47	0.098
ENU	4	208	23	0.111
UV-TMP	4	135	29	0.215

The total number of pool screens initiated (using a target window of 2800–3400 bp) and the total number of isolated deletions were tabulated for each mutant library.

cue of the observed phenotype with a wild-type copy of the target gene (usually in the form of a cosmid or plasmid clone), is necessary to establish whether a putative loss-of-function phenotype is actually due to the deletion. In addition, reverse genetics may lead to no obvious phenotype at all. Earlier estimates suggested that up to one-half of *C. elegans* genes may cause no obvious phenotype when null (Park and Horvitz 1986). When no phenotype is seen on isolation of the mutant animal, it can be difficult to tell whether this lack of effect is because (1) the randomly generated molecular lesion does not produce loss or reduction of gene function, (2) the gene is functionally redundant, or (3) the phenotype requires specific environmental conditions for its expression or is otherwise too subtle to be distinguished by conventional methods. The complementary use of RNA-mediated interference (Fire et al. 1998) might be helpful in cases in which deletion mutants display a confusing pleiotropy or an apparent wild-type phenotype.

At the current state of development, this method requires ~2 weeks for identification of a deletion and isolation of mutant worms. The rate of deletion identification is about one target per week, per person, without automation. This rapid approach to gene inactivation allows one to simultaneously approach a group of candidate genes, whether they belong to a single gene family or may be functionally related in a putative signaling pathway. Thus, deletion mutants of different, but potentially related, genes may be studied as a group for comparative phenotype analysis, genetic interactions, etc., rather than individually. For example, genetic and phenotypic analysis have revealed overlapping functions between the *C. elegans* presenilins *sel-12* and *hop-1* (Li and Greenwald 1997; Westlund et al. 1999) and in the human Niemann–Pick type-C disease gene homologs *npc-1* and *npc-2* (M. Sym, pers. comm.).

An obvious advantage of this target-selected approach is that specific regions of functional importance can be targeted to maximize the likelihood of obtaining null or severe loss-of-function mutations. On the other hand, this method is less well suited than traditional noncomplementation genetic screens for obtaining weaker (hypomorphic) alleles, which are often useful in genetic pathway analysis.

A procedure such as this one that uses random mutagenesis followed by deletion screening is ideally suited for organisms like *C. elegans*, in which pre-existing genomic sequence information is extensive but homologous recombination or transposon insertion methods are not practical. By improving the sensitivity of detection in more complex mutagenized genome pools, it may be possible to produce a clonal library of individually arrayed mutants, thereby eliminating the need to sort through thawed worms for the

deletion. Furthermore, by increasing the density of *C. elegans* growth and DNA microarrays, and automating the PCR set up and analysis, it should be feasible to pursue the systematic disruption of all of the predicted genes in this model animal, as is currently being done, by different methods, for all of the genes in *S. cerevisiae* (Dujon 1998).

METHODS

Bioinformatics

C. elegans homologs of non-nematode genes and proteins were identified by BLAST search (Altschul et al. 1990) of finished *C. elegans* sequences available at the National Center for Biotechnology Information (www.ncbi.nlm.nih.gov) and unannotated sequences available at the Sanger Centre (www.sanger.ac.uk/Projects/C_elegans/blast_server.html) and the Washington University Genome Sequencing Center (genome.wustl.edu/gsc/blast/blast_servers.html). Putative ORFs of unannotated sequences were determined with the GeneFinder algorithm available from the Baylor College of Medicine Human Genome Center (dot.imgen.bcm.tmc.edu:9331/gene-finder/gf.html). Nested PCR primer pairs were designed to flank a target window of 2.8–3.2 kb with MacVector 6.0. Whenever possible, the target window was chosen to include the most proximal exons, except in those cases in which the first intron was >1 kb in size.

Construction of Mutagenized Nematode and Corresponding DNA Libraries

Wild-type Bristol N2 nematodes were cultured by standard methods (Sulston and Hodgkin 1988). A synchronized population of L₄ larvae was treated with one of the following four mutagens: 25 mM ethylmethane sulfonate for 4 hr, 0.4 mM ethylnitrosourea for 4 hr, 1 mM diepoxyoctane for 4 hr, or 30 µg/ml trimethylpsoralen (Yandell et al. 1994), followed by UV irradiation at 3.5 uW/cm² for 15–30 sec (Anderson 1995). The efficiency of mutation was evaluated by quantitating the frequency of F₂ resistance to ivermectin (Rand and Johnson 1995); libraries were discarded if the proportion of genomes carrying newly induced ivermectin resistance mutations was <0.5%. F₁ eggs of mutagenized worms were collected for 4 hr, then allowed to hatch for 16 hr. Hatched larvae were distributed to 96-well microtiter plates at ~20 worms per well in 50 µl of NGM culture medium containing 1% (by volume) *Escherichia coli* HB101 as food (Sulston and Hodgkin 1988). A library from each chemical mutagenesis comprised 48 microtiter plates and included the arrayed progeny of nearly 10⁵ F₁ animals representing 2 × 10⁵ mutagenized genomes. Worms were grown in wells until the food was exhausted, generating, on the average, 100 F₂ progeny per F₁ animal. One-half of the worms by volume were removed from each well to make genomic DNA; of these, one-half were transferred to equivalent wells of V-bottom 96-well microtiter plates and one-half were pooled with other samples from the same 96-well plate (plate pools) into microfuge tubes. Worms in V-bottom wells were digested for up to 16 hr with 10 µl/well of worm lysis buffer containing 50 mM KCl, 10 mM Tris-Cl (pH 8.3), 2.5 mM MgCl₂, 0.45% Tween 20, 0.45% NP-40, 0.01% (wt/vol) gelatin and 200 µg/ml proteinase K. These worm DNA plates were stored without further purification at –80°C. Plate pools were digested for 4 hr with 0.4 ml of lysis

buffer containing 100 mM NaCl, 100 mM Tris-Cl (pH 8), 50 mM EDTA, 1% SDS, 1% β -mercaptoethanol and 100 μ g/ml proteinase K (Birren et al. 1997). Plate pool genomic DNA was then isolated with the Purgene kit (Gentra Systems) according to the manufacturer's protocol, followed by ethanol precipitation and transferral to sealed microtiter plates for storage at 4°C. The remaining half of the worms in the original plates were frozen in the wells by adding an equal volume of freezing solution, mixing, and graduated freezing to -80°C (Sulston and Hodgkin 1988). Survival following freezing was measured by counting motile animals thawed from 10 selected wells from each of 2 plates, and libraries were discarded if <100 viable animals per well on average were recovered.

PCR Screening for Deletions

Genomic DNA plate pools were screened by nested PCR with AmpliTaq Gold (Perkin-Elmer) and primer concentrations of 50 μ M for 35 cycles of 94°C for 30 sec, 58°C for 1 min, and 72°C for 1.5–2.5 min. Aliquots (1 μ l) of template DNA from two libraries (48 plate pools each) were added to a 96-well PCR plate, and up to 8 libraries were screened simultaneously for a given gene target in a Tetrad thermocycler (MJ Research, Watertown, MA). Potential deletions detected by gel electrophoresis of PCR products were confirmed by quadruplicate PCR of the original library plate pool. On identification of a positive plate, PCR was performed on the unpooled DNA plates to identify the positive well. Primary screen data and deletion sizes were available for 127 screened targets.

Isolation of Mutant Animals and Deletion Analysis

Deletions were mapped by restriction enzymes on the basis of the known genomic DNA cosmid sequence, and the mapping information was used to design three new PCR primers, two of which flanked the deletion and a third from within the deletion. Worms were thawed from the positive wells onto NGM plates and then picked individually into microtiter plate liquid cultures. Following growth and reproduction in liquid medium, aliquots of progeny were removed for PCR with flanking primers to identify clones bearing the deletion mutation. Positive lines were transferred to agar plates, and single worms were tested by multiplex PCR with both the flanking and internal primers to distinguish homozygous from heterozygous animals in subsequent generations. Mutant animals were subsequently outcrossed with wild-type N_2 animals to remove other potential mutations.

ACKNOWLEDGMENTS

We thank Bob Barstead and Ron Plasterk for sharing their technical results, Elizabeth Link and Marie Sutherlin for reviewing the manuscript, the *C. elegans* Genome Sequencing Consortium for primary sequence data and on-line search engines, and our many collaborators in the *C. elegans* community who provided biological and genetic expertise for the selection of targets.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

REFERENCES

Ahringer, J. 1997. Turn to the worm. *Curr. Opin. Genet. Dev.* **7**: 410–415.
 Altschul, S.F., W. Gish, W. Miller, E.W. Myers, and D.J. Lipman.

1990. Basic local alignment search tool. *J. Mol. Biol.* **215**: 403–410.
 Anderson, P. 1995. Mutagenesis. in *Caenorhabditis elegans: Modern biological analysis of an organism* (ed. H.F. Epstein and D.C. Shakes), pp. 31–58. Academic Press, San Diego, CA.
 Anderson, P. and S. Brenner. 1984. A selection for myosin heavy chain mutants in the nematode *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci.* **81**: 4470–4474.
 Birren, B., E.D. Green, S. Klapholz, R.M. Myers, and J. Roskams. 1997. *Genome analysis: A laboratory manual*. Vol. 1: *Analyzing DNA*, pp. 20–22. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 Blaxter, M.L. 1998. *Caenorhabditis elegans* is a nematode. *Science* **282**: 2041–2046.
 Blumenthal, T. and K. Steward. 1997. RNA processing and gene structure. In *C. elegans II* (ed. D.L. Riddle, T. Blumenthal, B.J. Meyer, and J.R. Priess), pp. 117–145. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
 The *C. elegans* Sequencing Consortium. 1998. Genome sequence of the nematode *C. elegans*: A platform for investigating biology. *Science* **282**: 2012–2018.
 De Stasio, E., C. Lephoto, L. Azuma, C. Holst, D. Stanislaus, and J. Uttam. 1997. Characterization of revertants of *unc-93 (e1500)* in *Caenorhabditis elegans* induced by N-ethyl-N-nitrosourea. *Genetics* **147**: 597–608.
 Dujon, B. 1998. European Functional Analysis Network (EUROFAN) and the functional analysis of the *Saccharomyces cerevisiae* genome. *Electrophoresis* **19**: 617–624.
 Fire, A., S. Xu, M.K. Montgomery, S.A. Kostas, S.E. Driver, and C.C. Mello. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**: 806–811.
 Friedberg, E.C., G.C. Walker, and W. Siede. 1995. DNA Repair and Mutagenesis. ASM Press, Washington, D.C.
 Gil, E., E.M. Link, L.X. Liu, C.D. Johnson, and J.A. Lees. 1999. Regulation of the insulin-like developmental pathway of *Caenorhabditis elegans* by a homolog of the PTEN tumor suppressor gene. *Proc. Natl. Acad. Sci.* **96**: 2925–2930.
 Gregoire, F.M., N. Chomiki, D. Kachinskas, and C.H. Warden. 1998. Cloning and developmental regulation of a novel member of the insulin-like gene family in *Caenorhabditis elegans*. *Biochem. Biophys. Res. Comm.* **249**: 385–390.
 Hart, A.C., J. Kass, J.E. Shapiro, and J. Kaplan. 1999. Distinct signalling pathways mediate touch and osmosensory responses in a polymodal sensory neuron. *J. Neurosci.* **19**: 1952–1958.
 Hieter, P. and M. Boguski. 1997. Functional genomics: It's all how you read it. *Science* **278**: 601–602.
 Hobert, O., K. Tessmar, and G. Ruvkun. 1999. The *C. elegans* *lim-6* LIM homeobox gene regulates neurite outgrowth and function of particular GABAergic neurons. *Development* **126**: 1547–1562.
 Jansen, G., E. Hazendonk, K.L. Thijssen, and R.H.A. Plasterk. 1997. Reverse genetics by chemical mutagenesis in *Caenorhabditis elegans*. *Nat. Genet.* **17**: 119–121.
 Jansen, G., K.L. Thijssen, P. Werner, M. van der Horst, E. Hazendonk, and R.H.A. Plasterk. 1999. The complete family of genes encoding G proteins of *Caenorhabditis elegans*. *Nat. Genet.* **21**: 414–419.
 Li, X. and I. Greenwald. 1997. HOP-1, a *Caenorhabditis elegans* presenilin, appears to be functionally redundant with SEL-12 presenilin and to facilitate LIN-12 and GLP-1 signaling. *Proc. Natl. Acad. Sci.* **94**: 12204–12209.
 Miklos, G.L.G. and G.M. Rubin. 1996. The role of the genome project in determining gene function: Insights from model organisms. *Cell* **86**: 521–529.
 Mushegian, A.R., D.E. Bassett Jr., M.S. Boguski, P. Bork, and E.V. Koonin. 1997. Positionally cloned human disease genes: Patterns of evolutionary conservation and functional motifs. *Proc. Natl. Acad. Sci.* **94**: 5831–5836.
 Mushegian, A.R., J.G. Garey, J. Martin, and L.X. Liu. 1998. Large-scale taxonomic profiling of eukaryotic model organisms: A comparison of homologous and orthologous proteins encoded

- by the human, fly, nematode, and yeast genomes. *Genome Res.* **8**: 590–598.
- Oliver, S.G. 1996. From DNA sequence to biological function. *Nature* **379**: 597–600.
- Park, E.C. and H.R. Horvitz. 1986. Mutations with dominant effects on the behavior and morphology of the nematode *Caenorhabditis elegans*. *Genetics* **113**: 821–852.
- Plasterk, R.H.A. 1995. Reverse genetics: From gene sequence to mutant worm. In *Caenorhabditis elegans: Modern biological analysis of an organism* (ed. H.F. Epstein and D.C. Shakes), pp. 59–80. Academic Press, San Diego, CA.
- Pulak, R.A. and P. Anderson. 1988. Structure of spontaneous deletions in *Caenorhabditis elegans*. *Mol. Cell. Biol.* **8**: 3748–3754.
- Rand, J.B. and C.D. Johnson. 1995. Genetic pharmacology: Interactions between drugs and gene products in *Caenorhabditis elegans*. In *Caenorhabditis elegans: Modern biological analysis of an organism* (ed. H.F. Epstein and D.C. Shakes), pp. 187–204. Academic Press, San Diego, CA.
- Shaham, S. 1998. Identification of multiple *Caenorhabditis elegans* caspases and their potential roles in proteolytic cascades. *J. Biol. Chem.* **273**: 35109–35117.
- Sluder, A.E., S.W. Mathews, D. Hough, V.P. Yin, and C.V. Maina. 1999. The nuclear receptor superfamily has undergone extensive proliferation and diversification in nematodes. *Genome Res.* **9**: 103–120.
- Sonnhammer, E.L.L. and R. Durbin. 1997. Analysis of protein domain families in *Caenorhabditis elegans*. *Genomics* **46**: 200–216.
- Sulston, J. and J. Hodgkin. 1988. Methods. In *The nematode Caenorhabditis elegans* (ed. W.B. Wood), pp. 587–606. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Westlund, B., D. Parry, R. Clover, M. Basson, and C.D. Johnson. 1999. Reverse genetic analysis of *Caenorhabditis elegans* presenilins reveals redundant but unequal roles for *sel-12* and *hop-1* in Notch-pathway signalling. *Proc. Natl. Acad. Sci.* **96**: 2497–2502.
- Yandell, M.D., L.G. Edgar, and W.B. Wood. 1994. Trimethylpsoralen induces small deletion mutations in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci.* **91**: 1381–1385.
- Zwaal, R.R., A. Broeks, J. van Meurs, J.T.M. Groenen, and R.H.A. Plasterk. 1993. Target-selected gene inactivation in *Caenorhabditis elegans* by using a frozen transposon insertion mutant bank. *Proc. Natl. Acad. Sci.* **90**: 7431–7435.

Received May 18, 1999; accepted in revised form July 15, 1999.

High-Throughput Isolation of *Caenorhabditis elegans* Deletion Mutants

Leo X. Liu, Jill M. Spoerke, Evan L. Mulligan, et al.

Genome Res. 1999 9: 859-867

Access the most recent version at doi:[10.1101/gr.9.9.859](https://doi.org/10.1101/gr.9.9.859)

References

This article cites 28 articles, 18 of which can be accessed free at:
<http://genome.cshlp.org/content/9/9/859.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Focused on your science.

USC
SCIENTIFIC
SIDE BY SCIENCE

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>