

VOLUME 1, SUBJECT INDEX

- Absorbance ratios, effect of guanidinium-phenol-chloroform RNA extraction on in RT-PCR, 286–290
- Acholeplasma, method of PCR detection and RFLP differentiation based upon 16S rRNA genes, 202–204
- Affinity capture, sequencing templates derived from, 222–228
- Allele-specific DNA amplification, 5–16
- Allele-specific oligonucleotide hybridization, detection of mutated *ras* genes, 211–216
- Allele-specific PCR amplification
detection of mutated *ras* genes, 211–216
HLA typing, 91–98
- Alleles, preferential PCR amplification of, 241–250
- Allelic differences, single nucleotide primer extension as quantitative assay for, 160–163
- Alu-PCR products, cloning of using uracil DNA glycosylase, 120–123
- Amplicons, generation of in cold cycle, 1–4
- Amplification, see also specific names
isothermal transcription-based, 25–33
preferential, of alleles, 241–250
single tube, detection and identification of multiple mycobacterial pathogens, 269–273
systems, comparison of, 25–33
- Anchor PCR, cloning of genomic DNA sequences flanking transgene integration site, 129–135
- Archeological remains, amplification of DNA from 107–110
- Archeology, molecular, 107–110
- Archival material, paraffin-embedded, PCR amplification from, 46–50
- Artifacts
fragments, in PCR product, method to eliminate, 171–174
generation of with degenerate oligonucleotide sequences, 57–62
RT-PCR, from processed pseudogenes, 70–71
- Aspartylglucosaminidase mRNA, method for determining amount of in tissues, 234–240
- Asymmetric PCR, sequencing templates derived from, 222–228
- Automation, of genetic screening methods, 91–98
- Azidothymidine, quantitative amplification method to assess antiviral activity in cell culture, 257–262
- Blood, see also Guthrie spots
dried, PCR analysis of DNA from, 91–98
preparation of DNA from for PCR screening of transgenes, 72–74
- Bovine corneal protein 54, and PCR artifacts generated in cloning of using DOS, 57–62
- Branch-capture reaction, 5–16
- Capture PCR, and isolation of DNA fragments adjacent to known sequence, 111–119
- cDNA inserts, PCR protocol for isolating, 195–198
- cDNA library, PCR protocol for identifying differentially expressed genes from, 195–198
- cDNA signals, amplified, ratio of as reliable measure of test gene expression, 164–170
- Chain-termination, compared with chemical degradation method for sequencing PCR-amplified DNA, 222–228
- Chemical degradation, compared with chain-termination method for sequencing PCR-amplified DNA, 222–228
- Chemical genetics, see Vectorette PCR
- Chromosomal DNA fragment, cloning with ligase chain reaction, 5–16
- Chromosome 7, human, YAC-based STS content mapping of cystic fibrosis region, 77–90
- Chromosomes
microdissected, PCR-mediated cloning of *HpaII* tiny fragments from, 229–237
STS content mapping, 77–90
- Cloning
Alu-PCR, using uracil DNA glycosylase, 120–123
genomic DNA sequences flanking transgene integration site using single-site PCR, 129–135
PCR-mediated, *HpaII* tiny fragments from microdissected human chromosomes, 222–228
PCR products, rapid and reliable methods, 140–141
of full-length Sendai virus mRNA, 293–296
- Clostridium perfringens*, DNA probe and PCR primers for, 51–56
- Cold deoxyribonuclear fusion, 1–4
- Competitor DNA fragments, generation of for quantitative PCR, 136–139
- Computer program, OSP for PCR and DNA sequencing primer selection, 120–123
- Contamination
DNA, single-step elimination of before RT-PCR, 279–282
in archeological specimens, 107–110
in environmental microbiology samples, 151–159
minimization with capture PCR method, 111–119
sources of in RT-PCR, 70–71
- CPCR, see Capture PCR
- CYP21B gene, selective amplification and sequencing of, 181–186
- Cystic fibrosis, newborn screening for, 91–98
- Cystic fibrosis gene, multiplex PCR for rapid detection of mutations, 297–298
- Cystic fibrosis region, human chromosome 7, YAC-based STS content mapping of, 77–90
- Degenerate oligonucleotide sequences, use of in PCR and generation of artifacts, 57–62
- Denaturants, use of for sequencing PCR products, 222–228
- Deoxyribonuclear fusion, cold, 1–4
- Detection
influenza A and B viruses in respiratory secretions, 263–268
multiple mycobacterial pathogens in single tube, 269–273
- Dideoxy sequencing, direct, of double-stranded PCR products, 255–256
- DNA
adjacent to known sequence, cloning of using single-site PCR, 129–135
allele-specific, quantitative assay of with SNUPE, 160–163
amplification from archeological remains, 107–110
ancient, PCR-based methods of study, 217–221
damage of in PCR, 17–24
double-stranded, amplification, 171–174
fossil, PCR-based study of, 107–110
genomic, direct amplification from autoclaved infectious microorganisms, 74
method for amplification from dried blood spots, 91–98
method for microextraction from dried blood spots, 91–98
mitochondrial, PCR-based methods of study, 217–221
PCR-amplified, sequencing methods, 222–228
polymorphic, random amplification of for fingerprinting plants, 274–276

- preparation from whole blood of mice for PCR screening of transgenes, 72–74
- DNA amplification, allele-specific, 5–16
- DNA analysis, PCR, from dried blood spots of filter paper blotters, 91–98
- DNA detection, ligase mediated, 5–16
- DNA diagnostics, use of ligase chain reaction, 5–16
- DNA fragments, adjacent to known sequence, amplification of with capture PCR, 111–119
- DNA ligases
characterization, 5–16
cloning, 5–16
- DNA microextraction, from plant leaf discs, 175–180
- DNA polymerase
and fidelity in PCR, 17–24, 63–69
error discrimination by, 17–24
thermostable, in environmental microbiology, 151–159
- DNA probes, 16s/23s ribosomal spacer region as target for in eubacteria, 51–56
- DNA sequence analysis, detection of mutated *ras* genes, 211–216
- DOS, see Degenerate oligonucleotide sequences
- Double-stranded PCR product, sequencing methods, 222–228
- Duchenne muscular dystrophy, newborn screening for, 91–98
- Environmental microbiology, and PCR, 151–159
- Enzymes, 3SR, 25–33. See also *specific names*
- Erratum, 149, 210, 299
- Error discrimination, during PCR, 17–24
- Error rates, in PCR, 17–24
- Eubacteria, 16s/23s ribosomal spacer region as target for DNA probes, 51–56
- Evolution
domesticated species, PCR-based study, 107–110
PCR, direct sequencing, and the comparative approach, 217–221
- Evolutionary analysis, PCR-based, for HLA polymorphism studies, 91–98
- 5' end, mRNA, determination with PCR-enhanced method, 43–45
- Fidelity, optimization of in PCR, 63–69
- Fingerprinting, plants, random amplification of DNA for, 274–276
- Fixation methods, and PCR amplification, 46–50
- Forensics
HLA typing in, 91–98
identification, from dried blood specimens, 91–98
- Gene expression
analysis with RNA-PCR, 283–285
in environment, PCR detection, 151–159
reporter, ratio of amplified cDNA signals as reliable measure of, 164–170
- Genes, differentially expressed, PCR protocol for identifying from cDNA library, 195–198
- Genetic screening, automation of, 91–98
- Genetically engineering microorganism, PCR detection of those released in the environment, 151–159
- Genome
human, isolation of mapping of regions with YAC clones, 77–90
plant, RAPD and other PCR-based analyses from DNA extracted from small leaf discs, 175–180
- Genomic DNA template, double-stranded, PCR amplification and direct sequencing, 171–174
- Genomic walking, use of vectorette PCR, 39–42
- Giardia, PCR detection, 151–159
- Glyceraldehyde-3-phosphate dehydrogenase, use of oligonucleotide primers for, 283–285
- Glycoprotein phosphorylation sites, herpesvirus, site-directed mutagenesis of using recombination PCR, 205–207
- Guanidinium-phenol-chloroform RNA extraction, effect of on A260/A280 and accuracy of mRNA quantitation by RT-PCR, 286–290
- Guthrie spots
and automation of genetic screening methods, 91–98
PCR DNA analysis of, 99–106
- Hepatitis C virus, direct amplification of RNA from human serum, 291–292
- HLA class II typing, PCR based, 91–98
- HLA typing, uses of, 91–98
- HLA-DQ α locus, preferential amplification of alleles at, 241–250
- Hot start PCR, 1–4
use in environmental microbiology, 151–159
- Hpa*II tiny fragments, PCR-mediated cloning of from microdissected human chromosomes, 229–237
- HTF, see *Hpa*II tiny fragments
- Human immunodeficiency virus, quantitative gene amplification procedure to monitor infection and antiviral activity in cell culture, 257–262
- Human leukocyte antigen class II molecules, PCR-based typing, 91–98
- Infectious disease, diagnosis from dried blood specimens, 91–98
- Influenza A and B viruses, detection in respiratory secretions with PCR, 263–268
- Insertion mutations, rapid mapping of with REP-Tn5 PCR, 187–194
- J-PCR, see Junction PCR
- Junction-PCR, analysis of transgenic animal pedigrees, 208–210
- Klenow-kinase-ligase, method of cloning PCR products, 140–141
- Legionella*, PCR detection, 151–159
- Ligase chain reaction, 5–16
compared with 3SR and PCR, 25–33
methods, 5–16
variations, 5–16
- Linker and adaptor cloning, method for PCR products, 140–141
- Microbiology, environmental, and PCR, 151–159
- Microorganisms
autoclaved infectious, direct amplification of genomic DNA from, 74
identification with PCR-based methods, 217–221
indicator, PCR detection of, 151–159
indigenous, PCR detection, 151–159
viable but nonculturable, PCR detection in the environment, 151–159
- Minisequencing, solid-phase, and PCR, quantification of mRNA by, 234–240
- Misincorporation, in PCR product, method to eliminate, 171–174
- Mitochondrial DNA
amplification of from archeological remains, 107–110
PCR-based methods of study, 217–221
- Molecular archeology, 107–110
- Molecular mechanisms, value of comparative studies, 217–221
- Mollicutes, method of PCR detection and RFLP differentiation based upon 16S rRNA genes, 202–204
- mRNA
determination of approximate molar concentration using competitor DNA fragments, 136–139
determination of 5' end sequence with PCR-enhanced method, 43–45
quantification by PCR and solid-phase minisequencing, 234–240
semi-quantitative PCR of, 144–145

- Multigene families, method for specific amplification and PCR sequencing, 181–186
- Multiplex amplification system and environmental monitoring of microorganisms, 151–159
 identification of multiple mycobacterial pathogens, 269–273
 for rapid detection of cystic fibrosis gene, 297–298
- Mutagenesis, site-directed, of glycoprotein phosphorylation sites by recombination PCR, 205–207
- Mutant allele frequency, ascertainment from dried blood specimens, 91–98
- Mutant-enriched PCR, detection of mutated *ras* genes, 211–216
- Mutations
 activating *ras*, screening for with PCR-primer introduced restriction analysis, 146–148
 characterization with PCR-SSCP, 34–38
 cystic fibrosis gene, multiplex PCR for rapid detection of, 297–298
E. coli::Tn5 insertion, rapid mapping of with REP-Tn5 PCR, 187–194
 point, identification of in p53 cDNAs by direct PCR sequencing, 199–201
ras, PCR-based approaches for detection, 211–216
- Mycobacterial pathogens, multiple, detection of in single tube, 269–273
- Mycoplasma, method of PCR detection and RFLP differentiation based upon 16S rRNA genes, 202–204
- Newborn screening, PCR-based, 91–98.
 See also Guthrie spots
- Nuclear sequences, amplification of from archeological remains, 107–110
- Nucleic acids, purification from environmental microorganisms for PCR amplification, 151–159
- Oligonucleotide Selection Program, computer program for PCR and DNA sequencing primers, 120–123
- Oligonucleotide typing, HLA class II genes, 91–98
- OSP, see Oligonucleotide Selection Program
- p53 cDNAs, identification of point mutations in by direct PCR sequencing, 199–201
- p53 pseudogene, detection of using PCR, 251–254
- Paraffin-embedded tissues, PCR amplification from, 46–50
- Pathogens, water-borne, PCR detection, 151–159
- PCR, compared with 3SR and LCR, 25–33
- PCR products
 rapid and reliable cloning methods, 140–141
 sequencing of, 222–228
- PCR templates, dry, slow inactivation of by UV light, 142–143
- PCR-PIRA, see PCR-primer introduced restriction analysis
- PCR-primer introduced restriction analysis, screening for activating *ras* mutations with, 146–148
- PCR-restriction fragment length polymorphism, difficulties of for HLA typing, 91–98
- PCR-single-stranded conformation polymorphism, 34–38
- Phosphorylation sites, herpesvirus glycoprotein, site directed mutagenesis of using recombination PCR, 205–207
- Phylogeny
 application of PCR methods to, 217–221
 reconstruction with PCR-based methods, 217–221
- Plants
 leaf discs, DNA microextraction method, 175–180
 random amplification of polymorphic DNA for fingerprinting, 274–276
- Point mutations, identification of in p53 cDNAs by direct PCR sequencing, 199–201
- Polymerase chain reaction. See PCR; *specific methods*
- Polymorphic DNA, random amplification of for fingerprinting plants, 274–276
- Polymorphisms, HLA class II, PCR detection, 91–98
- Population studies, PCR-based, 107–110
- Preferential amplification, of alleles, 241–250
- Primer introduced restriction analysis, and PCR, method for screening for activating *ras* mutations, 146–148
- Primers
 computer program for selection of, 120–123
 for *Clostridium perfringens*, 51–56
 for Sendai virus mRNA cloning, 293–296
 oligonucleotide, use of for glyceraldehyde-3-phosphate dehydrogenase, 283–285
- Pseudogene, processed
 artifacts from in RT-PCR, 70–71
 detection of using PCR, 251–254
- Quantitation
 allelic differences, using single nucleotide primer extension assay, 160–163
 gene amplification procedure to assess HIV infection and antiviral activity in cell culture, 257–262
 generation of competitor DNA fragments for, 136–139
 mRNA, accuracy by RT-PCR, 286–290
 mRNA, by PCR and solid-phase minisequencing, 234–240
 PCR, and mRNAs, 144–145
- Random amplified polymorphic DNA, and analysis of plant genomes from DNA extracted from small leaf discs, 175–180
- RAPD, see Random amplified polymorphic DNA
- ras* genes, mutated, PCR-based methods of detection, 211–216
- ras* mutations, activation, screening for with PCR-primer introduced restriction analysis, 146–148
- Recombination PCR, and site-directed mutagenesis of herpesvirus glycoprotein phosphorylation sites, 205–207
- REP-Tn5 PCR, and rapid mapping of *E. coli*::Tn5 insertion mutations, 187–194
- Reporter gene expression, PCR-based assay for, 164–170
- Restriction fragment length polymorphism, detection of mutated *ras* genes, 211–216
- Reverse transcriptase PCR
 and accuracy of mRNA quantitation, 286–290
 artifacts from processed pseudogenes, 70–71
 elimination of contaminating DNA before, 279–282
- Ribosomes, 16s/23s spacer region as target for DNA probes in eubacteria, 51–56
- RNA
 allele specific, quantitative assay of with SNuPE, 160–163
 hepatitis C virus, direct amplification from human serum, 291–292
- RNA-PCR, and use of oligonucleotide primers for glyceraldehyde-3-phosphate dehydrogenase, 283–285
- RNase mismatch cleavage, detection of mutated *ras* genes, 211–216
- RPCR, see Recombination PCR
- RT, see Reverse transcriptase
- 16S rRNA genes, method of PCR detec-

- tion and RFLP differentiation of Mollicutes based upon, 202–204
- 16s/23s ribosomal spacer region, eubacteria, as target for DNA probes, 51–56
- Self-sustained sequence replication, 25–33
 compared with PCR and LCR, 25–33
- Semi-quantitative PCR, of mRNAs, 144–145
- Sendai virus mRNA, full-length, PCR cloning of, 293–296
- Sensitivity
 PCR, conditions for increase of, 1–4
 PCR-SSCP, 34–38
- Sequence data, DNA, value of comparative studies, 217–221
- Sequence-tagged site content mapping, human chromosomes, 77–90
- Sequencing
 direct, of double-stranded PCR products, avoiding strand reassociation in, 193–194
 direct, of PCR-amplified double-stranded genomic DNA template, 171–174
 direct dideoxy, and sequencing of double-stranded PCR products, 255–256
 direct PCR, and identification of point mutations in p53 cDNAs, 199–201
 from single-stranded templates derived from PCR products, 222–228
 PCR-amplified DNA, 222–228
 PCR-based, CYP21B gene, 181–186
 PCR-based, multigene families, 181–186
 sensitivity of for mixed allelic samples, 199–201
- Single nucleotide primer extension assay, and quantitative measurement of allelic differences, 160–163
- Single-site PCR
 and cloning of genomic DNA sequences flanking transgene integration site, 129–135
 comparison with anchor PCR, 129–135
- Single-stranded conformation polymorphism, 34–38
 detection of mutated *ras* genes, 211–216
- Single-stranded DNA binding proteins, use of for sequencing PCR products, 222–228
- Site-directed mutagenesis, herpesvirus glycoprotein phosphorylation sites, use of recombination PCR for, 205–207
- SNuPE, see Single nucleotide primer extension
- Soil, purification of nucleic acids from, 151–159
- Solid-phase minisequencing, and PCR, quantification of mRNA by, 234–240
- Spiroplasma, method of PCR detection and RFLP differentiation based upon 16S rRNA genes, 202–204
- SSCP, see Single-stranded conformation polymorphism
- Steroid 21-hydroxylase deficiency, PCR sequencing analysis of multigene family, 181–186
- Strand reassociation, avoiding in direct sequencing of double-stranded PCR products, 193–194
- STS, see Sequence-tagged site
- Systematics, molecular, application of PCR methods to, 217–221
- 3' overhangs, generation of with UDG, 120–123
- 3SR, see Self-sustained sequence replication
- T7
 DNA polymerase, modified, and fidelity in PCR, 63–69
 promoter sequence and amplimer, sequencing templates derived from, 222–228
- Taq* DNA polymerase
 and fidelity in PCR, 63–69
 use of for sequencing PCR products, 222–228
- Taxonomy, use of DNA sequences as molecular tags for species identity, 217–221
- Temperature variation, and preferential amplification, 241–250
- Template preparation, for direct dideoxy sequencing, 255–256
- Thermolabile DNA polymerases, and direct sequencing of double-stranded PCR products, 193–194
- Thermophilic organisms, 5–16
- Thermostable enzymes, in ligase chain reaction, 5–16. See also *Taq* DNA polymerase; Vent DNA polymerase
Thermus aquaticus, see also *Taq*
- Tissues, paraffin-embedded, PCR amplification from, 46–50
- Tn5 insertion mutations, rapid mapping of with REP-Tn5 PCR, 187–194
- Transcription, isothermal, amplification system, 25–33
- Transgene
 integration site, cloning of genomic flanking DNA sequences using single-site PCR, 129–135
 method for preparation of DNA from blood of mice for screening, 72–74
- Transgenic animals, analysis of with junction-PCR, 208–210
- Transplantation, use of HLA typing for donor selection, 91–98
- UDG, see Uracil DNA glycosylase
- Ultraviolet light, and slow inactivation of dry PCR templates, 142–143
- Uracil DNA glycosylase, cloning of Alu-PCR products using, 120–123
- Ureaplasma, method of PCR detection and RFLP differentiation based upon 16S rRNA genes, 202–204
- Variable number tandem repeat, preferential amplification at, 241–250
- Vectorette PCR, 39–42
- Vectors, cloning of PCR products into and sequencing, 222–228
- Vent DNA polymerase, and fidelity in PCR, 63–69
- Vibrio vulnificus*, PCR detection, 151–159
- Virus, influenza A and B, PCR detection in respiratory secretions, 263–268
- VNTR, see Variable number tandem repeat
- Water, purification of nucleic acids from microorganisms in, 151–159
- YAC, see Yeast artificial chromosomes
- Yeast artificial chromosomes
 and STS content mapping, 77–90
 use of capture PCR in, 111–119
 use of vectorette PCR for sequencing termini, 39–42

VOLUME 1, AUTHOR INDEX

- Abramow-Newerly, W., 208
Allen, M., 91
Arnold, C., 39
Balachandran, R., 277
Baptist, E.W., 57
Barany, F., 5
Barry, T., 51, 75
Basile, A.A., 283
Baur, W., 286
Bej, A.K., 151
Belloni, E., 297
Betts, R.F., 263
Bevan, I.S., 222
Blankenstein, T., 136
Bos, J., 211
Brian, David A., 43
Cariani, E., 291
Chang, Y.F., 70
Cheng, J., 199
Clapoff, S., 208
Classen, U., 229
Colleran, G., 51
Collier, S., 181
Cone, R.W., 142
Connolly, R., 286
Coonrod, D., 263
Cooper, D.L., 57
Cousins, D., 269
Cremonesi, L., 297
Cruess, D.F., 286
Dai, A., 160
Davidson, J.N., 263
Deng, S., 202
Deragon, J.M., 175
Diamantstein, T., 136
Dias, C., 63
Dieffenbach, C.W., 283, 286
Dilworth, D.D., 279
Donofrio, J.C., 263
Dritschilo, A., 171
Dunican, L.K., 51
Dveksler, G.S., 283
Eckert, K.A., 17
Erlich, G.D., 277
Erlich, H.A., 241
Fahy, E., 25
Fairfax, M.R., 142
Ferrari, M., 297
Ford-Lloyd, B.V., 274
Gannon, F., 51, 75
Gingeras, T.R., 25
Glennon, M., 51
Golay, J., 144
Green, E.D., 77
Green, P., 77, 124
Greer, C.E., 46
Grose, C., 205
Gupta, K.C., 293
Gupta, P., 277
Gyllensten, U., 91
Haas, M., 199
Ham, R.G., 70
Hayashi, K., 34
Hesse, S., 229
Higuchi, R., 241
Hillier, L., 124
Hiruki, C., 202
Hodgson, I.J., 39
Hofmann, M.A., 43
Holodniy, M., 257
Horsthemke, B., 229
Hulla, J.E., 251
Ikonen, E., 234
Introna, M., 144
Jacobson, D.R., 146
Jones, D.H., 205
Jung, M., 171
Kasid, U., 171
Katzenstein, D.A., 257
Keohavong, P., 63
Kere, J., 255
Klemish, S.W., 195
Kocher, T.D., 217
Kunkel, T.A., 17
Kwoh, D.Y., 25
Lagerström, M., 111
Landegren, U., 111
Landry, B.S., 175
LeBon, J.M., 160
Ling, L.L., 63
Lorens, J.B., 140
Lüdecke, H.J., 229
Lund, J.K., 46
Lupski, J.R., 187
MacGregor, G.R., 129
Magnani, C., 297
Mahbubani, M.H., 151
Malmgren, H., 111
Manninen, T., 234
Manos, M.M., 46
McCabe, E.R.B., 99, 187
McCarrey, J.R., 279
Menon, R.S., 70
Merigan, T.C., 257
Metcalf, M.A., 142
Mischke, D., 193
Montanaro, V., 255
Moskovits, T., 146
Mullis, K.B., 1
Munthali, M., 274
Mutchler, K.J., 195
Newbury, H.J., 274
Nisson, P.E., 120
Overbeek, P.A., 129
Pääbo, S., 107
Parik, J., 111
Passerini, F., 144
Peltonen, L., 234
Pettersson, U., 111
Platzer, C., 136
Primi, D., 291
Rapley, R., 222
Rashtchian, A., 120
Ravaggi, A., 291
Riggs, A.D., 160
Robertson, J.A., 202
Roder, J.C., 208
Russo, A.F., 195
Seia, M., 297
Shankarappa, B., 277
Sharefkin, J.B., 286
Singer-Sam, J., 160
Srivastava, A.K., 255
St. Clair, J., 70
Stemke, G.W., 202
Stewart, J., 111
Strachan, T., 181
Subramanian, P.S., 187
Syvänen, A-C., 234
Tassabehji, M., 181
Thilly, W.G., 63
Tilmans, I., 193
Überla, K., 136
van Mansfeld, A.D.M., 211
Versalovic, J., 187
Walker, M.R., 222
Walsh, P.S., 241
Wanner, R., 193
Ward, M., 208
Watkins, P.C., 120
Wilton, S., 269
Winbert, G., 72
Winters, M.A., 257
Xu, X., 293
Yamaguchi, M., 286
Yao, Z., 205


Custom

Since 1984

PCR* PRIMERS

Left Primer

Bio-Synthesis
5'-CAT ACGAT-3'



FULL SERVICE FOR YOUR PCR WORK
PRIMER/PROBE DESIGN
BIOTIN/ALKAL-PHOS. OLIGOS


- . Full Technical Support
- . 48 hrs delivery
- . Purified Oligos
- . Gel picture Q.C on each Oligo
- . No set up charge
- . 10.0 O.D's (>330 µgs) delivered



Includes **4.** **50**
Base

Right Primer

Bio-Synthesis
5'-ACT CTCATC-3'



* PCR process covered by Hoffman LaRoche patents.


THE BIO-LADDER™

A complete ladder for PCR*, defines a new standard of accuracy in DNA size analysis.

" Sizing DNA like never before !"
BIO-SYNTHESIS BIO-LADDER gives you clean, sharp, bright bands in precise 100 bp increments.

- * 50-1,000 bp, an optimum range for amplified products.
- * Bands in increments of 100 to facilitate sizing, even includes a 50 bp band.
- * Extra bright 500 bp band ensures easy orientation on gel.
- * Ready to load - no preheating needed.
- * Two convenient sizes :

Cat # 54-1020	20 runs	\$40. ⁰⁰
# 54-1050	50 runs	\$75. ⁰⁰



Gel picture. A 2.5 % agarose gel was loaded with 5µl(1 run) of BIO-LADDER. Gel is run at 100 volts in 0.5 TBE (Tris Borate-EDTA) buffer containing ethidium bromide at 1.0 µg/ml.
Lane 1: Bio-Ladder sizing marker
Lane 2: 95 bp HLA DQα gene amplified product
Lane 3: 324 bp HLA DQα gene amplified product

BIO-SYNTHESIS, INC. (800) 227-0627

Fax: (214) 420-0442

In TX: (214) 420-8505

Results...



Pure and Simple.

Pure... NuSieve® GTG agarose. A one-of-a-kind, high purity, low melting temperature agarose which resolves small DNA fragments of 10 bp to 1 kb. Ideal for separating small DNA, including PCR products. Yields "biologically active" DNA for reliable cloning in the presence of remelted agarose gel.

Simple... NuSieve GTG agarose. An easy-to-cast, one-component gel system which shortens DNA prep time and eliminates DNA recovery steps.

Only from FMC... NuSieve GTG agarose. One of FMC's certified Genetic Technology Grade™ product line which is compatible with such procedures as restriction digestions, ligations and blotting. Ask for technical information today.

For Research Only. Not for use in diagnostic procedures. NuSieve GTG agarose — U.S. Patent No. 3,956,273 — FMC Corporation. **FMC**, NuSieve and Genetic Technology Grade are trademarks of FMC Corporation. Copyright © 1990 FMC Corporation. All Rights Reserved. The GeneAmp™ Polymerase Chain Reaction (PCR) Process — U.S. Patent No. 4,683,202 — Cetus Corporation

FMC

FMC BioProducts, 5 Maple Street, Rockland, ME 04841, 800-521-0390
FMC BioProducts Europe, Risingevej 1, DK-2665 Vallensbaek Strand, Denmark, 45-42-73-11-22

Reader Service No. 216