

Analysis and correction of crosstalk effects in pathway analysis

Michele Donato¹, Zhonghui Xu², Alin Tomoiaga³, James G. Granneman⁴,
Robert G. MacKenzie⁴, Riyue Bao⁵, Nandor Gabor Than⁶, Peter H. Westfall³,
Roberto Romero², Sorin Draghici^{1,7*}

¹Computer Science Department, Wayne State University

²Perinatology Research Branch, NICHD/NIH

³Center for Advanced Analytics and Business Intelligence, Texas Tech University

⁴Center for Integrative Metabolic and Endocrine Research, Wayne State University

⁵Department of Biological Sciences, Wayne State University

⁶Department of Obstetrics and Gynecology, Wayne State University, School of Medicine

⁷Department of Clinical and Translational Science, Wayne State University

July 30, 2013

*To whom the correspondence should be addressed. E-mail: sorin@wayne.edu, Fax: +1-313-577-6679

Abstract

Identifying the pathways that are significantly impacted in a given condition is a crucial step in understanding the underlying biological phenomena. All approaches currently available for this purpose calculate a p-value that aims to quantify the significance of the involvement of each pathway in the given phenotype. These p-values were previously thought to be independent. Here we show that this is not the case, and that many pathways can considerably affect each other's p-values through a "crosstalk" phenomenon. Although it is intuitive that various pathways could influence each other, the presence and extent of this phenomenon have not been rigorously studied and, most importantly, there is no currently available technique able to quantify the amount of such crosstalk. Here, we show that all three major categories of pathway analysis methods (enrichment analysis, functional class scoring, and topology-based methods) are severely influenced by crosstalk phenomena. Using real pathways and data, we show that in some cases pathways with significant p-values are not biologically meaningful, and that some biologically meaningful pathways with non-significant p-values become statistically significant when the crosstalk effects of other pathways are removed. We describe a technique able to *detect*, *quantify*, and *correct* crosstalk effects, as well as *identify independent functional modules*. We assessed this novel approach on data from four real experiments coming from three phenotypes involving two species. This method is expected to allow a better understanding of individual experiment results, as well as a more refined definition of the existing signaling pathways for specific phenotypes.

Introduction

The correct identification of the signaling and metabolic pathways involved in a given phenotype is a crucial step in the interpretation of high-throughput genomic experiments. Most approaches currently available for this purpose treat the pathways as independent. In fact, pathways can affect each other's p-values through a phenomenon we refer to as *crosstalk*. This crosstalk may be due to the regulatory interactions among different pathways or to the gene overlap among pathways. In this work we will use the term crosstalk to refer to the effect that pathways exercise on each other due to the presence of overlapping genes. Although it is intuitive that various pathways could influence each other, especially when they share genes, the presence and extent of this phenomenon have not been rigorously studied and, most importantly, there is no currently available technique able to quantify the amount of such crosstalk. There are three major categories of methods that aim to identify significant pathways: enrichment analysis (e.g. Fisher's Exact Test - hypergeometric) (Tavazoie et al., 1999; Draghici et al., 2003), functional scoring (e.g. GSEA) (Mootha et al., 2003; Subramanian et al., 2005), and topology-based methods (e.g. impact analysis) (Draghici et al., 2007; Tarca et al., 2009). Another classification of gene set analysis methods is based on the definition of the null hypothesis, and divides the methods into competitive and self-contained (Goeman and Bühlmann, 2007; Nam and Kim, 2008). In this work we focus on competitive methods, and in particular on the Fisher Exact Test, although the problems identified likely apply also for self-contained methods.

Here we show that the results of all these methods are affected by crosstalk effects, and that this phenomenon is related to the structure of the pathways. We propose the first approach that can: i) *detect crosstalk* when it exists, ii) *quantify its magnitude*, iii) *correct for it*, resulting in a more meaningful ranking among pathways in a specific biological condition, and iv) identify *novel functional modules* that can play an independent role and have different functions than the pathway they are currently located on. This method is expected to allow a better understanding of individual experiment results, as well as a more refined definition of the existing signaling pathways for specific phenotypes.

Crosstalk effects

In order to demonstrate the existence and assess the extent of crosstalk effects, we conducted a systematic exploration of this phenomenon. We constructed a reference set of genes from all genes present in at least one KEGG signaling pathway (2,963 genes at the time). Then, each pathway was used as a “bait”, choosing from it a number of genes that would make it significant at a chosen significance level ($\alpha = 0.01$ after Bonferroni correction). Other random genes were selected up to a constant number ($n=100$) of “differentially expressed” (DE) genes (see Supplementary materials for details). Under these circumstances, the research hypothesis is true for the bait, while the null hypothesis is true for all other pathways. We repeated this selection 1,000 times for each pathway P_i , and each time we computed the Fisher Exact Test p-value (Tavazoie et al., 1999), SPIA (impact analysis) (Tarca et al., 2009), and GSEA (Subramanian et al., 2005) p-values for all pathways from the KEGG database (Kanehisa et al., 2004). With these results, we constructed the empirical distributions of the False Discovery Rate (FDR)-corrected p-values corresponding to each prey P_j . The distributions of the p-values for all three methods are severely skewed towards zero, showing that all methods produce a significant number of false positives due to crosstalk effects (Fig. S1 in Supplementary Materials). We hypothesized that crosstalk is mostly due to the common genes between pathways. If this were true, we would expect to see a strong coupling between pairs of pathways with many genes in common and a weak coupling between pathways that do not share any genes. In order to test this hypothesis, we calculated the Jaccard similarity index (Jaccard, 1910) for each pair of pathways, as well as the Pearson correlation between the Jaccard index and the p-values of the preys (Figs. 1 and 2). The data shows a very strong correlation (Pearson correlation index of 0.87 for Fisher’s Exact Test, 0.62 for GSEA and 0.83 for SPIA), which confirms our hypothesis that the crosstalk can be explained by the presence of genes that are involved in more than one pathway.

Results

Crosstalk analysis and correction. The method we propose for correcting for crosstalk effects takes as input a set of reference pathways and a list of genes that are DE in the given condition. The crosstalk analysis is composed of three steps. The first step is the computation of a crosstalk matrix, i.e. the analysis of pairwise crosstalk interactions between pathways in the given condition, resulting in a heat map showing the p-value of each pathway when the genes from each other pathway are removed. This matrix allows the visualization of various cases of crosstalk effects and provides information on the extent of crosstalk effects in the condition analyzed. The second step is the module detection, that allows the identification of sub-pathways that appear to be involved in the specific condition studied independently of the pathway they belong to. The final step is the maximum impact estimation, where the biological impact of each gene is assigned to only one of the pathways to which the gene belongs, for those genes that belong to more than one pathway. The result of the last two steps is essentially a new set of pathways that can include: a) original pathways as found in the literature, b) novel functional modules that might be relevant to the given condition, and c) pathways modified by the removal of a such a functional module. The new set of pathways is not affected by crosstalk effects, and new p-values are computed for each pathway, building a ranked list from which the false positives due to crosstalk have been removed (see Methods).

Fat remodeling in mice. We include here the results of the crosstalk analysis in an experiment investigating cellular and metabolic plasticity of white fat tissue (WAT), where the classical over-representation analysis (ORA) produced a number of false positives, and failed to rank highly pathways that were known to be involved in the given condition. In this experiment, the chronic activation of WAT β -adrenergic receptors by certain physiological and pharmacological conditions transforms the tissue into one resembling brown fat, a thermogenic organ (Granneman et al., 2005; Li et al., 2005; Mottillo et al., 2007). The data set was obtained from a microarray analysis of white fat from mice treated with low dose (0.75 nmol/hr) CL 316,243 (CL) for 0, 3 and 7 days. Here we discuss only the list of DE genes coming from the comparison between expression levels of genes at days 3 and 0 (for the comparison between days 7 and 0, see Supplementary Materials).

The top 20 pathways ranked by ORA and their associated FDR-corrected p-values are shown in Fig. 3a. In this figure, pathways highlighted in red represent pathways not related with the phenomenon in analysis, while pathways highlighted in green are those for which we know, with reasonable confidence, that they are involved in the given phenomenon. The white background indicates pathways for which we do not have conclusive information on their involvement (or lack of) with the phenomenon in analysis. The three most significant pathways in the comparison between days 3 and 0 were *Parkinson's*, *Alzheimer's* and *Huntington's* diseases. The fourth pathway in the ranked list is *Leishmaniasis*. The first three pathways describe degenerative diseases of the central nervous system that have no connection to fat remodeling. *Leishmaniasis* describes the signaling involved in a disease spread by the bite of certain species of sand flies. Clearly, this pathway is also unlikely to give insights about the fat remodeling phenomenon. While other pathways such as *Phagosome* (Newman et al., 1982), *PPAR Signaling* (Granneman et al., 2005), and *Cell cycle* (Lee et al., 2012), are definitely more related to the phenomenon of fat remodeling, their presence in the middle of a ranked list dominated by false positives (6 false positives in the 10 pathways significant at 1%) illustrates how the crosstalk effects make the classical ORA unable to find the truly relevant pathways.

In order to analyze and eliminate the crosstalk effects we computed the *crosstalk matrix* as described in the Methods section. The analysis of the matrix illustrates some interesting examples of crosstalk effects. Fig. 4 represents a detail of the entire matrix. In this figure, the high significance of *Parkinson's* (bright red in row 1, column 1) disappears when the crosstalk due to *Alzheimer's* is eliminated (green in row 1, column 2). This indicates that *Parkinson's* is a false positive, since its significance is due exclusively to genes from *Alzheimer's*. Furthermore, the high significance of *Alzheimer's* (bright red in row 2, column 2) also disappears when the crosstalk effect of *Parkinson's* is eliminated (green in row 2, column 1). This means that *Alzheimer's* significance is also due only to the genes in common with *Parkinson's*. Essentially, the analysis tells us that the genes in common between the two pathways are activated independently of either pathway, which suggests that these genes constitute an independent functional module. The same phenomenon involves the *Cardiac Muscle Contraction* and *Huntington's disease* pathways. The same independent functional module is responsible for the changes shown in areas marked with *a* in Fig. 4.

An inspection of these genes and their signaling mechanisms reveals that this module is composed by genes present in mitochondria, organelles involved in all pathways above. The fact that this module is strongly activated in this fat remodeling experiment that is not related to any of the above conditions (*Alzheimer's*, *Parkinson's*, *Huntington's*), suggests that this should be considered as an independent pathway, dedicated to mitochondrial activity. Fig. 5 shows a representation of this new pathway. In order to investigate the involvement of mitochondria in this condition, epididymal white fat of control and CL-treated (CL-7d) mice were stained with fluorescent Alexa-555 conjugated to streptavidin and imaged by spinning disc confocal microscopy. Fig. 6 shows a comparison between the control (left) and CL-treated mice (right). The right panel of this figure

shows a massive generation of new mitochondria after 7 days of treatment, demonstrating *in vivo* that indeed, the mitochondrial pathway is central in this experiment.

Another very interesting phenomenon can be observed in Fig. 4 (circle *b*). Here, *Toll-like Receptor Signaling* (*TLR* pathway becomes *more significant* when the *Rig-I Like Receptor Signaling* (*RLR* pathway (not Here, *Toll-like Receptor Signaling* (*TLR*) pathway becomes *more significant* when the *Rig-I Like Receptor Signaling* (*RLR*) pathway (not significant on its own) is removed. The *TLR* pathway is the generic pathway involved in the immune response. The *RLR* pathway is the antiviral innate immunity pathway, which includes the mechanisms specifically aimed at the detection of exogenous DNA or RNA. In essence, the crosstalk analysis tells us that in the fat remodeling experiment, the immune system has been activated but this immune response is *not* due to the presence of foreign genetic material. This is exactly what happens here. The CL treatment causes the death of some white fat cells (Granneman et al., 2005). In turn, this causes an immune response in which macrophages are required to dispose of the dead cells (Li et al., 2001). Such subtle distinctions between various triggers that activated the immune response are not possible with any classical analysis methods, and it is remarkable that a data analysis method was able to provide this type of insight.

We then applied the proposed Maximum Impact Estimation described in the Methods section to the data. The corrected p-values are shown in Fig. 3b. The ranking based on these crosstalk corrected p-values is greatly improved. The most significant pathway is now the newly discovered mitochondrial pathway shown in Fig. 5 and validated by the in-situ hybridization shown in Fig 6. The new p-values also indicate the *Phagosome* pathway as one of the pathways related to this phenomenon (Newman et al., 1982). Third in the list is an independent module shared by *Cell Cycle* and *Oocyte Meiosis*. This can be thought of as a pathway related to the creation of new cells. Finally, the true involvement of *PPAR signaling* pathway in the phenomenon of fat remodeling has been previously demonstrated (Granneman et al., 2005) After removing the influence of the mitochondrial crosstalk, the *Parkinson's*, *Alzheimer's*, and *Huntington's* pathways are not significant anymore (now ranked 60th, 61st and 54th, respectively, data not shown). Also, after removing the crosstalk from *Phagosome*, *Leishmaniasis* is not significant anymore (now ranked 62nd, data not shown).

Cervical ripening. The second data set analyzed was obtained from a recent study that investigated the transcriptome of uterine cervical ripening in human pregnancy before the onset of labor at term (Hassan et al., 2009). The tissue analyzed is the human uterine cervix, the lower part of the uterus extending from the isthmus of the uterus into the vagina. It is mainly composed of smooth muscle and extracellular matrix, which consists of collagen, elastin, proteoglycans, and glycoproteins (Leppert, 1995; Ulbjerg et al., 1983b). The uterine cervix has an essential function in the maintenance of pregnancy and also in parturition (Hassan et al., 2006, 2009, 2010). Cervical ripening is a critical component of the common terminal pathway of parturition, which includes the extensive remodeling of the cervix (Hassan et al., 2009). Disorders of cervical ripening can lead to premature or protracted cervical change, complicating term (e.g. protracted dilatation or arrest of dilatation) or preterm gestations (e.g. premature cervical dilation in the second trimester) (Hassan et al., 2009). The state of cervical ripening has traditionally been assessed by clinical examination (Bishop score or its modifications (Bishop, 1964)), which includes the digital examination of the cervix for its consistency, dilatation, effacement, and position. This method has also been used to predict the likelihood that a patient would go into spontaneous labor. The goal of this experiment was to examine the relationship between human cervical ripening and the cervical transcriptome, aiming to improve our understanding of the biology of cervical ripening at term. This study in-

cluded pregnant women who underwent elective C-section at term with an unripe (n=11) or ripe cervix (n=11). Cervical biopsies were obtained from these women transvaginally, from the anterior lip of the uterine cervix following C-section. Microarray analysis was performed on RNA isolated from these cervical tissue specimens using Affymetrix GeneChip HGU133Plus2.0 arrays (Hassan et al., 2009).

On this dataset we performed the comparison between gene expression levels from cervical tissues obtained from women with an unripe (n=11) or ripe cervix (n=11) using the classical ORA. The results are shown in Fig. 7a. Pathways with a p-value smaller than 0.05 after FDR correction were *Focal adhesion*, *ECM-receptor interaction*, *Amoebiasis*, *Cell adhesion molecules (CAMs)*, *Small cell lung cancer*, and *Dilated cardiomyopathy*.

There is plenty of experimental evidence that biological processes described by the pathways *Focal Adhesion*, *ECM-Receptor Interaction*, and *Cell Adhesion Molecules* are related to cervical ripening. The relation between these pathways and the phenomenon in analysis was revealed by studies on humans and animals showing the involvement of extra-cellular matrix metabolism and cell adhesion molecules in cervical ripening (Leppert, 1995; Leppert et al., 1986; Mahendroo et al., 1999; Ulbjerg et al., 1983a,b). However, the pathway *Amoebiasis* describes the biological process of infection from a parasite that invades the intestinal epithelium. Amoeba infection involves the parasite attachment to the intestinal mucus layer, followed by disruption and death of host epithelial cells. This process is completely unrelated to the physiological condition of cervical ripening in term pregnancy. The same is true for the *Small Cell Lung Cancer* pathway. Clearly, the top ranked pathways include some describing complex phenomena that are unrelated to the studied condition. Also, the significant pathways known to be involved in the process of cervical ripening are somewhat general pathways describing cellular interactions.

The analysis of the crosstalk matrix (see Fig. S6 in Supplementary Materials) shows that there is a independent functional module among the top three pathways in the ranking. This novel module includes the genes present in the interaction between the cellular transmembrane protein integrin and three important ECM components, collagen, laminin, and fibronectin. The KEGG pathways involved in the identification of this pathway are *Focal adhesion*, *ECM-receptor interaction*, and *Amoebiasis*. Henceforth, we will refer to this pathway as the *Integrin-Mediated ECM Signaling* (Fig. 8).

Very interestingly, the independent functional module found in this condition is, in fact, the exact same module found in the hormone treatment experiment described in the Supplementary Materials. Interestingly, the KEGG pathways involved in the identification of this functional module are slightly different between the two phenotypes. While in this phenotype this module was found from the interaction of *Focal adhesion*, *ECM-receptor interaction* and *Amoebiasis*, in the hormone treatment the last pathway is replaced by *Pathways in Cancer*. The fact that the same module was found to be activated and statistically significant in two different phenotypes, from the interaction of different sets of canonical pathways, further supports the idea that this module describes an independent mechanism and should therefore be considered as an independent pathway.

Further analysis of the crosstalk matrix shows that the *Small Cell Lung Cancer* loses significance when the crosstalk effects of the first three pathways are removed (bright green loss of significance in first 3 columns of row 5 in Fig. S6). This allows us to conclude that it is a false positive in the classical ORA, with its ORA significance due exclusively to crosstalk effects.

The ranking of pathways with the p-values corrected for cross-talk by our analysis is shown in Fig. 7b. The first pathway is *Integrin-mediated ECM Signaling* with an FDR corrected p-value of $2.9e^{-13}$. *Cell Adhesion Molecules* is now the second in ranking, with an FDR corrected p-value of 0.004. The false positives in the classical ORA results, *Amoebiasis* and *Small Cell Lung Cancer*, are not significant anymore. The biological significance of the pathway *Dilated Cardiomyopathy*

may be linked to the fact that 10%-15% of the uterine cervix is constituted of smooth muscle, and cervical ripening involves alterations of this component. The last significant pathway at the 5% significance threshold is *Leukocyte Transendothelial Migration*. Although human and animal studies (Hassan et al., 2009) have shown that cervical ripening does not require activation of a typical inflammatory response, and influx of inflammatory cells into the cervix, the significance of this pathway may reflect the beginning of later inflammatory events typical of parturition (Timmons et al., 2009; Word et al., 2007).

In addition to the data sets described above, we analyzed a data set coming from the comparison of expression levels at days 7 and 0 in the fat remodeling experiment, a data set investigating the effect of various types of hormones on the endometrium of healthy post-menopausal women who underwent hysterectomy (Hanifi-Moghaddam et al., 2007), as well as a data set requested by one of the reviewers, investigating the correlation between gene expression values “*with MiniMental Status Examination (MMSE) and neurofibrillary tangle (NFT)*” in subjects with Alzheimer’s disease (Blalock et al., 2004). The latter dataset was analyzed to identify significant pathways from both KEGG and Reactome databases (see the Supplementary Results section in the Supplementary materials).

Conclusions

These results show that the novel approach proposed for the detection and correction for crosstalk effects allows not only the elimination of most of the false positives present in the results of the classical ORA, but also the identification of novel functional sub-pathways that are specifically involved in the condition studied, giving useful insights on the phenomenon in analysis that are not captured by existing techniques.

We assessed this novel approach on data from four real experiments coming from three phenotypes involving two species. In all cases, this approach was able to eliminate most false positives, as well as correctly identify as significant pathways that had been biologically proven to be involved in the given condition, yet not found to be significant by the classical analysis. We also found several independent functional modules including a *mitochondrial activity* module active in a different stage of fat remodeling in mice, and an *integrin-mediated ECM signaling* found to be involved in hormone treatment in post-menopausal women and cervical ripening in pregnant women. Interesting, the later module was extracted independently from the crosstalk interactions of two different groups of pathways, in the two conditions analyzed.

This approach is a departure from the current paradigm that considers the pathways as static models, independent of the phenotype. In the view proposed here, various specific modules, or sub-pathways, can be dynamically linked to specific conditions. When such independent functional modules are identified in independent conditions, such as the *integrin-mediated ECM signaling* above, these modules could be considered as candidate new pathways.

Methods

Pathway analysis in the presence of overlapping pathways: the crosstalk matrix. Our goal is to develop an approach that can detect and quantify the crosstalk between pathways. The main issue we are trying to address here is the fact that in the presence of overlapping pathways (i.e. for all pathways databases available today) crosstalk phenomena increase the probability of false positives, i.e. increase the number of pathways reported as significant but that in reality are not interesting (borrowing terminology from Brad Efron, we call pathways that have lesser biological

significance “not interesting” even though they might be statistically significant with a large enough sample size). To better understand the approach we are going to present, let us briefly review the classical Fisher Exact Test approach described above. Fig. 9a represents the contingency table used for assessing the significance of a pathway P_i by the classical over-representation (ORA) approach. The table divides genes as either being in the pathway or not, versus being considered DE or not DE (NDE); n_i represents the number of DE genes on P_i , while n represents the total number of DE genes, and m_i represents the number of NDE genes on P_i while m represents the total number of NDE genes. It follows that $n_i + m_i = |P_i|$ represents the number of genes on P_i , while with $n + m$ we represent the total number of genes.

The reasoning behind the ORA is that if the number of DE genes on a pathway is much higher than expected by chance, then the pathway is likely to be biologically interesting. In order to take into account the effect of the overlap on the significance of the two pathways we consider the effect of the removal of the overlapping part on the significance of the pathways. This is achieved as follows: let us consider two overlapping pathways P_i and P_j . With the notation $P_{i \setminus j}$ we define the set of elements in P_i excluding the intersection with P_j ; in the same way, with the notations $n_{i \setminus j} + m_{i \setminus j}$ we represent the number of genes that are in pathway P_i but not in pathway P_j , and with $n_{i \setminus j}$ the number of DE genes that are on pathway P_i but not in pathway P_j . We then consider the contingency table shown in Fig. 9b, whose bottom margin is identical to that of Fig. 9a.

With this contingency table, we compute for every pair of pathways $[i, j]$ the p-value of $P_{i \setminus j}$. Since this computation yields an $k \times k$ matrix, where k is the number of pathways, the results are most conveniently represented using a heat map of the negative log p-values. Each cell (i, j) of this matrix characterizes the significance of pathway P_i when we remove the effect of pathway P_j . The rows and the columns are ordered by the original p-values of the pathways, which are placed on the diagonal. We will refer to this matrix as the *crosstalk matrix*. This matrix is useful for identifying the effects of crosstalk among pathways.

An example of the crosstalk matrix can be found in Fig. 10. We will refer to the part of the matrix above the horizontal significance threshold as the *significance strip*. The *non-significance strip* will be the part below the horizontal significance threshold. The *significance quadrant* will be the part of the significance strip to the left of the significance threshold. Using these terms, we can identify and discuss several interesting phenomena that are not captured by any of the existing pathway analysis methods.

A first interesting case is when a pathway P_i is reported as significant by the classical analysis, but it *loses* its significance when the effect of another pathway P_j is removed. This is represented, in the crosstalk matrix, by a non significant p-value (green square) in the significance strip. In this case P_i is unlikely to be biologically meaningful, since its significance is most likely due to a crosstalk from P_j .

A second interesting case is when a pathway P_i that is *not* significant for the classical analysis *becomes* significant when the crosstalk effect of another pathway P_j is removed. This is represented in the crosstalk matrix by a significant p-value (red square) in the non-significant strip. The meaning of this is that pathway P_j was *masking* the significance of P_i , indicating that a phenomenon likely to be biologically meaningful is happening in the part of P_i which is not in common with P_j .

A third and last interesting case is a symmetric (with respect to the diagonal) decrease in significance of pathways in the significance quadrant. This indicates the presence of an independent functional sub-module, common to both P_i and P_j , that is responsible for their significance. Note that the activity of this module is tightly related to the condition studied.

The maximum impact estimation: an expectation maximization technique for the assessment of the significance of signaling pathways in presence of crosstalk. The crosstalk matrix is a useful tool for the interpretation of the effect of crosstalk between pathways. However, the ultimate goal of the analysis of signaling pathways is to provide a meaningful ranking among pathways, as well as a p-value quantifying the likelihood that a certain pathway is involved in the phenomenon in analysis. Here, we developed a correction method for the ranking of pathways that takes in account the overlaps between pathways.

The main idea is that, if there is no crosstalk, then there is no ambiguity in the ORA significance calculations. In such a case, if genes in a path are over-represented, it cannot be a false positive caused by crosstalk. Our approach is therefore to infer an underlying pathway impact matrix where each gene contributes to one and only one pathway, hence is devoid of crosstalk, and then to perform the ORA analysis using that impact matrix. Since this underlying pathway impact matrix is not observed directly, it is inferred through likelihood-based methods, and estimated using the EM algorithm. The corrected ranking is computed using ORA analysis with the underlying pathway impact matrix, shown as follows.

Let us consider the DE indicator vector Y , representing the differential expression of genes, and the membership matrix X describing the membership of each gene in each one of k pathways $P_1 \dots P_k$. The vector Y is defined as follows:

$$Y_i = \begin{cases} 1 & \text{if } g_i \text{ is DE} \\ 0 & \text{if } g_i \text{ NDE} \end{cases}$$

and each cell $X_{i,j}$ of the matrix X is defined as follows:

$$X_{ij} = \begin{cases} 1 & \text{if } g_i \text{ belongs to } P_j \\ 0 & \text{if } g_i \text{ does not belong to } P_j \end{cases}$$

The matrix $Y|X$ obtained by combining the vector Y with the X matrix is shown in the example in Fig. 11.

In many analysis methods, the membership matrix X is also interpreted as the *impact matrix*: if $X_{ij} = 1$, then gene g_i *impacts* pathway P_j . In ORA, for example, each gene is considered to have the same full impact on all pathways the gene belongs to. Crosstalk effects result from the fact that a gene can belong to more than one pathway, but in principle, it can potentially have a different biological impact on each such pathway. Our aim is to identify the pathway where the biological impact of such a shared gene is maximum. We do so by estimating the maximum impact pathway using an expectation maximization approach (see Supplementary materials).

Identification of independent functional modules. The maximum impact estimation procedure alone is not be able to identify overlapping modules responsible for the entire significance of other pathways, as in the situations represented by case 3 in the section describing the crosstalk matrix. In such cases the overlap should be considered as a separate pathway that is more likely to be biologically meaningful in the condition under analysis. An additional step is needed in order to correctly deal with this situation. In this additional step, we extract certain significant overlaps from the list of pathways, and include them in the list as *independent functional modules*. An independent functional module is a module for which there is evidence of an activity independent of the pathways it resides in, for the given condition. If an independent module is found in more than one, possibly unrelated, conditions, this module is considered as a *candidate novel pathway*.

A module must satisfy certain conditions in order to be treated as an independent functional module. Let us assume that we are analyzing the overlap between the pathways P_i and P_j ; the first

condition is that both pathways are significant (after FDR correction for multiple comparisons) at a certain threshold α . The threshold alpha is the significance threshold chosen by the user. Typical values for this threshold are 0.01 and 0.05. This condition limits the search to the significance quadrant of the crosstalk matrix. The second condition is that the overlap $P_i \cap_j$ itself must be significant at α (after FDR correction). The third condition is that the sub-pathways obtained by removing the overlap from both original pathways, indicated by $P_i \setminus_j$ and $P_j \setminus_i$, must *not* be significant at α (after FDR correction). If we denote with $p(P)$ the p-value of a generic pathway P , then the conditions can be summarized as follows:

1. $p(P_i) < \alpha, p(P_j) < \alpha$
2. $p(P_i \cap_j) < \alpha$
3. $p(P_i \setminus_j) \geq \alpha, p(P_j \setminus_i) \geq \alpha$

This pairwise procedure might yield modules that are similar one to each other, for example in cases where a module is contained in three or more pathways. That could be solved with a three-way or n-way search, but we opted for another approach for limiting the number of new modules. Once all interesting pairwise modules are created, we test for similarity among modules. The index used for similarity is a modified Jaccard Similarity index mJS defined as follows:

$$mJS = \frac{|M_1 \cap M_2|}{\min(|M_1|, |M_2|)} \quad (1)$$

where M_1 and M_2 are two modules obtained with the search criteria explained above. We merge any two modules similarity is greater than a certain threshold st . Once the modules are merged, the similarity among all the modules (including the newly created one) is computed again, and the merging procedure is applied again until there are no more modules that can be merged.

This newly created modules are removed from all pathways with which they overlap, and this list of modified pathways is used in the EM procedure. For the data sets analyzed in this work, we used an st threshold of 0.25. The procedure used to select this threshold can be found in the Supplementary Materials.

After applying the module discovery and the EM approach, the result is a modified membership matrix that can be used to perform the desired type of analysis. This matrix now includes three types of pathways: i) original pathways as found in the literature, ii) novel functional modules that are impacted in the given condition independently from the pathways they belong to, and iii) the pathways from which such independent modules have been removed. If the same independent module is found in several conditions, in other words if this module is active independently from its parent pathways in several different phenotypes, such a module should be considered a good candidate for a novel pathway.

Additional aspects of the crosstalk correction, such as a detailed description of the module detection step, false negative rate, and pathway size bias can be found in the Methods section of the Supplementary Materials (module detection step and false negative rate), and in Figs. S11 and S12 (pathway size bias).

Acknowledgements. This work has been partially supported by the following grants: NIH RO1 RDK089167, R42 GM087013, and NSF DBI-0965741 (to S.D.), by the Robert J. Sokol Endowment in Systems Biology, and by the Perinatology Research Branch, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NIH, DHHS. Any opinions, findings, and conclusions or recommendations expressed in this material are those of

the author(s) and do not necessarily reflect the views of the NSF, NIH, or any other of the funding agencies.

Author contribution. M.D. and S.D. identified the problem, conceived the study, and wrote the manuscript. M.D. and Z.X. developed the crosstalk correction based on the EM. S.D., M.D., P.H.W., and A.T. developed the crosstalk matrix analysis. J.G.G and R.G.M. performed the fat remodeling experiment, helped interpreting the results, and edited the fat remodeling experiment section in the manuscript. R.B. helped interpreting the results and edited the hormone treatment experiment section in the manuscript. N.G.T. helped interpreting the results and edited the cervical ripening section in the manuscript. P.H.W. edited the crosstalk matrix analysis and the crosstalk correction section in the manuscript. S.D., P.H.W, and R.R. supervised the study.

Disclosure declaration. The authors declare that they have no competing financial interests.

Figure legends

Figure 1: Pathway coupling in the Fisher Exact Test p-values. Left panel: a number of random genes were chosen from a “bait” pathway i such that its Fisher Exact Test p-value is 0.01. Other genes were chosen randomly from all other pathways (acting as preys), up to a constant number ($n = 100$). The elements $[i, j]$ where $i \neq j$ represent the mean of the distribution of p-values for 1000 random trials using pathway i as bait and pathway j as prey. The elements $[i, i]$ (on the diagonal) represent the classical Fisher Exact Test p-value of pathway i . The data show that a considerable number of pathways influence each other through a “coupling” of the p-values. For instance, row 3 of the matrix shows that when pathway 3 is chosen to be significant, several other pathways (e.g. columns 57 to 70) also tend to be significant (dark shades of blue represent significant p-values). Right panel: each point represents the average of the p-values of all the random trials for pairs with the same Jaccard index. The lines represent the fitting of linear and a quadratic models. Both models show a strong dependence between the p-value coupling and the Jaccard index. Similar results were obtained for GSEA and impact analysis (see Fig. 2).

Figure 2: Pathway coupling in the impact analysis (left panel) and GSEA (right panel). Each point represents the mean of all p-values of all random trials for pairs with the same Jaccard Index. The lines represent the fitting of linear and quadratic models. Both models show a strong dependence between the p-value coupling and the Jaccard Index.

- (a) The top 20 pathways resulting from classical ORA before correction for crosstalk. The top four pathways are not related to fat remodeling.
- (b) The top 20 pathways after correction for crosstalk. Pathways ranked 1, 3, and 5 are modules that are functioning independently of the rest of their pathways in this particular condition. Starred pathways are pathways edited by removing such modules. Note the lack of any obvious false positive above the significance threshold(s).

Figure 3: The results of the ORA analysis in the fat remodeling experiment for the comparison between days 3 and 0, before (left) and after (right) correction for crosstalk effects. All p-values are FDR-corrected. The lines show the significance thresholds: blue - 0.01, yellow - 0.05. Pathways highlighted in red represent pathways not related with the phenomenon in analysis, while pathways highlighted in green are those for which we know, with reasonable confidence, that they are involved in the given phenomenon. The white background indicates pathways for which we do not have conclusive information on their involvement (or lack of) with the phenomenon in analysis.

Figure 4: Detail of the crosstalk matrix: comparison between days 3 and 0 in the CL treatment. Areas marked with *a* correspond to functional modules that are activated independently from the pathways they belong to. The cell marked with *b* corresponds to a specific part of the *TLR* pathway that is responsible for the immune response to host genetic material. cells on the diagonal contain the p-values of the classical ORA, ordered from the most significant one to the least significant one. The cell $P_{i,j}$ contains the p-value of pathway P_i after the effect of P_j is removed. The color of each cell represents the p-value: bright red for p-values close to zero, bright green for p-values close to 1.

Figure 5: Mitochondrial activity pathway. This independent functional module is responsible for the incorrect identification of the pathways *Parkinson's disease*, *Alzheimer's disease*, *Huntington's disease*, and *Cardiac Muscle Contraction* by the classical ORA.

Figure 6: Epididymal white adipose tissue of a control mouse (left) and a mouse treated with CL for 7 days (right). Treatment with CL for 7 days triggered massive mitochondrial biogenesis, demonstrating *in vivo* that indeed, the mitochondrial pathway is central in this experiment. The white bar represents a 20 microns length.

- (a) Top 20 pathways reported by ORA before correction for crosstalk. Pathways like *Amoebiasis* and *Small Cell Lung Cancer* are not related to this phenotype.
- (b) The top 20 pathways reported by ORA after the crosstalk analysis. After the correction neither *Amoebiasis* nor *Small Cell Lung Cancer* are significant anymore. At the same time, *Cell Adhesion Molecules* and the *Integrin-mediated ECM Signaling* have an increased significance. Starred pathways are pathways edited by removing such module.

Figure 7: The results of the ORA for the cervical ripening experiment, before (left) and after (right) the correction for crosstalk effects. All p-values are FDR-corrected. The lines show the significance thresholds: blue - 0.01, yellow - 0.05.

Figure 8: The novel *Integrin-Mediated ECM Signaling*. This new module was found to be independently activated and statistically significant in two different conditions: hormone treatment of post-menopausal women and cervical ripening in normal pregnancies. Genes shown in red were found to be differentially expressed in the hormone treatment experiment.

(a) Standard over-representation approach contingency table; $n_i + m_i$ and $n + m$ represent, respectively, the number of genes belonging to pathway P_i and the total number of genes. n_i and n represent, respectively, the number of differentially expressed genes belonging to pathway P_i and the total number of DE genes.

(b) Contingency table for over-representation approach taking in account the overlap between pairs of pathways; $P_i \setminus j$ represents the set of elements in P_i excluding the intersection with P_j ; with the notations $n_{i \setminus j} + m_{i \setminus j}$ we represent the total number of genes that are in pathway P_i but not in pathway P_j , and with $n_{i \setminus j}$ the number of DE genes that are on pathway P_i but not in pathway P_j .

Figure 9: A comparison of the classical over-representation analysis (left) with the crosstalk matrix analysis proposed here (right).

Figure 10: Example of a crosstalk matrix. On the diagonal we find the classical over-representation analysis, ordered by p-value. The blue line represents the 0.01 significance level, while the black line represents the 0.05 significance level. The p-values in the matrix have been log-transformed (base 10 log) and the sign of the result has been inverted. The color of the cell represents the p-value: bright red for p-values close to zero, bright green for p-values close to 1.

Figure 11: Example of a DE/membership matrix; the column Y represents the indicator of differential expression of the various genes (1 for the n DE genes and 0 for the m NDE). Column P_j represents the membership indicator for pathway j . Row g_i describes gene i in terms of its differential expression and its membership to the various pathways.

Figures




Figure 1




Figure 2

rank	pathway	pval(fdr)	rank	pathway	p(fdr)
1	Parkinson's disease	2.0e-06	1	Mitochondrial Activity	8.1e-10
2	Alzheimer's disease	3.6e-06	2	Phagosome	9.3e-09
3	Huntington's disease	3.4e-05	3	Cellcycl+Oocyte	5.8e-08
4	Leishmaniasis	0.0003	4	PPAR signaling pathway	0.001
5	Phagosome	0.0006	5	Compl. C.C.+Systemic L.E.	0.002
6	Cell cycle	0.0011	6	* Cytok.-cytok. rec. int.	0.043
7	Oocyte meiosis	0.0016	7	Toll-like receptor signaling	0.051
8	Cardiac muscle contraction	0.0016	8	MAPK signaling pathway	0.115
9	Toll-like receptor	0.0018	9	B-cell receptor signaling	0.145
10	PPAR signaling pathway	0.0018	10	Lysosome	0.187
11	Chemokine signaling pathway	0.0154	11	Nat. killer cell med. cytotox.	0.187
12	Lysosome	0.0211	12	* Cell cycle	0.229
13	B cell receptor	0.0252	13	Calcium signaling pathway	0.229
14	Systemic lupus erythematosus	0.0292	14	Cell adhesion molecules	0.258
15	Compl. and coag. cascades	0.0342	15	NOD-like receptor signaling	0.258
16	Cytokine-cytokine rec. inter.	0.0346	16	Vasc. smooth muscle contr.	0.424
17	Chagas disease	0.0466	17	Dilated cardiomyopathy	0.424
18	Progest. med. oocyte matur.	0.0530	18	* Oocyte meiosis	0.432
19	Fc epsilon RI signaling pathway	0.0548	19	Type I diabetes mellitus	0.432
20	Leukocyte transendoth. migr.	0.0548	20	Wnt signaling pathway	0.476

(a)

(b)

Figure 3




Figure 4




Figure 5




Figure 6

rank	pathway	p(fdr)	rank	pathway	p(fdr)
1	Focal adhesion	$1.1e-08$	1	Integrin mediated ECM Signal.	$2.90e-13$
2	ECM-receptor interaction	$1.1e-08$	2	Cell adhesion molecules (CAMs)	0.0041
3	Amoebiasis	$1.2e-06$	3	Dilated cardiomyopathy	0.0041
4	Cell adhesion molecules	0.009	4	Leukocyte transendothelial migr.	0.0134
5	Small cell lung cancer	0.015	5	TGF-beta signaling pathway	0.2228
6	Dilated cardiomyopathy	0.015	6	Endocrine/other f.r. <i>Ca</i> reabs.	0.5791
7	Viral myocarditis	0.066	7	Insulin signaling pathway	0.9182
8	TGF-beta signaling path.	0.098	8	Alzheimer's disease	1
9	Prion diseases	0.1555	9	Vascular smooth muscle contr.	1
10	Leukocyte transend. migr.	0.1869	10	Glutamatergic synapse	1
11	Pathways in cancer	0.1869	11	Mineral absorption	1
12	Nat. killer c. med. cytotox.	0.2202	12	Nat. killer cell mediated cytotox.	1
13	Malaria	0.2202	13	Calcium signaling pathway	1
14	Adherens junction	0.3711	14	Complement and coag. casc.	1
15	Arr. right ventr. cardiom.	0.3711	15	MAPK signaling pathway	1
16	Calcium signaling pathway	0.3712	16	HTLV-I infection	1
17	Cholinergic synapse	0.6605	17	* * Focal adhesion	1
18	Vascular smooth muscle contr.	0.6605	18	* ECM-receptor interaction	1
19	Glutamatergic synapse	0.6969	19	* * Amoebiasis	1
20	HTLV-I infection	0.6969	20	Small cell lung cancer	1

(a)

(b)

Figure 7




Figure 8

	DE	NDE	Total
P_i	n_i	m_i	$n_i + m_i$
P_i^c	$n - n_i$	$m - m_i$	$(n + m) - (n_i + m_i)$
Total	n	m	$n + m$

(a)

	DE	NDE	Total
$P_{i \setminus j}$	$n_{i \setminus j}$	$m_{i \setminus j}$	$n_{i \setminus j} + m_{i \setminus j}$
$P_{i \setminus j}^c$	$n - n_{i \setminus j}$	$m - m_{i \setminus j}$	$(n + m) - (n_{i \setminus j} + m_{i \setminus j})$
Total	n	m	$n + m$

(b)

Figure 9:




Figure 10

	Y	P_1	P_2	P_3	\dots	P_k
g_1	1	0	1	1	\dots	0
g_2	1	0	1	0	\dots	0
g_3	1	1	0	0	\dots	1
\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots
g_{n-1}	1	0	0	1	\dots	0
g_n	1	0	1	0	\dots	0
g_{n+1}	0	0	0	1	\dots	0
g_{n+2}	0	1	0	1	\dots	0
\vdots	\vdots	\vdots	\vdots	\vdots	\ddots	\vdots
g_{n+m-1}	0	1	0	0	\dots	0
g_{n+m}	0	0	0	0	\dots	0

Figure 11

References

- Bishop, E. H., 1964. Pelvic Scoring For Elective Induction. *Obstetrics and gynecology*, **24**:266–268.
- Blalock, E. M., Geddes, J. W., Chen, K. C., Porter, N. M., Markesbery, W. R., and Landfield, P. W., 2004. Incipient Alzheimer’s disease: microarray correlation analyses reveal major transcriptional and tumor suppressor responses. *Proceedings of the National Academy of Sciences of the United States of America*, **101**(7):2173–2178.
- Draghici, S., Khatri, P., Martins, R. P., Ostermeier, G. C., and Krawetz, S. A., 2003. Global functional profiling of gene expression. *Genomics*, **81**(2):98–104.
- Draghici, S., Khatri, P., Tarca, A. L., Amin, K., Done, A., Voichita, C., Georgescu, C., and Romero, R., 2007. A systems biology approach for pathway level analysis. *Genome Research*, **17**(10):1537–1545.
- Goeman, J. J. and Bühlmann, P., 2007. Analyzing gene expression data in terms of gene sets: methodological issues. *Bioinformatics*, **23**(8):980–987.
- Granneman, J. G., Li, P., Zhu, Z., and Lu, Y., 2005. Metabolic and cellular plasticity in white adipose tissue I: effects of beta3-adrenergic receptor activation. *American Journal Of Physiology-Endocrinology And Metabolism*, **289**(4):E608–616.
- Hanifi-Moghaddam, P., Boers-Sijmons, B., Klaassens, A. H. A., van Wijk, F. H., den Bakker, M. A., Ott, M. C., Shipley, G. L., Verheul, H. A. M., Kloosterboer, H. J., Burger, C. W., *et al.*, 2007. Molecular analysis of human endometrium: short-term tibolone signaling differs significantly from estrogen and estrogen plus progestagen signaling. *Journal of Molecular Medicine-jmm*, **85**(5):471–480.
- Hassan, S. S., Romero, R., Haddad, R., Hendler, I., Khalek, N., Tromp, G., Diamond, M. P., Sorokin, Y., and Malone, J., 2006. The transcriptome of the uterine cervix before and after spontaneous term parturition. *Am J Obstet Gynecol*, **195**(3):778–786.
- Hassan, S. S., Romero, R., Tarca, A. L., Nhan-Chang, C.-L., Mittal, P., Vaisbuch, E., Gonzalez, J. M., Chaiworapongsa, T., Ali-Fehmi, R., Dong, Z., *et al.*, 2010. The molecular basis for sonographic cervical shortening at term: identification of differentially expressed genes and the epithelial-mesenchymal transition as a function of cervical length. *Am J Obstet Gynecol*, **203**(5):472.e1–472.e14.
- Hassan, S. S., Romero, R., Tarca, A. L., Nhan-Chang, C.-L., Vaisbuch, E., Erez, O., Mittal, P., Kusanovic, J. P., Mazaki-Tovi, S., Yeo, L., *et al.*, 2009. The transcriptome of cervical ripening in human pregnancy before the onset of labor at term: Identification of novel molecular functions involved in this process. *The Journal of Maternal-Fetal and Neonatal Medicine*, **22**(12):1183–1193.
- Jaccard, P., 1910. Étude anatomique de bois comprimés. *Bulletin de la Société Vaudoise des Sciences Naturelles*, **37**:547–579.
- Kanehisa, M., Goto, S., Kawashima, S., Okunom, Y., and Hattori, M., 2004. The KEGG resource for deciphering the genome. *Nucleic Acids Research*, **32**(Database issue):277–280.

- Lee, Y.-H., Petkova, A. P., Mottillo, E. P., and Granneman, J. G., 2012. In vivo identification of bipotential adipocyte progenitors recruited by Beta3-adrenoceptor activation and high-fat feeding. *Cell Metabolism*, **15**(4):480–491.
- Leppert, P. C., 1995. Anatomy and physiology of cervical ripening. *Clinical obstetrics and gynecology*, **38**(2):267–279.
- Leppert, P. C., Cerreta, J. M., and Mandl, I., 1986. Orientation of elastic fibers in the human cervix. *Am J Obstet Gynecol*, **155**(1):219–224.
- Li, M., Carpio, D. F., Zheng, Y., Bruzzo, P., Singh, V., Ouaz, F., Medzhitov, R. M., and Beg, A. A., 2001. An essential role of the NF-kappa B/Toll-Like Receptor pathway in induction of inflammatory and tissue-repair gene expression by necrotic cells. *The Journal of Immunology*, **166**(12):7128–7135.
- Li, P., Zhu, Z., Lu, Y., and Granneman, J. G., 2005. Metabolic and cellular plasticity in white adipose tissue II: role of peroxisome proliferator-activated receptor-alpha. *American Journal Of Physiology-Endocrinology And Metabolism*, **289**(4):E617–626.
- Mahendroo, M. S., Porter, A., Russell, D. W., and Word, R. A., 1999. The parturition defect in steroid 5alpha-reductase type 1 knockout mice is due to impaired cervical ripening. *Molecular endocrinology (Baltimore, Md.)*, **13**(6):981–992.
- Mootha, V. K., Lindgren, C. M., Eriksson, K.-F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., *et al.*, 2003. PGC-1 alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nature Genetics*, **34**(3):267–273.
- Mottillo, E. P., Shen, X. J., and Granneman, J. G., 2007. Role of hormone-sensitive lipase in beta-adrenergic remodeling of white adipose tissue. *Am J Physiol Endocrinol Metab*, **293**(5):E1188–97.
- Nam, D. and Kim, S.-Y., 2008. Gene-set approach for expression pattern analysis. *Briefings in bioinformatics*, **9**(3):189–197.
- Newman, S. L., Henson, J. E., and Henson, P. M., 1982. Phagocytosis of senescent neutrophils by human monocyte-derived macrophages and rabbit inflammatory macrophages. *The Journal of Experimental Medicine*, **156**(2):430.
- Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., Paulovich, A., Pomeroy, S. L., Golub, T. R., Lander, E. S., *et al.*, 2005. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proceeding of The National Academy of Sciences of the USA*, **102**(43):15545–15550.
- Tarca, A. L., Draghici, S., Khatri, P., Hassan, S. S., Mittal, P., sun Kim, J., Kim, C. J., Kusanovic, J. P., and Romero, R., 2009. A novel signaling pathway impact analysis (SPIA). *Bioinformatics*, **25**(1):75–82.
- Tavazoie, S., Hughes, J. D., Campbell, M. J., Cho, R. J., and Church, G. M., 1999. Systematic determination of genetic network architecture. *Nature Genetics*, **22**:281–285.
- Timmons, B. C., Fairhurst, A.-M., and Mahendroo, M. S., 2009. Temporal changes in myeloid cells in the cervix during pregnancy and parturition. *J Immunol*, **182**(5):2700–2707.

- Uldbjerg, N., Ekman, G., Malmström, A., Olsson, K., and Ulmsten, U., 1983a. Ripening of the human uterine cervix related to changes in collagen, glycosaminoglycans, and collagenolytic activity. *American Journal of Obstetric and Gynecology*, **147**(6):662–666.
- Uldbjerg, N., Ulmsten, U., and Ekman, G., 1983b. The ripening of the human uterine cervix in terms of connective tissue biochemistry. *Clinical obstetrics and gynecology*, **26**(1):14–26.
- Word, R. A., Li, X.-H., Hnat, M., and Carrick, K., 2007. Dynamics of cervical remodeling during pregnancy and parturition: mechanisms and current concepts. *Seminars in reproductive medicine*, **25**(1):69–79.



Analysis and correction of crosstalk effects in pathway analysis

Michele Donato, Zhonghui Xu, Alin Tomoiaga, et al.

Genome Res. published online August 9, 2013

Access the most recent version at doi:[10.1101/gr.153551.112](https://doi.org/10.1101/gr.153551.112)

Supplemental Material <http://genome.cshlp.org/content/suppl/2013/09/17/gr.153551.112.DC1>

P<P Published online August 9, 2013 in advance of the print journal.

Accepted Manuscript Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.

Creative Commons License This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 3.0 Unported), as described at <http://creativecommons.org/licenses/by-nc/3.0/>.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>
