



Benchmarking bulk and single-cell variant calling approaches on Chromium scRNA-seq and scATAC-seq libraries

Matthew V J Wiens, Hossein Farahani, R Wilder Scott, et al.

Genome Res. published online August 15, 2024

Access the most recent version at doi:[10.1101/gr.277066.122](https://doi.org/10.1101/gr.277066.122)

P<P Published online August 15, 2024 in advance of the print journal.

Accepted Manuscript Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.

Creative Commons License This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <https://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Comprehensive immune receptor profiling.
Discover the **DriverMap™ AIR Assay** difference.

LEARN
MORE



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Published by Cold Spring Harbor Laboratory Press

Benchmarking bulk and single-cell variant calling approaches on Chromium scRNA-seq and scATAC-seq libraries

Matthew Wiens¹, Hossein Farahani¹, R. Wilder Scott¹, T. Michael Underhill^{1,2 *}, Ali Bashashati^{1,3 *}

1: School of Biomedical Engineering, University of British Columbia

2: Department of Cellular & Physiological Sciences, University of British Columbia

3: Department of Pathology & Laboratory Medicine, University of British Columbia

* These authors supervised the study

Corresponding Author Contact:

ali.bashashati@ubc.ca

Running Title:

Single-cell variant calling on Chromium libraries

Abstract

Single-cell sequencing methodologies such as scRNA-seq and scATAC-seq have become widespread and effective tools to interrogate tissue composition. Increasingly, variant callers are being applied to these methodologies to resolve the genetic heterogeneity of a sample, especially in the case of detecting the clonal architecture of a tumor. Typically, traditional bulk DNA variant callers are applied to the pooled reads of a single-cell library to detect candidate mutations. Recently, multiple studies have applied such callers on reads from individual cells, with some citing the ability to detect rare variants with higher sensitivity. Many studies apply these two approaches to the Chromium (10x Genomics) scRNA-seq and scATAC-seq methodologies. However, Chromium-based libraries may offer additional challenges to variant calling compared to existing single-cell methodologies, raising questions for the validity of variants obtained from such a workflow. To determine the merits and challenges of various variant-calling approaches on Chromium scRNA-seq and scATAC-seq libraries, we use sample libraries with matched bulk whole-genome-sequencing to evaluate the performance of callers. We review caller performance, finding that bulk callers applied on pooled reads significantly outperform individual-cell approaches. We also evaluate variants unique to scRNA-seq and scATAC-seq methodologies, finding patterns of noise but also potential capture of RNA-editing events. Finally, we review the notion that variant calling at the single-cell level can detect rare somatic variants, providing empirical results that suggest resolving such variants is infeasible in single-cell Chromium libraries.

1 Introduction

2 Single-cell sequencing technologies have emerged as one of the most powerful interrogative tools
3 currently available for studying cell populations in a variety of biological scenarios. Single-cell DNA-
4 sequencing (scDNA-seq) tools revealed new insights into the ancestral hierarchy and sub-clonal
5 composition of various cancers (Navin et al. 2011; Gawad et al. 2016). Additionally, other single-cell
6 approaches, such as single-cell RNA-sequencing (scRNA-seq) and single-cell ATAC-sequencing (scATAC-
7 seq), have resulted in major advances in our understanding of intercellular heterogeneity, cellular
8 differentiation, and cell-cell interactions (Jaitin et al. 2014; Buenrostro et al. 2015). These single-cell
9 sequencing technologies are increasingly being paired with short variant calling approaches to
10 interrogate the unique haplotype state of individual cells (Zhou et al. 2020; Lu et al. 2021; Fan et al.
11 2018; Ding et al. 2019; Petti et al. 2019; Vu et al. 2019; Poirion et al. 2018). This pairing enables
12 investigators to extract additional information from these libraries regarding the single-cell genetic
13 state, thus saving the time and money otherwise required to obtain scDNA-seq libraries. For instance,
14 using scRNA-seq, the clonal composition of a tumor sample and expression phenotypes of individual
15 subclones have been determined independently of scDNA-seq (Zhou et al. 2020; Lu et al. 2021; Ding et
16 al. 2019). An additional benefit is that modality specific sequence-altering effects such as RNA editing
17 can be investigated at the single-cell level (Picardi et al. 2017).

18 The most popular single-cell methodology is the Chromium platform provided by 10x Genomics which
19 employs a droplet-encapsulation-based approach to sequence many thousands of cells in parallel
20 (Zheng et al. 2017). Single-cell alternative sequencing modalities, especially libraries generated by
21 Chromium pipelines, pose new challenges for existing variant callers due to unique biological and
22 technical limitations. To start, cell loadings in Chromium libraries range anywhere from 500 to over
23 50,000 cells, and at these high loadings most cells have low coverage so that they may be sequenced in

1 a single run. As a result, alternative allele visibility can be poor at the single-cell level, where cells will
2 often only have a single read at a given site. Additionally, the sparse coverage provided by approaches
3 such as scATAC-seq and scRNA-seq limits the regions in which variants can be detected. ATAC-seq is
4 sparse due to its targeted sampling approach, while Chromium scRNA-seq only sequences one end of a
5 mRNA transcript, significantly reducing overall gene coverage. Modality-specific effects, such as allelic
6 dropout, can further impact the accuracy of the allele called in each individual cell. Finally, at higher cell
7 loadings, beads may contain more than one cell, which can complicate the analysis. Very few variant
8 callers have been designed to address these unique challenges.

9 Nonetheless, multiple studies have performed variant calling on libraries generated by the Chromium
10 3'/5' scRNA-seq and scATAC-seq chemistries (Lu et al. 2021; Fan et al. 2018; Massarat et al. 2021; N. M.
11 et al. 2021; Petti et al. 2019; Dou et al. 2023). A wide variety of variant calling approaches have been
12 applied. These fall into one of two categories: treating the reads as having been called in bulk (i.e.,
13 applied to the pooled reads) and disregarding cell-level information; or leveraging cell level information
14 and inter-cell statistics to improve variant calling. The latter approach is of interest due to its potential
15 to discover rare variants visible only at the level of small groups of cells. Indeed, studies have claimed
16 that cell-level variant calling reveals a wealth of variants not visible in whole genome and exome
17 sequencing (N. M. et al. 2021). However, the extent to which such observed variants occur due to
18 biological variations versus false positives resulting from artifactual sources requires further
19 investigation. Additionally, although one recent study has performed benchmarking of pooled callers as
20 part of a comparison with their purpose-built single-cell variant caller (Dou et al. 2023), there are no
21 publications that focus on comparing the performance of pooled and individual-cell variant calling
22 approaches which have been used in studies with Chromium-derived single-cell sequencing libraries
23 (Dou et al. 2023).

In this study, we aim to provide a comprehensive comparison of commonly used single-cell and pooled variant calling approaches on matched whole genome sequencing (WGS) and 10x Chromium scRNA-seq and scATAC-seq libraries derived from mouse tumor samples. We additionally aim to investigate rare variants which might be detected only in single-cell modalities and only at the individual-cell level to determine to what extent real rare variants may be extracted from these modalities.

Results

Benchmark Datasets

Three tumor samples derived from a genetically engineered mouse model of epithelioid sarcoma featuring an induced adulthood deletion of *Smarb1* were acquired and each sample underwent Chromium scRNA-seq and scATAC-seq, as well as Illumina WGS, library generation (see Supplemental Table S1). Additionally, a control sample underwent Chromium scRNA-seq and scATAC-seq. Single cell sequencing was performed at loadings of around 7000-15000 cells per sample. This loading is typical for Chromium pipelines and serves to reproduce the low per-cell coverage typical of this single-cell methodology.

Benchmark Metrics

Our unique experimental setting allowed us to capture both real variants that would be present in many cells as well as rare variants that would be present in small fraction of cells. To identify real variants, the consensus of multiple variant callers (HaplotypeCaller (Poplin et al. 2018), strelka2 (Kim et al. 2018), SAMtools (Li 2011) and VarScan (Koboldt et al. 2009)) on the matched WGS libraries was utilized (referred to as *common ground truth variants*; Table 1). Given that WGS is limited in its capability to detect variants present in a small number of cells, as compared to other modalities, we would expect that such rare variants would most often be undetected or indistinguishable from noise in WGS data. As

such, variant calls identified in scRNA-seq or scATAC-seq data without significant evidence in WGS were treated as potential rare variants (referred to as *rare positive variants*). We define a number of variant categories and metrics (Table 1; Supplemental Figure S1) to facilitate comparison of various variant calling algorithms. These are described in more detail in Methods (see section “Variant Categories and Metrics”).

Variant Categories	Other Names	Definition
common ground truth	WGS variants	Variants derived from whole genome sequencing
pooled read		Variants derived from pooled-read bulk calling on single-cell sequencing
individual-cell variants		Variants derived from individual-cell calling on single-cell sequencing
pooled true positive	TP	Pooled-read variant which has a matching WGS variant
pooled false negative	FN	WGS variant not detected in pooled-read variant calling on single-cell sequencing
rare positive	RP	variant that was detected in single-cell sequencing which was not present in the common ground truth set
union of rare positive (URP)	scRNA-RP / scATAC-RP	Union of rare positive variant calls for all callers performed on pooled scRNA-seq or scATAC-seq reads
multimodal rare positive	MRP	URP variants seen in both scATAC-seq and scRNA-seq modalities
single cell unique (scu)	scRNA-scu / scATAC-scu	variant detected at the single-cell level in single-cell sequencing which was not called on pooled reads nor found in WGS

Table 1: Quick reference for definitions of variant categories defined and used frequently

throughout this study

As our primary metric to compare various variant-calling approaches on single-cell sequencing, we use a precision-recall measure that evaluates missed variants (pooled false negatives) and extra variants that have been called (rare positives) for each caller. We used the quality value for each variant generated by each caller as the score threshold to generate precision and recall across different levels. We further computed the area under the precision-recall curve (AUPRC) to compare the overall performance of each caller. Precision and recall were computed for variants across multiple samples to provide an overall score. We additionally use metrics termed the common true positive rate (cTPR) and common false discovery rate (cFDR) to evaluate the performance of variant callers with different filtering

approaches. Refer to Methods (section “Variant Categories and Metrics”) for more details on these metrics.

Benchmarking Variant Callers

We consider variant calling approaches for single-cell datasets as falling into two categories: pooled and individual-cell methods. Bulk callers have been tested previously on Chromium scRNA-seq and scATAC-seq libraries at the pooled read level (Liu et al. 2019; Massarat et al. 2021). However, no studies have tested the application of bulk callers to individual cells for Chromium libraries. Given that such approaches are increasingly being used (Lu et al. 2021; Ding et al. 2019), it is essential to benchmark their performance against bulk methods and confirm their utility.

Typically, for individual-cell calling, the intersection of variants called across multiple cells is used to derive a consensus list that can be filtered by quality. To investigate the potential of these approaches, we apply Strelka2 (Kim et al. 2018) and GATK HaplotypeCaller (Poplin et al. 2018) on individual BAM files for reads corresponding to a single cell. We then take the intersection of variants from individual cells and define the quality of each variant as the maximum quality value of the variant over the cells in which it was found.

Based on previous benchmarking studies applying bulk callers to Chromium single-cell pooled reads (Liu et al. 2019; Massarat et al. 2021), we test Strelka2, SAMtools (Danecek et al. 2021), GATK HaplotypeCaller, and FreeBayes (Garrison and Marth 2012) as the most promising options, given their previously tested performance. We run them on the pooled reads for each single-cell library according to their recommended best-practices approach for each modality.

It is notable that there are many purpose-built options for individual-cell variant calling, such as Monovar (Zafar et al. 2016), SCcaller (Dong et al. 2017), SciPhi (Singer et al. 2018), ProSolo (Lähnemann

et al. 2021), ScanSNV (Luquette et al. 2019), scAllele (Quinones-Valdez et al. 2022), and Monopogen (Dou et al. 2023). Many of these callers were developed to address allelic bias due to multiple-displacement amplification used in scDNA-seq methodologies. As a result of their tailored statistical modeling approaches and traditional application to high-coverage single cells, they have been dismissed as inapplicable to Chromium single-cell methodologies (Lu et al. 2021). However, the ability of these tools to adaptively model changing allele frequency might be helpful to correct for allelic imbalance due to preferential visibility of one chromosome in scRNA-seq and scATAC-seq, thus improving variant detection. The major question is whether they will transfer to the low read coverage per cell of Chromium-based sequencing. To verify this viability, we benchmark SCcaller, which among the caller options has a varying window allelic modeling approach suitable for RNA sequencing.

It should be mentioned that there are recently developed callers developed specifically for improving sensitivity in single cell modalities through the use of additional population-level information to overcome the limitations of single-cell sequencing. Examples include Monopogen (Dou et al. 2023) which leverages population level statistics such as linkage disequilibrium to help differentiate real variants from noise, and scAllele (Quinones-Valdez et al. 2022) which leverages splicing patterns to help inform variant confidence. While we do not benchmark them in this paper, approaches like these may have significant promise for Chromium-based single-cell modalities.

Given that single-cell data has the potential to capture variants which show evidence in a low number of cells and thus have a low presentation frequency, we first investigated the performance of variant callers at different *variant allele frequencies* (VAF, the fraction of alternate-allele reads over total reads at a variant site). We can see from Figures 1c and 1d that, for both scATAC-seq and scRNA-seq, the majority of variants either have very low VAF (less than 0.01) or very high VAF (equal to 1.0). As such, a failure to detect low-VAF variants will miss many WGS variant sites being lost. All callers suffer a drop in

sensitivity at low VAF, especially below VAF of 0.01 (Figures 1a and 1b). However, two of the individual-cell callers, SCcaller and Strelka, perform quite well at low VAF compared to all other callers, possibly highlighting the potential of individual-cell callers to pick up lower-frequency variants. Based on the results of Figure 1, in our further analyses we kept only those variants with VAF of 0.01 or greater to filter out the large number of very-low-frequency variants missed by all callers.




Figure 1: Common True Positive Rate averaged over samples for different variant-calling approaches, for scRNA-seq (a) and scATAC-seq (b), binned by VAF (e.g. the point VAF=0.5 would include variants from 0.5 to the next bin, which is 0.6); The number of WGS variants with coverage in the single-cell sequencing modality, for scRNA-seq (a) and scATAC-seq (b) binned by VAF

Figures 2a and 2b demonstrate the performance of variant calling approaches on scRNA-seq and scATAC-seq respectively based on precision-recall curves. We can see that pooled read calling can

achieve good sensitivity in both cases, reaching up to 90.5% and 76.9% recall in each modality respectively. Based on AUPRC, we find that Strelka demonstrates the best performance on scRNA-seq, while SAMtools demonstrates best performance on scATAC-seq (see Table 2). Notably, in scRNA-seq, most callers have lower precision than in scATAC-seq, indicating that extra variants are consistently called in scRNA-seq that are not present in the common ground truth set. This is especially notable when variants are called in non-exonic and exonic regions (see supplemental Figure S2) and raises the importance of limiting scRNA-seq variant calling to exonic regions. In later sections, we investigate what proportion of these rare positive variants are noise/artifacts, biological factors unique to sampling RNA, or true biological variants. The fact that callers perform well at detecting common ground truth variants indicates that the WGS variants are well-represented at the pooled read level in both modalities.

Contrary to bulk callers applied to the pooled reads, the same callers applied at the individual cell level do not effectively resolve common ground-truth variants. Both Strelka SC and HaplotypeCaller SC demonstrate low recall at most filtering thresholds, only detecting common ground truth variants when quality filters are so stringent that only a handful of variants remain (corresponding to the left side of each curve). On scATAC-seq datasets, the single-cell callers do manage to match pooled read calling for precision at the very extremum of their quality threshold, indicating that variant calling at the single-cell level is able to resolve common ground-truth variants, but at the cost of extremely low recall.

HaplotypeCaller, on the other hand, performs poorly at any filtering threshold, indicating a very low sensitivity to common ground-truth variants.

	Strelka	HaplotypeCaller	SAMtools	FreeBayes
scRNA-seq	0.655	0.575	0.743	0.295
scATAC-seq	0.612	0.482	0.546	0.416

Table 2: Area under precision recall curve (AUPRC) for bulk callers on pooled reads, in exonic regions for the case of scRNA-seq




Figure 2: Precision-Recall curves averaged over samples for different variant-calling approaches, for scRNA-seq (a) and scATAC-seq (b); Individual cell calling (Strelka SC, HaplotypeCaller SC) shown with dashed line; dots signify the recommended quality filter level for each caller (QUAL > 30 for HaplotypeCaller, QUAL > 20 for others, no dot for single-cell callers which do not have a recommended filter level)

A common expectation of variant callers applied at the individual cell level is the ability to resolve rare variants which have low alternate allele evidence. To determine the ability of variant callers to detect low-read-support variants, we look at the variants called on the single-cell libraries without applying any filtering to the variant sites determined by each caller. As we can see in Figure 3a and 3b, certain bulk callers considerably outperform others, with Strelka consistently demonstrating the highest sensitivity for both scRNA-seq and scATAC-seq. Additionally, the individual cell application of Strelka (Strelka SC) appears to match the performance of Strelka when called in bulk. On the other hand, HaplotypeCaller calling on individual cells (HaplotypeCaller SC) demonstrates poor performance at detecting common

- 1 ground-truth variants across all read coverage levels, indicating that the caller has decreased sensitivity
- 2 when applied at the single-cell level.




Figure 3: Average over samples of common true positive rate (cTPR) (**a, b**) and common false discovery rate (cFDR) (**c, d**) versus alternate allele coverage for variants binned by total (e.g. pooled) alt-allele coverage (e.g. the point coverage=400 would include variants from 400 to the next bin, which is 500), for each of the variant-calling approaches used, for both scRNA-seq (**a, c**) and scATAC-seq (**b, d**)

1 Individual-cell application of bulk callers demonstrate extremely high numbers of rare positive variants,
2 as is evidenced by the cFDR statistics in Figure 3c and 3d. HaplotypeCaller SC demonstrates lower FDR
3 than Strelka SC, possibly due to its decreased sensitivity. Pooled read callers (with the exception of
4 FreeBayes) demonstrate similar performance in cFDR across allele coverage levels, indicating more rare
5 positives for variants with low alternate allele coverage.

6 By comparison to other calling approaches, SCcaller significantly underperforms, mostly due to a high
7 cFDR, especially in scRNA-seq. While the cTPR of the caller is decent, being outperformed only by
8 Strelka, it appears that SCcaller is indiscriminate in its sensitivity. Given that SCcaller has been shown to
9 obtain good results on MDA scDNA-seq datasets, this may be primarily due to the low coverage present
10 in the cells. Additionally, Strelka SC also generates a very high cFDR across different allele coverages in
11 the case of scRNA-seq. Overall, this result supports the suspicion of others that existing single-cell
12 variant callers are not suitable for Chromium data, although this may not be the case for all such callers.

13 Overall, these results agree with previous studies demonstrating that a wide range of variants present in
14 a sample are visible in libraries generated by Chromium scRNA-seq and scATAC-seq modalities (Liu et al.
15 2019; Massarat et al. 2021). This enables such modalities to supplement, or act independent of, DNA-
16 sequencing to discover biologically relevant variant sites. The choice of caller has a significant impact on
17 the ability to resolve these variants. Applying bulk callers to individual cells results in extremely high rare
18 positive rates likely owing to a large number of false-positive calls, making such approaches impractical
19 for calling biologically relevant variants. The potential veracity of such rare positives is discussed in later
20 sections. Applying bulk callers to pooled reads can provide quite useful results, even at standard quality
21 levels (see dots on Figure 2). Strelka, HaplotypeCaller, and SAMtools perform well on both scRNA-seq
22 and scATAC-seq, with small differences such that it is difficult to say which one is best. Depending on
23 whether precision or recall is more important, and depending on which modality is used, Strelka or

SAMtools might give optimal performance. There appears to be trade-offs between callers, and we hope the data we have presented here will help inform researchers in the field as to which approach they would prefer.




Figure 4: Comparing the composition of URP and TP variants by VAF and read depth for scRNA-seq (a, b) and scATAC-seq (c, d), number of variants with read depth over 100 are summed in the right-most bin

Investigating Rare Positive Variants in Pooled Reads

Previously we noted that bulk callers applied to pooled reads detect variants in addition to those found in the common ground-truth set. It is possible that some of these rare positive variants are either real and not detected in WGS or due to other biological factors. ATAC-seq and RNA-seq modalities frequently sample specific regions to a depth much higher than the 40x coverage we obtained for our WGS. This deeper coverage may enable higher alternate allele visibility and greater confidence in variant calling at a given locus. However, rare positive variants may just as well be the result of artifacts due to noise/error or modality-specific sequence alterations. Considering that explicit verification of these variants would likely require high-depth sequencing to confirm, we use indirect approaches to draw conclusions about the portion of rare positive variants that are potentially real.

To examine the composition of rare positive variants in general, we defined *union rare positive* (URP) variants as the union of rare positive variant calls for all callers performed on pooled scRNA-seq or scATAC-seq reads. We quantified the number of URP variants without any filtering except which demonstrate evidence (at least one alt-allele read) in WGS (Table 3). This is to account for any variants that did not pass filters for calling in our common ground-truth set (i.e., WGS), but which gave sufficient evidence in scRNA-seq or scATAC-seq to be called. We find that, on average, 9.8% of URP variants called in ATAC-seq (scATAC-RPs) are present in at least 5 reads in WGS, with 6.8% present in at least 10 reads. On the other hand, only 2.2% and 1.2% of scRNA-seq URP variants (scRNA-RPs) show evidence of being present in WGS at depths greater than 5 or 10 reads respectively. Thus, it appears that a portion of scATAC-RPs may be real and possibly missed in the common ground truth set, while this is significantly less so for scRNA-RP variants. This raises a concern that scRNA-RPs are enriched for noisy or artifactual variants.

1 It should be noted that scRNA-seq appears to generate more URPs than scATAC-seq. If we normalize by
 2 the total coverage of each modality (see Table 3), we find that scRNA-seq has 2.9x more URP variants
 3 per megabase than scATAC-seq ($p=0.087$ by 2-tail paired t -test, $n=3$), despite scATAC-seq detecting
 4 similar or more ($\sim 1.8x$) common ground-truth variants than scRNA-seq ($p=0.068$ by 2-tail paired t -test,
 5 $n=3$). The extra variants we see may either be real and due to differences in coverage location, or due to
 6 higher error rates in scRNA-seq.

	scRNA-seq			scATAC-seq		
	EPSRC1	EPSRC2	EPSRC4	EPSRC1	EPSRC2	EPSRC4
Coverage, Mb	466.8	385.7	435.8	2373.2	2015.6	2435.4
WGS Variants, count (/Mb)	120428 (258.4)	122854 (318.5)	84070 (192.9)	1111673 (468.4)	846406 (419.9)	1186267 (487.1)
URP Variants, count (/Mb)	120815 (258.8)	101475 (263.1)	200310 (459.6)	330710 (139.3)	169171 (83.9)	271228 (111.4)
scu Variants, count (/Mb)	661 (1.416)	761 (1.973)	2079 (4.771)	25 (0.011)	9 (0.004)	48 (0.020)

Table 3: Counts of Variants Found in Each Modality

8 To determine if rare positive variants tend to have a lower read coverage or lower VAF distribution,
 9 potentially indicating noise as a cause for these variants, we broke down scRNA-RP and scATAC-RP
 10 variants by VAF (Figures 4a and 4c) and read depth (Figures 4b and 4d) in comparison with pooled TP
 11 variants for control. Notably, while URP variants in both modalities appear to be slightly more distributed
 12 towards lower VAFs as compared to TP variants, there appears to be no meaningful difference that
 13 would indicate VAF as a major correlating factor for rare positive variants. Likewise, the read depth
 14 distribution of URP for both modalities is relatively similar to that of TP variants, with TP variants
 15 presenting in a higher portion of low-read-coverage sites than URP variants, especially for scATAC-seq.
 16 The tendency of URPs to occur in regions of higher coverage for scATAC-seq may be indicative of a
 17 systematic source for their generation, which we attempt to explore more in the next section.




Figure 5: SBS 96 trinucleotide mutation breakdown for (a) common ground-truth, scATAC-RP, and scRNA-RP variants as seen in EPSRC1 which is representative of the group trend, and (b) the SBS1536 breakdown of scRNA-RP variants in exonic regions only, demonstrating that the signature is maintained in exonic regions and is characterized by repeat-extensions of all base patterns. Images were generated using the SigProfiler suite.

1 scATAC-seq and scRNA-RP variants, as well as common ground-truth variants. The COSMIC (Alexandrov
2 et al. 2020) signatures serve as a reference of well-known mutational signatures that might compose
3 some part of the variants seen in our samples.

4 We find that scATAC-RP variants feature a trinucleotide mutational breakdown similar to that seen in
5 WGS (Figure 5a), namely with a high proportion of variants which may be explained as a composition of
6 the COSMIC signature SBS5 (Figure 6a). scATAC-RPs in EPSRC1 also appear to match the SBS45, and
7 SBS46 signatures, which have been linked to sequencing errors. scATAC-seq also features a prominent
8 CTG->CCG rate (Figure 5a), which does not match any known signatures that we have reviewed and may
9 be due to a preparation/sequencing error. As such, it appears that, while scATAC-RPs may contain real
10 variants, there is a substantial risk of including noise based on the signatures observed.

11 On the other hand, the trinucleotide mutational breakdown of scRNA-RPs variants differs substantially
12 from that of WGS, demonstrating extremely high incidence of repeat extension mutations of the form
13 YY[X]YY>YY[Y]YY, especially in poly(T)/poly(C) regions (Figure 5a, b). These mutations are present even if
14 we reduce the list of variants to those found in exonic regions, which we would expect to have higher
15 quality due to increased coverage by scRNA-seq (Figure 5b). According to SigProfiler, the signature of
16 these mutations does not match any yet established signature, although they are somewhat similar to
17 SBS58, another COSMIC potential sequencing artifact signature (Figure 6a, 6b).

18 These previous results suggest that many URP variants are due to noise. One major advantage of our
19 dataset is the matched scRNA-seq and scATAC-seq, which allows rare variants to be validated by
20 confirming presence in both modalities. We use the intersection of scRNA-RP and scATAC-RP variants as
21 a validation set for rare positive variants, which we term multimodal rare positive (MRP) variants. We
22 use estimated MRP and URP variant statistics (Table 4, Definition in Methods - Variant Categories and
23 Metrics, which have minimum threshold of 5 supporting reads) alongside pileup overlap statistics to

extrapolate the number of valid URP variants we would expect to see in each modality. We find that the rate of MRP variants per megabase is extremely small. If we extrapolate this rate to the URP variants unique to each modality, we find that they would make up around 2% and 14% of unique URP variants in scRNA-seq and scATAC-seq, respectively. This agrees well with the portion of rare positive variants seen with evidence of the alternate allele in WGS discussed previously. It should be noted that the true number of overlapping rare variants may be larger due to regions where one modality has very low coverage and thus does not sample the rare alternate allele. Overall, these results suggest that rare variants detectable through pooled-read variant calling in scRNA-seq and scATAC-seq make up only a small portion of rare positive variants.

	scRNA-seq	scATAC-seq	Overlapping
Total Coverage, MB	177.4	1260.0	128.0
URP/MRP Variants, count	33180	35317	1012
MRP Variant Rate, /MB			8.1 ± 0.8
Predicted Rare Variants, using MRP Variant Rate	728.7 (0.021)	4926.5 (0.138)	

Table 4: scRNA-seq and scATAC-seq Overlapping URP Variants

Overall, it appears that many of the URP variants may be explained as due to noise. Some of the URP variants show evidence of being real rare mutations indicating that scATAC-seq and scRNA-seq de-novo calling may yield variants not visible in typical WGS experiments. However, they make up such a small portion of the URP variant set that they would be infeasible to separate out from the majority of noisy variants in any practical scenario. The increased incidence of scRNA-RPs compared with scATAC-seq, as well as characteristic repeat extension mutations in scRNA-seq but not in scATAC-seq, has strongly suggests that most of the scRNA-RPs are false positives generated by RNA-modality-specific effects. These effects may be the result of scRNA-seq-specific qualities such high coverage in low-complexity regions (3' UTR, 5'UTR and intronic regions) and poly-adenylation, although further investigation would

be required to confirm this. In scRNA-seq, though these false positives are frequent, it may be possible to reduce their incidence. We recommend others consider removing repeat-extension variants of the forms TT[C]TT>TT[T]TT, GG[C]NG>GG[G]NG, and CC[T]CN>CC[C]CN. On the other hand, scATAC-RPs seem to contain a more prominent mixture of real variants among the noisy variants. As we have not discovered an easy way to separate these, it appears that noise must simply be accepted.


Single-cell-unique Variants

As described previously, there is an abundance of variants that are unique to single-cell sequencing modalities, which tend to present in a low number of reads and hence a low number of cells. Previous studies have attributed these variants to real mutations unique to small cell populations caused by high mutation rates in cancer. Given the incidence of systematic errors in regions of high coverage or motif repeats (Stoler and Nekrutenko 2021; Tørresen et al. 2019; Li 2014), variant loci called on loci with a low number of supporting reads should be treated with caution. Before any claims of rare single-cell-unique variants may be made, it is essential to determine what portion of these variants may be the result of noise or error.

To estimate the potential of rare single-cell-unique variants in our single-cell libraries, we quantify scu-variants in scRNA-seq and scATAC-seq. In line with previous results, we find abundance of scu-variants (average of 1180 per sample at this filtering level) for scRNA-seq (see Table 3).

To evaluate whether these scu-variants are the result of rapid mutations characteristic of cancer, we extract the scu-variants from an unmatched control for which we have scRNA-seq and scATAC-seq, but no WGS. To account for the lack of matched WGS, we filter out all variants seen in the intersection of the common ground truth set from each of the sarcoma samples. We find considerably more scu-variants are generated (6600 for CTRL vs average of 1170 for EPSRC). Repeating the analysis performed by (N. M. et al. 2021), if we examine the breakdown of scu variant locations and impact annotated by

1 SnpEff (Cingolani et al. 2012) as seen in Figure 7, we find a breakdown very similar to that of the original
 2 paper, with 3' UTR variants being extremely common and an abundance of missense and intergenic
 3 variants (see Figure 7a). The rare positive variants called on pooled reads (referred to as URP variants in
 4 Table 1) also demonstrate breakdowns similar to (N. M. et al. 2021) (see Figure 7b). Although our
 5 breakdowns are not directly comparable to previous results due to a difference in the annotation tool
 6 used (SnpEff rather than SeattleSeq), these similarities give confidence that we have reproduced the
 7 scu-variants seen in previous studies.



9 **Figure 7:** Breakdown of variant annotations for (a) scu-variants and (b) rare positive variants

10 derived from scRNA-seq for each sample

11 Notably, the distribution of both scu-variants and rare positive variants in our wild-type control closely
 12 follows that of the sarcomas. A Chi-squared test of the category counts for scu-variants across samples
 13 results in a p-value of 0.0004 (see Table 5), which is mostly due to the decreased incidence of intergenic
 14 and non-coding variants seen in EPSRC1. If intergenic and non-coding categories are removed, or
 15 EPSRC1 is removed from comparison, the p-value shifts to non-significant values of 0.357 and 0.359

respectively, indicating strong agreement. Because intergenic and non-coding regions typically have poor coverage in scRNA-seq, we are inclined to conclude these extra variants in EPSRC1 are due to noise, possibly resulting from differences in the coverage profile of the scRNA-seq library sampled from EPSRC1. This overall result demonstrates that scu-variants are found in both cancerous and healthy tissue consistently across the same genomic locations and therefore cannot be easily attributed to low frequency somatic mutations associated with cancer development as previously described.

	EPSRC1	EPSRC2	EPSRC4	CTRL
3_prime_UTR_variant	135	106	333	450
missense_variant	33	31	66	106
synonymous_variant	10	9	26	47
5_prime_UTR_variant	3	5	24	40
stop_gained	3	0	3	5
other	6	6	19	34
Total	190	157	471	682

Table 5: Breakdown of scRNA-seq scu-variants by predicted location/impact, “other” represents all variants not in one of the existing categories

The abundance of scu-variants seen in Chromium scRNA-seq appears to be modality-specific, as performing the same analysis on scATAC-seq returns on average ~230x less variants for the same filtering levels (average 27 scu-variants, see Table 3). This is surprising considering the significantly higher breadth of coverage seen in scATAC-seq vs scRNA-seq. Although the two modalities cover different genomic regions, with scRNA-seq focusing on the exome, we would not expect the background somatic mutation rate to vary so drastically by region. This discrepancy further implies that the abundant scu-variants seen in scRNA-seq are the result of RNA-specific artifacts.

Overall, while we do not dismiss the possibility that scu-variants may capture rare cancerous somatic mutations, the abundance of scu-variants in scRNA-seq seen in both normal and cancerous tissue suggests that isolating rare somatic variants from background noise in scRNA-seq would be a considerable challenge. The lack of scu-variants in scATAC-seq is further evidence that those seen in scRNA-seq are due to noise. As such, we conclude that the utility of scu-variants as candidate rare somatic variants is questionable at best.

RNA-Editing Events

RNA-editing events are a type of post-transcriptional modification that can alter the sequence of an mRNA transcript to modify the function or localization of the resulting protein (Nishikura 2010). RNA-editing is widespread among eukaryotic organisms and specific modifications are often tissue-specific (Picardi et al. 2015; Song et al. 2004). There is a considerable interest in furthering our understanding of editing sites, such as the resulting impact on function and the context in which it occurs, as RNA-editing has been suggested as a safer method of genomic alteration for cancer therapy (Reardon 2020). As a result, there is a demand for approaches that can resolve RNA-editing at the cell-type level and link this to cell activity.

RNA-sequencing can capture RNA-editing events that occurs on mRNA strands as they are reverse-transcribed to cDNA for tagging and amplification. Moreover, scRNA-seq methods may resolve this at the single-cell level, enabling cell-type specific characterization of editing sites at a much finer level of detail than can be achieved with many bulk-RNA-seq libraries. Thus, scRNA-seq is uniquely suited to provide additional insight into RNA-editing. However, some studies have questioned the ability to resolve RNA-editing events in tag-based scRNA-seq approaches such as Chromium (Chen et al. 2019) We set out to determine if RNA-editing events are visible in Chromium scRNA-seq libraries and to determine if they may resolve cell-type-specific effects.

1 Prior work has amalgamated libraries of common RNA-editing sites in mice (Mansi et al. 2021). Using
2 overlap between scRNA-RP variants derived from our scRNA-seq datasets and the known RNA-editing
3 sites, we attempted to characterize the visibility of RNA-edits in Chromium 3' scRNA-seq. Across our 4
4 scRNA-seq samples we find a total of 4942 unique variant sites (average 1870 per sample) matching
5 RNA-editing events in the REDportal database, out of a total of 107,095 variants (average 1.75%
6 detection rate). For context, only 2 and 31 matches were discovered for common ground-truth and
7 scATAC-RP variants respectively. The coverage of editing sites seen in scRNA-seq can vary, with most
8 demonstrating low coverage. However, many editing sites appear in a sufficient quantity of reads such
9 that many cells within a sample demonstrate visibility of the editing site (Figure 8a). Given that editing
10 alleles often appear on a fraction of reads (Figure 8a), it is possible that a considerable number of editing
11 sites may have relevance in cell-type-specific analysis of editing events.

12 Most editing sites did not present in an adequate number of cells to infer cell-type-specific editing.
13 However, we found one site at Chr 7:130985524 presenting as an A to G base substitution that
14 demonstrated differential presentation of the RNA edit based on the expression profile of the cells in
15 the sample. This was found in EPSRC1, and we display the presentation of the RNA-edit allele
16 presentation frequency per cell, termed VAF, on the UMAP embedding of expression generated using
17 the Seurat R package (Figure 8d). For comparison, we used the expression profile of cells to perform
18 unsupervised Louvain clustering for any cell-type heterogeneity within our tumor sample (Figure 8b). As
19 we can see, the VAF varies considerably across the sample, with clusters 5 and 2 being enriched for the
20 RNA-edit and clusters 3 and 4 showing an absence of the edit ($p=4.0e-14$ by Fisher's exact test using
21 alternate allele and reference allele counts across cluster categorization). As can be seen in Figure 8c,
22 this is not due to differences in expression levels, as the gene on which the edit is located (*Htra1*) is
23 expressed in all involved clusters.




Figure 8: (a) Summary of RNA-editing events seen across scRNA-seq samples, demonstrating high-incidence of editing sites and presence in both low and high portions of reads, (b, c, d) UMAP embedding of cells in EPSRC1 scRNA-seq sample based on per-cell expression profile, with (b) clusters derived from Louvain clustering of normalized log counts per million for each gene after regressing cell-cycle effects to demonstrate the heterogeneity of cells, (c) expression level of the gene on which the RNA-edit of interest is located ("LOG(CPM)", logarithm of counts per million), demonstrating a uniform expression profile, (d) RNA-edit allele presentation frequency (VAF) differing across cells in a pattern similar to expression-based cell heterogeneity

Overall, this constitutes evidence that RNA-editing events may be seen in Chromium scRNA-seq libraries and used to infer heterogeneous presentation of RNA-editing. Although only one differentially presented site was found in our samples, there are many other sites with high coverage. Given that our samples are relatively homogeneous with respect to cell-type compared to other studies due to all being sampled from a mesenchymal tumor phenotype, it is possible that performing the same analysis on Chromium scRNA-seq libraries composed of multiple distinct cell types would yield additional cell-type-specific presentation of RNA-edits. However, we realize that a fair number of the variants found to match known PTM sites may be false positives due to noise discussed in previous sections. Furthermore, without a ground truth dataset of PTMs to compare, we cannot measure the accuracy of the approach. Thus, significant further study is required to confirm whether the results we have found represent real measurable RNA-editing events in Chromium scRNA-seq.

Discussion

Previous studies have applied various variant-calling approaches to Chromium-based scRNA-seq and scATAC-seq libraries without full consideration of the potential false-positives and artifacts generated by such approaches. In particular, traditional callers applied to pooled and individual cells are frequently used. Our analysis shows that a false discovery rate of around 0.4 and 0.2 for Chromium scRNA-seq and scATAC-seq may respectively be expected for most pooled read callers on these modalities. Of the pooled read calling approaches, it appears that Strelka2, GATK HaplotypeCaller, and SAMtools Call appear to offer the best performance in both modalities.

Variant calling in Chromium single-cell sequencing modalities such as scRNA-seq and scATAC-seq is increasingly being done with the application of bulk variant callers at the individual cell level. The major perceived advantage of variant calling at the individual cell is the ability to resolve rare variants not otherwise detected in variant calling on pooled reads. However, based on the results of this study, such

1 approaches do not exceed the performance of pooled read calling for the detection of infrequently
2 presented variants and are limited by their high false-discovery rates. Additionally, the excess of variants
3 unique to individual-cell callers appear to mostly consist of noise or artifacts specific to the sequencing
4 modality. We provide evidence of this fact in contrast to previous studies claiming that such variants
5 might be rare somatic variants present in a small sub-group of cells. While we do not refute the
6 presence of rare somatic variants detected with the single-cell approach, we reason that it would be
7 difficult to separate these from the number of false-positive variants generated without additional
8 validation with high-depth DNA sequencing. As a result, we conclude that variant calling at the single-
9 cell level offers no benefits over calling on the pooled reads in Chromium scRNA-seq and scATAC-seq,
10 and we recommend others to rely on variant callers such as Strelka, SAMtools, and HaplotypeCaller,
11 which demonstrate good performance in both modalities.

12 Through further investigation of variants discovered by variant calling on pooled reads, we find that
13 scRNA-seq generates a significant number of novel variants that are likely due to noise. We track this
14 down to repeat-extension variants, possibly due to the poly(A) tail or increased coverage in 3' and 5' low
15 complexity regions characteristic of the Chromium scRNA-seq modality. We recommend others
16 performing variant calling in Chromium scRNA-seq to remove variants with XX[Y]XX>XX[X]XX signatures,
17 especially if there is evidence that these compose a significant portion of the variants detected.

18 Finally, we find that Chromium scRNA-seq appears to detect instances of RNA-editing on the sequenced
19 transcripts. The incidence of these events is extensive and are found in many cells. We show that such a
20 signature can be investigated at the single-cell level to resolve cell-type-dependent RNA-editing for a
21 particular locus, provided there is sufficient coverage in cells. As such, we conclude that Chromium
22 scRNA-seq may be a useful tool for investigating RNA-editing events at the single-cell and cell-type level.

1 It should be noted that we are not discounting the possibility of rare somatic mutations in our data.
2 However, based on the number of likely artifactual variants we see both in tumors and wild-type
3 samples, it would be very difficult to resolve such variants if they exist without high-depth DNA-
4 sequencing for confirmation.

5 Of novel variants called on scRNA-seq, many of them appear to be due to repeat-extension artifacts
6 characteristic of Illumina sequencing. We believe this is due to higher coverage in low complexity
7 regions found in the 3' and 5' untranslated regions of genes. However, a certain portion of these
8 variants may be associated with RNA-editing events seen in other scRNA-seq methodologies. Our
9 preliminary results suggest that Chromium 3' scRNA-seq may be used to detect and characterize inter-
10 cell variations in RNA-editing which could significantly accelerate our understanding of this
11 phenomenon.

12 It is worth mentioning that it is possible that any of the variant calling approaches applied herein may be
13 considerably improved by a different application of filtering approaches. For ease of comparison, we did
14 not test filtering approaches exhaustively for all callers. Additionally, the quality metric used for
15 individual-cell callers, namely maximum overall variant quality, may not be optimal, but is most similar
16 to what is typically used. Of alternative approaches we tested, such as minimum number of cells in
17 which the variant was visible or average quality, we found it to perform the best. More work to
18 investigate other filtering techniques may improve results.

19 Overall, variant calling in Chromium scRNA-seq and scATAC-seq have great promise and good results can
20 be achieved using basic variant calling approaches. We offer a broad review and empirical evidence of
21 the performance of such approaches which should be helpful to anyone wishing to apply them to their
22 own sequencing libraries.

Methods

Mouse Model and Tumor Samples

All samples used in this study were selected from an inbred mouse line derived from the C57BL/6J line. *Smarchb1* (*Smarchb1^{ff}*, control *Smarchb1^{f/+}*) deletion was induced in the *Hic1^{CreERT2}* background (Scott et al. 2019), including a *B6.Cg-Gt(ROSA)26Sortm^{14(CAG-tdTomato)Hze}/J* (Jax stock number 007914) reporter allele, following administration of tamoxifen (TAM) at ~ 8 weeks of age. Epithelioid-like tumors appeared in various regions of the mouse with a latency of 10-14 months post-TAM weeks. From this study, three tumor samples from different animals labelled as EPSRC1, EPSRC2, and EPSRC4 were taken for further processing. Tumor resections were gently minced using scissors into 4 mm pieces and enzymatically dissociated using a cocktail of 1.5 U/mL Collagenase D (Roche 11 088 882 001) and 2.4 U/mL Dispase II (Roche 04 942 078 001) for 60 minutes. The resultant cell suspension was passed through a 40um filter and centrifuged at 500g for 5 mins. Cells were then incubated for 30 mins on ice with the following lineage antibody cocktail [anti-CD45-Alx647 (Ablab 67-0047-01, 1:400), anti-CD31-APC (BD Biosciences 551262, 1:400), anti-CD11b-Alx647 (Ablab 67-0055-01, 1:500), anti-F4/80-Alx647 (Ablab 67-0035-05, 1:500), anti-CD117-APC (eBioscience 17-1172-82, 1:500) and anti-Ter119-647 (Ablab 67-0031-01, 1:200) dissolved in FACS buffer (PBS with 2% FBS and 2mM EDTA] before fluorescence-activated cell sorting (FACS) against lineage markers to reduce the number of contaminating cell types and for tdTomato positive cells on an Influx (BD) sorter. Samples were then prepared and sequenced using Chromium-based scRNA-seq and scATAC-seq workflows (10x Genomics) as well as WGS (Illumina). This resulted in a highly enriched collection of tumor cells for each sample.

Sample Preparation and Sequencing

For scRNA-seq, sorted cells were prepared using the Chromium Next GEM Single Cell 3' v3 kit (10x Genomics). Cells were sequenced on the NextSeq 500 platform (Illumina). Reads were processed first

using Cell Ranger v3.0.2 (10x Genomics) on default settings. Cell Ranger performs a variety of preprocessing steps including unique molecular identifier (UMI) and cell barcode (CB) identification and denoising, adapter trimming, and duplicate marking, as well as alignment with STAR (Dobin et al. 2013). On average, there were 7900 cells per sample and 62000 reads per cell.

For scATAC-seq, sorted cells were prepared using the Chromium Single Cell ATAC v1 kit (10x Genomics). Cells were again sequenced on the NextSeq 500 platform (Illumina). Reads were processed using Cell Ranger ATAC v1.1.0 (10x Genomics) on default settings which implements BWA-MEM for alignment (Li 2013) and custom cell barcode identification and cell filtering steps. There was an average of 12400 cells per sample and between 5000 and 22000 median fragments per cell.

Concomitantly to single-cell capture, additional cells were centrifuged, flash frozen and banked. Thawed cells were further processed for WGS. Control tissue was obtained from age matched *Hic1^{CreERT2}*; *R26^{tdTomato}*, *Smabcb1^{+/+}* controls and processed for WGS. Samples were prepared using the DNA PCR-Free Library Tagmentation Prep kit (Illumina). The libraries were sequenced on the HiSeq X platform (Illumina). Reads were aligned using BWA-MEM (Li 2013).

Variant Categories and Metrics

The full set of variant category definitions can be seen in Table 1. As a gold standard of common or high-allele-frequency variants which should be called in single-cell, we use the consensus of multiple variant callers on the matched WGS libraries for each sample. This gives a high confidence list of biological common variants present in each sample. Given that WGS is limited in its capability to detect variants present in a small number of cells, as compared to other modalities, we would expect that such rare variants would most often be undetected or indistinguishable from noise in WGS data. As such, we use WGS to generate what we call the *common ground truth set* given that it should accurately represent the content of high-allele-frequency variants in the sample.

Metric	Formula	Threshold
Precision	$\frac{TP}{TP + RP}$	across quality filter levels
Recall	$\frac{TP}{TP + FN}$	across quality filter levels
cTPR	$\frac{TP}{TP + FN}$	across alt-allele coverage levels
cFDR	$\frac{RP}{RP + TP}$	across alt-allele coverage levels

Table 6: Quick reference for definitions of metrics used frequently throughout this study

If a variant in the common ground truth set is not detected in single cells but the reads in that sample show sufficient evidence for alternate allele, it is defined as a *pooled false negative* and used to calculate precision. Furthermore, we define a *rare positive* as a variant that was detected in single-cell sequencing which was not present in the common ground truth set. It is possible that rare positives are rare variants detected due to the high depth and allelic imbalance present in scRNA-seq and scATAC-seq. However, it is also likely that many are false positives due to noise or artifacts. Despite the potential veracity of rare positives, we use them to calculate recall out of convenience.

We define *union rare positive* (URP) variants as the union of rare positive variant calls for all callers performed on pooled scRNA-seq or scATAC-seq reads. We additionally required them to pass a weak quality filter (QUAL > 10) to remove low-quality variants. To ensure equivalent comparison of modalities, variants in all regions were considered except where otherwise noted.

To determine the performance of callers in regions of low vs high alternate allele read coverage, we define a *common true positive rate* (cTPR) based on the portion of pooled true positive and pooled false negative variants (Table 6). Correspondingly, we define a *common false discovery rate* (cFDR) using the portion of rare positives and pooled true positive variants.

1 We cannot confirm rare-positive variants with our WGS libraries as they are not of high enough
2 coverage to resolve low-frequency variants with confidence. However, there is a potential approach to
3 estimate the rare variants present in our samples. A notable advantage of our dataset is the availability
4 of matched scRNA-seq and scATAC-seq for a specific sample. Given the much higher coverage and
5 potential for visibility of rare alleles in these modalities, variants seen in both scRNA-seq and scATAC-seq
6 but not WGS are arguably very likely to be real rare variants. Thus, to provide an estimate of the
7 incidence of rare variants in the absence of a high-coverage WGS or WES resource for verification, we
8 can leverage the overlap of URP variants in scRNA-seq and scATAC-seq. We use the URP variant seen in
9 scRNA-seq and scATAC-seq, keeping those that have at least 5 supporting reads in each modality and an
10 alternate allele frequency of at least 0.5% to remove sequencing noise. We then count the number of
11 base pairs with at least 5 reads of coverage in scRNA-seq, scATAC-seq, and overlapping between the two
12 to provide an estimate for the rare mutation rate per megabase. We term this the *estimated rare*
13 *variant rate*.

14 For comparison with existing studies (N. M. et al. 2021), we define a scu-variant as any variant that is
15 called at the single-cell level in scRNA-seq or scATAC-seq but is not called on pooled reads nor found in
16 WGS. To keep in line with previous studies, we restrict this definition to the intersection of scu-variants
17 called by both GATK HaplotypeCaller and strelka2 with quality greater than 100 and 40 respectively. We
18 further restrict this to variants which show no evidence of being called at a statistically significant level
19 in WGS or by bulk callers on pooled reads.

20 *WGS Common Ground Truth Variant Calling*

21 To ensure high quality of called variants, WGS-derived common ground truth variants were taken as the
22 intersection of multiple callers. We used GATK HaplotypeCaller (Poplin et al. 2018), strelka2 (Kim et al.
23 2018), SAMtools (Li 2011) and VarScan (Koboldt et al. 2009) in their germline calling modes on default

settings. For GATK, we performed recommended best-practices steps prior to variant calling including Picard MarkDuplicates and GATK Base Quality Score Recalibration. Filtering was then done for all caller results to keep high-quality variants. For strelka2 and VarScan, this involved keeping variants which passed all filtering steps. For GATK HaplotypeCaller, variants were filtered using the recommended hard filtering approach: for SNVs, $QD > 2.0$, $QUAL > 30.0$, $SOR < 3.0$, $FS < 60.0$, $MQ > 40.0$, $MQRankSum > -12.5$, $ReadPosRankSum > -8.0$; for Indels, $QD > 2.0$, $QUAL > 30.0$, $FS < 200.0$, $ReadPosRankSum > -20.0$. For SAMtools, variants were filtered using the recommended hard filtering approach of $QUAL \geq 20.0$. Variants were added to the common ground truth set for a sample if they passed filters for at least three of the four callers.

Variant Calling on Pooled Reads from Single-Cell Libraries

Variant calling was performed on pooled reads from scRNA-seq and scATAC-seq libraries using Strelka2, SAMtools, GATK HaplotypeCaller, and FreeBayes. For HaplotypeCaller, pooled reads were pre-processed according to modality-specific recommended best practices. For scRNA-seq, this included SplitNCigarReads and Base Quality Score Recalibration, while for scATAC-seq, the DNA-seq variant-calling best practices approach was assumed and thus only Base Quality Score Recalibration was performed. SAMtools and HaplotypeCaller were run in their default configuration. Strelka2 was run using its new --rna mode, in which many filters are loosened, and a new scoring model is used. This was applied for both scRNA-seq and scATAC-seq, as it was reasoned that these new filters, especially relating to allele frequencies, would enable better sensitivity in both modalities.

Variant Calling on Individual Cell Reads from Single-Cell Libraries

To prepare reads to be called individually at the single-cell level, a custom Python script was used to split SAM/BAM read files by cell barcode into individual files for each cell. Libraries were split using the whitelist of cell barcodes generated by the CellRanger pipeline on its default settings for scRNA-seq and

scATAC-seq respectively, which uses a set of custom criteria for filtering out cells which are deemed to be low quality. All reads not having cell barcodes on the list being discarded. After splitting, variant callers were run on each individual cell BAM file to generate a list of variants which were then annotated with SnpEff (Cingolani et al. 2012). A custom Python script was used to collect variants called in each individual cell into one file and summarize collective statistics for each variant. A quality score was assigned to each variant, defined as the maximum quality seen for that variant across all cells in the sample.

Two bulk variant callers, Strelka2 and GATK HaplotypeCaller, were applied at the single cell level. For HaplotypeCaller, reads were preprocessed according to recommended best practices prior to read splitting. Strelka2 was run in its --rna mode.

Additionally, SCcaller (Dong et al. 2017) was run as a purpose-built single-cell variant caller on each individual cell. SCcaller requires a list of known variants ideally called on matched DNA-seq for the sample in question to estimate allele frequencies in each region. For this purpose, the list of variants called on WGS was used for each sample. Additionally, SCcaller averages local allele frequencies following a moving window of length specified by the user. To keep this window on the same order of length as that of a protein, and thereby model the allelic bias associated with each protein, a window size of 5kb was used.

Defining scu-variants

Following previous studies, a single-cell-unique variant was defined as a variant which was detected in two independent individual-cell callers with sufficient quality score in at least one cell to pass a filtering threshold. The two callers used were Strelka and HaplotypeCaller, both applied at the single-cell level. Additionally, scu-variants had to be unique to single-cell calling and therefore any variants detected in WGS or by the application of variant callers on pooled reads were removed. We defined a bulk variant

as the union of variants called on the pooled single cell reads using Strelka, HaplotypeCaller, and SAMtools. The quality filters for scu-variants generated by Strelka and HaplotypeCaller were set to 40 and 100 respectively, and these were found to be the quality past which the number of variants stopped decreasing significantly for any further increase in quality.

Coverage and False Negative Quantification

To determine if a variant found in WGS was visible in single-cell libraries, Vartrix by 10x Genomics was used to count the number of reads per cell containing a variant of interest. Possorted BAM files from CellRanger were run on Vartrix using the consensus variants from matched WGS. Reads with mapping quality less than 10 were removed. For scRNA-seq, reads have molecular barcodes and were thus deduplicated using the umi mode; in scATAC-seq, reads were deduplicated with the no-duplicates option. Variant loci were counted by number of alternate/reference allele reads per cell. Missing alleles were defined as a locus which did not show evidence of the allele seen in WGS but had coverage. False negative calls were defined as loci which showed evidence of the alternate allele but were not called as such by the variant caller.

For situations where GATK HaplotypeCaller was used for variant calling, all the above steps were run to generate a count on the BAM file after preprocessing according to GATK recommended best practices for each modality.

To check variants found in single-cell libraries were visible in WGS libraries, GATK ASEReadCounter was used. Variants were treated as heterozygous and used to count the presentation of alternate and reference alleles on reads passing default quality filters.

To determine the regions and depth of coverage of the genome for BAM files, BEDTools (Quinlan and Hall 2010) was used. From this toolkit, functions genomecov, merge, and intersect were used to

determine regions of genome coverage, combine regions with coverage to contiguous sets, and determine overlapping regions of coverage between two BAM files respectively.

Benchmark Scoring

We use quality filtering measures at a variety of thresholds to generate precision-recall curves. For bulk-calling methods, the default variant quality returned is used. For individual-cell calling, a list of variants is available for each cell. To merge these into one list, the union of variants across all cells was taken and any variants appearing in multiple cells were counted once. The maximum of variant call qualities over all cells in which the variant was detected was used as an aggregate quality measure.

For scATAC-seq, variants in any region are considered. As the validity of scRNA-seq results may extend only to exonic regions, we include the results for scRNA-seq limited to exonic regions in Figures 1, 2, and 3, but also include the results for scRNA-seq in all regions in Supplemental Figure S2, which demonstrate a substantially increased incidence of false positives.

Expression Level Analysis and Plotting

Expression-level analysis was done in R version 4.0.3 (R Core Team) using the Seurat package version 4.1.0 (Hao et al. 2021) to perform loading and manipulation of expression counts generated by Cell Ranger for each sample.

Some plots demonstrating cell-cell similarity based on expression were generated using the Seurat package. Cells were filtered based on percent-mitochondrial rna, number of total visible genes, and number of RNA reads according to Seurat recommended practices. The 2000 most variable features were kept. Features were reduced to the first 50 components found by principal component analysis. The cell-cycle impact on gene expression was regressed out by identifying cell-cycle genes based on the Harvard Chan School Bioinformatics Core tinyatlas annotations followed by repeated PCA analysis. Cells

were clustered using Louvain clustering at 0.8 resolution and plotted using UMAP (McInnes et al. 2020) on all 50 PCA components.

Data Access

The scRNA-seq data used in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; <https://www.ncbi.nlm.nih.gov/geo/>) under accession number GSE213338.

The scATAC-seq data used in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; <https://www.ncbi.nlm.nih.gov/geo/>) under accession number GSE213503.

The WGS data used in this study have been submitted to the NCBI BioProject database (<https://www.ncbi.nlm.nih.gov/bioproject/>) under accession number PRJNA1099909.

The source code including all helper scripts is available at https://github.com/mvjw/variant_stats and is also attached in the supplemental file titled Supplemental Code.

Competing Interests

The authors declare no competing interests.

Acknowledgements

This work was supported by the following grants: Terry Fox New Frontiers Program Project Grant 1021 (TMU), National Institutes of Health grant CA231652 (TMU), and NSERC Discovery Grant (AB). We would also like to thank the following facilities for technical assistance: BRC-seq, ubcFLOW cytometry, AbLab, BRC genotyping and BRC transgenic unit.

The authors confirm contribution to the paper as follows: study conception and design: M. Wiens, A. Bashashati, H. Farahani, R. W. Scott, T. M. Underhill; data collection: R. W. Scott, T. M. Underhill; analysis and interpretation of results: M. Wiens, A. Bashashati, H. Farahani, R. W. Scott, T. M. Underhill;

- 1 draft manuscript preparation: M. Wiens; supervised the study: T. M. Underhill, A. Bashashati. All authors
- 2 reviewed the results and approved the final version of the manuscript.
- 3

References

- Buenrostro JD, Wu B, Litzenburger UM, Ruff D, Gonzales ML, Snyder MP, Chang HY, Greenleaf WJ. 2015. Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* **523**: 486–490.
- Chen G, Ning B, Shi T. 2019. Single-Cell RNA-Seq Technologies and Related Computational Data Analysis. *Front Genet* **10**. <https://www.frontiersin.org/article/10.3389/fgene.2019.00317> (Accessed April 4, 2022).
- Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w¹¹¹⁸; iso-2; iso-3. *Fly (Austin)* **6**: 80–92.
- Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM, et al. 2021. Twelve years of SAMtools and BCFtools. *GigaScience* **10**: giab008.
- Ding J, Lin C, Bar-Joseph Z. 2019. Cell lineage inference from SNP and scRNA-Seq data. *Nucleic Acids Res* **47**: e56.
- Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–21.
- Dong X, Zhang L, Milholland B, Lee M, Maslov AY, Wang T, Vijg J. 2017. Accurate identification of single nucleotide variants in whole genome amplified single cells. *Nat Methods* **14**: 491–493.
- Dou J, Tan Y, Kock KH, Wang J, Cheng X, Tan LM, Han KY, Hon C-C, Park W-Y, Shin JW, et al. 2023. Single-nucleotide variant calling in single-cell sequencing data with Monopogen. *Nat Biotechnol* 1–10.
- Fan J, Lee H-O, Lee S, Ryu D, Lee S, Xue C, Kim SJ, Kim K, Barkas N, Park PJ, et al. 2018. Linking transcriptional and genetic tumor heterogeneity through allele analysis of single-cell RNA-seq data. *Genome Res* **28**: 1217–1227.
- Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing. <http://arxiv.org/abs/1207.3907> (Accessed October 31, 2022).
- Gawad C, Koh W, Quake SR. 2016. Single-cell genome sequencing: current state of the science. *Nat Rev Genet* **17**: 175–188.
- Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C, Zager M, et al. 2021. Integrated analysis of multimodal single-cell data. *Cell* **184**: 3573–3587.e29.
- Jaitin DA, Kenigsberg E, Keren-Shaul H, Elefant N, Paul F, Zaretsky I, Mildner A, Cohen N, Jung S, Tanay A, et al. 2014. Massively Parallel Single-Cell RNA-Seq for Marker-Free Decomposition of Tissues into Cell Types. *Science* **343**: 776–779.

- 1 Kim S, Scheffler K, Halpern AL, Bekritsky MA, Noh E, Källberg M, Chen X, Kim Y, Beyter D, Krusche P, et al.
2 2018. Strelka2: fast and accurate calling of germline and somatic variants. *Nat Methods* **15**: 591–
3 594.
- 4 Koboldt DC, Chen K, Wylie T, Larson DE, McLellan MD, Mardis ER, Weinstock GM, Wilson RK, Ding L.
5 2009. VarScan: variant detection in massively parallel sequencing of individual and pooled
6 samples. *Bioinformatics* **25**: 2283–2285.
- 7 Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and
8 population genetical parameter estimation from sequencing data. *Bioinformatics* **27**: 2987–
9 2993.
- 10 Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
11 *ArXiv13033997 Q-Bio*. <http://arxiv.org/abs/1303.3997> (Accessed April 4, 2022).
- 12 Li H. 2014. Toward better understanding of artifacts in variant calling from high-coverage samples.
13 *Bioinformatics* **30**: 2843–2851.
- 14 Liu F, Zhang Y, Zhang L, Li Z, Fang Q, Gao R, Zhang Z. 2019. Systematic comparative analysis of single-
15 nucleotide variant detection methods from single-cell RNA sequencing data. *Genome Biol* **20**:
16 242.
- 17 Lu T, Park S, Zhu J, Wang Y, Zhan X, Wang X, Wang L, Zhu H, Wang T. 2021. Overcoming Expressional
18 Drop-outs in Lineage Reconstruction from Single-Cell RNA-Sequencing Data. *Cell Rep* **34**:
19 108589.
- 20 Mansi L, Tangaro MA, Lo Giudice C, Flati T, Kopel E, Schaffer AA, Castrignanò T, Chillemi G, Pesole G,
21 Picardi E. 2021. REDportal: millions of novel A-to-I RNA editing events from thousands of
22 RNAseq experiments. *Nucleic Acids Res* **49**: D1012–D1019.
- 23 Massarat AR, Sen A, Jaureguy J, Tyndale ST, Fu Y, Erikson G, McVicker G. 2021. Discovering single
24 nucleotide variants and indels from bulk and single-cell ATAC-seq. *Nucleic Acids Res* **49**: 7986–
25 7994.
- 26 McInnes L, Healy J, Melville J. 2020. UMAP: Uniform Manifold Approximation and Projection for
27 Dimension Reduction. *ArXiv180203426 Cs Stat*. <http://arxiv.org/abs/1802.03426> (Accessed April
28 4, 2022).
- 29 N. M. P, Liu H, Dillard C, Ibeawuchi H, Alsaedy T, Chan H, Horvath AD. 2021. Improved SNV Discovery in
30 Barcode-Stratified scRNA-seq Alignments. *Genes* **12**: 1558.
- 31 Navin N, Kendall J, Troge J, Andrews P, Rodgers L, McIndoo J, Cook K, Stepansky A, Levy D, Esposito D, et
32 al. 2011. Tumour evolution inferred by single-cell sequencing. *Nature* **472**: 90–94.
- 33 Nishikura K. 2010. Functions and Regulation of RNA Editing by ADAR Deaminases. *Annu Rev Biochem* **79**:
34 321–349.

- 1 Petti AA, Williams SR, Miller CA, Fiddes IT, Srivatsan SN, Chen DY, Fronick CC, Fulton RS, Church DM, Ley
2 TJ. 2019. A general approach for detecting expressed mutations in AML cells using single cell
3 RNA-sequencing. *Nat Commun* **10**: 3660.
- 4 Picardi E, Horner DS, Pesole G. 2017. Single-cell transcriptomics reveals specific RNA editing signatures in
5 the human brain. *RNA* **23**: 860–865.
- 6 Picardi E, Manzari C, Mastropasqua F, Aiello I, D’Erchia AM, Pesole G. 2015. Profiling RNA editing in
7 human tissues: towards the inosinome Atlas. *Sci Rep* **5**: 14941.
- 8 Poirion O, Zhu X, Ching T, Garmire LX. 2018. Using single nucleotide variations in single-cell RNA-seq to
9 identify subpopulations and genotype-phenotype linkage. *Nat Commun* **9**: 4892.
- 10 Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Auwera GAV der, Kling DE, Gauthier LD,
11 Levy-Moonshine A, Roazen D, et al. 2018. Scaling accurate genetic variant discovery to tens of
12 thousands of samples. 201178. <https://www.biorxiv.org/content/10.1101/201178v3> (Accessed
13 April 4, 2022).
- 14 Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
15 *Bioinformatics* **26**: 841–842.
- 16 Quinones-Valdez G, Fu T, Chan TW, Xiao X. 2022. scAllele: A versatile tool for the detection and analysis
17 of variants in scRNA-seq. *Sci Adv* **8**: eabn6398.
- 18 R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical
19 Computing, Vienna, Austria <https://www.R-project.org>.
- 20 Reardon S. 2020. Step aside CRISPR, RNA editing is taking off. *Nature* **578**: 24–27.
- 21 Scott RW, Arostegui M, Schweitzer R, Rossi FMV, Underhill TM. 2019. Hic1 Defines Quiescent
22 Mesenchymal Progenitor Subpopulations with Distinct Functions and Fates in Skeletal Muscle
23 Regeneration. *Cell Stem Cell* **25**: 797-813.e9.
- 24 Song W, Liu Z, Tan J, Nomura Y, Dong K. 2004. RNA Editing Generates Tissue-specific Sodium Channels
25 with Distinct Gating Properties *. *J Biol Chem* **279**: 32554–32561.
- 26 Stoler N, Nekrutenko A. 2021. Sequencing error profiles of Illumina sequencing instruments. *NAR*
27 *Genomics Bioinforma* **3**: lqab019.
- 28 Tørresen OK, Star B, Mier P, Andrade-Navarro MA, Bateman A, Jarnot P, Gruca A, Grynberg M, Kajava
29 AV, Promponas VJ, et al. 2019. Tandem repeats lead to sequence assembly errors and impose
30 multi-level challenges for genome and protein databases. *Nucleic Acids Res* **47**: 10994–11006.
- 31 Vu TN, Nguyen H-N, Calza S, Kalari KR, Wang L, Pawitan Y. 2019. Cell-level somatic mutation detection
32 from single-cell RNA sequencing. *Bioinformatics* **35**: 4679–4687.

1 Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, Ziraldo SB, Wheeler TD, McDermott GP,
2 Zhu J, et al. 2017. Massively parallel digital transcriptional profiling of single cells. *Nat Commun*
3 **8**: 14049.

4 Zhou Z, Xu B, Minn A, Zhang NR. 2020. DENDRO: genetic heterogeneity profiling and subclone detection
5 by single-cell RNA sequencing. *Genome Biol* **21**: 10.

6