



## Probing the eukaryotic microbes of ruminants with a deep-learning classifier and comprehensive protein databases

Ming Yan, Thea Os Andersen, Phil B. Pope, et al.

*Genome Res.* published online December 27, 2024

Access the most recent version at doi:[10.1101/gr.279825.124](https://doi.org/10.1101/gr.279825.124)

---

**P<P** Published online December 27, 2024 in advance of the print journal.

**Accepted Manuscript** Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.

**Creative Commons License** This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <https://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

**Email Alerting Service** Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

---

**Comprehensive** immune receptor profiling.  
Discover the **DriverMap™ AIR Assay** difference.

LEARN  
MORE



---

To subscribe to *Genome Research* go to:  
<https://genome.cshlp.org/subscriptions>

---

Published by Cold Spring Harbor Laboratory Press

**Probing the eukaryotic microbes of ruminants with a deep-learning classifier and comprehensive protein databases**

Ming Yan<sup>1,2</sup>, Thea O. Andersen<sup>3,4</sup>, Phillip B. Pope<sup>3,4,5</sup>, and Zhongtang Yu<sup>1,2\*</sup>

<sup>1</sup> Department of Animal Sciences, The Ohio State University, Columbus, OH, USA

<sup>2</sup> Center of Microbiome Science, The Ohio State University, Columbus, OH, USA

<sup>3</sup> Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Ås, Norway.

<sup>4</sup> Faculty of Biosciences, Norwegian University of Life Sciences, Ås, Norway.

<sup>5</sup> Centre for Microbiome Research, School of Biomedical Sciences, Queensland University of Technology (QUT), Translational Research Institute, Woolloongabba, QLD, Australia.

Correspondence note:

Zhongtang Yu, Department of Animal Sciences, The Ohio State University, Columbus, OH 43210, USA.

Email: [yu.226@osu.edu](mailto:yu.226@osu.edu), Tel: +1-614-292-3057, fax: +1-614-292-2929

## Abstract

Metagenomics, particularly genome-resolved metagenomics, has significantly deepened our understanding of microbes, illuminating their taxonomic and functional diversity and roles in ecology, physiology, and evolution. However, eukaryotic populations within various microbiomes, including those in the mammalian gastrointestinal (GI) tract, remain relatively underexplored in metagenomic studies due to the lack of comprehensive reference genome databases and robust bioinformatics tools. The GI tract of ruminants, particularly the rumen, contains a high eukaryotic biomass although a relatively low diversity of ciliates and fungi, which significantly impacts feed digestion, methane emissions, and rumen microbial ecology. In the present study, we developed GutEuk, a bioinformatics tool that improves upon the currently available Tiara and EukRep in accurately identifying eukaryotic sequences from metagenomes. GutEuk is optimized for high precision across different sequence lengths. It can also distinguish fungal and protozoal sequences, further elucidating their unique ecological, physiological, and nutritional impacts. GutEuk was shown to facilitate comprehensive analyses of protozoa and fungi within more than one thousand rumen metagenomes, revealing a greater genomic diversity among protozoa than previously documented. We further curated several ruminant eukaryotic protein databases, significantly enhancing our ability to distinguish the functional roles of ruminant fungi and protozoa from those of prokaryotes. Overall, the newly developed package GutEuk and its associated databases create new opportunities for in-depth study of GI tract eukaryotes.

## Introduction

The complex and multi-kingdom microbial ecosystem unique to the rumen supplies ruminants with up to 70% of their energy requirement through the digestion and fermentation of plant materials (Bergman, 1990). However, this process also results in methane emissions, which account for approximately 14% of the anthropogenic greenhouse gas emissions (Gerber et al., 2013). The rumen microbes also convert dietary nitrogen, including protein and non-protein nitrogen, into high-quality

microbial protein, contributing up to 80% of the protein metabolized in the small intestine (Wallace et al., 1997). While the rumen microbiome is primarily linked to ruminant nutrition and production, the lower gastrointestinal (GI) microbiome is more associated with animal health and host physiology (Huws et al., 2018). The intricate and interdependent multi-kingdom ecosystem comprises anaerobic bacteria, archaea, fungi, protozoa, and viruses, each occupying a unique niche. While the prokaryotic communities of the rumen and intestines have been extensively investigated through multi-omics technologies, analysis of the eukaryotic communities still primarily relies on amplicon sequencing and analysis of a phylogenetic marker (metataxonomics), mainly the 18S rRNA gene for protozoa and internal transcribed spacer (ITS) for fungi, which provide limited taxonomic resolution and functional information. The lack of a comprehensive genome database for rumen protozoa and fungi further limits omics analyses of these eukaryotes. Our understanding of the functional importance of the eukaryotic communities within the rumen is still primarily based on culture-based and genomic studies of a small number of species.

Unlike the human gut, where eukaryotes account for less than 1% of the gut microbiome (Huffnagle & Noverr, 2013), fungi and protozoa constitute up to 20% and 50% of the rumen microbiome biomass, respectively (Andersen et al., 2023; Edwards et al., 2017; Huws et al., 2018). Rumen fungi are specialized fiber degraders capable of producing diverse carbohydrate-active enzymes (CAZymes), especially those hydrolyzing recalcitrant plant cell wall materials (Hagen et al., 2020), complementing the CAZymes repertoire of rumen bacteria (Hagen et al., 2020; Peng et al., 2021). In addition, gut fungi produce diverse secondary metabolites with potential antimicrobial and therapeutic properties (Swift et al., 2021). Except for *Pecoramyces* sp. F1, which was isolated from the rumen (Jin et al., 2011), all available genomes of gut fungi from ruminants, *Anaeromyces robustus*, *Caecomyces churrovius*, *Neocallimastix californiae*, *Neocallimastix lanati*, and *Pecoramyces ruminantium*, were obtained exclusively from fecal samples, therefore also representing hindgut fungi. Although the rumen and the hindgut share some of their fungi, they have distinctive fungal communities (Fliegerova et al., 2015), and it remains unknown whether the fungal enzymatic repertoire differs between the rumen and the hindgut. While the

contribution of fungi to fiber degradation in the rumen is well-recognized, the functions of rumen protozoa are intricate and multifaceted, encompassing productive roles such as fiber degradation and rumen pH stabilization, alongside counterproductive roles such as promoting methane emissions and wasteful intra-ruminal recycling of microbial protein (Firkins et al., 2020; Yu et al., 2024). Indeed, rumen protozoa exhibit a linear relationship with methane emissions (Guyader et al., 2014). Due to their negative associations with feed efficiency and methane emissions, a few studies have attempted to specifically inhibit the dominant species of *Entodinium* (Park, Mao, et al., 2019; Park, Yang, et al., 2019). Alternatively, analyses of single-cell amplified genomes (SAGs) and subsequent proteomic analysis of rumen protozoa revealed an unexpectedly extensive repertoire of various CAZymes (Andersen et al., 2023; Li et al., 2022). Furthermore, recent studies have demonstrated an association between rumen protozoa and feed efficiency in beef cattle (Clemmons et al., 2021) and between intestinal protozoa and intestinal health in calves (Fu et al., 2023). Therefore, it is imperative to transcend beyond metataxonomic analysis of rumen protozoa, characterizing the genomes and enzymatic capacity of the eukaryotic fraction in the GI tract of ruminants. Moreover, a comprehensive protein database is needed to enable robust and unbiased analyses of protozoal gene/protein expression profiles in relation to key rumen functions and animal production, such as methane emissions and feed efficiency.

Single-cell genomics has proven valuable in unraveling the fundamental biological characteristics of protozoa (Li et al., 2022). However, it is not cost-effective for ecological studies of protozoal communities. Metagenomics has revolutionized the investigation of various microbiomes, including the GI microbiomes of ruminants, but most studies focus on bacteria and archaea while ignoring fungi and protozoa. Two machine learning-based tools (Tiara and EukRep) are available to identify eukaryotic sequences directly from metagenomic assemblies (Karlicki et al., 2022; Pronk & Medema, 2022; West et al., 2018). Studies using these tools have revealed the ecological importance of protists in aquatic environments (Alexander et al., 2023; Duncan et al., 2022; Kavagutti et al., 2023; Saraiva et al., 2023). However, it remains unclear whether the above tools can be used to analyze eukaryotes abundant in host-


associated ecosystems like the rumen. Moreover, the diversity and functional repertoire of the ruminant GI tracts, especially the less-studied hindgut and ruminant species, remains limited. Therefore, our first objective was to benchmark the existing tools for classifying ruminant gut eukaryotes whilst creating a new tool (referred to as GutEuk) to identify and classify eukaryotic microbes (fungi and protozoa) in metagenomes derived from the gut of ruminants. Using GutEuk, we also aimed to reanalyze over one thousand metagenomes and metatranscriptomes from ruminant GI tracts and to create protein databases for ruminant gut eukaryotes. This analysis included samples from the foregut (mostly rumen but also including reticulum, omasum, and abomasum) and hindgut (cecum, colon, and rectum) across various ruminant species (mostly domestic species including *Bos taurus* and *Ovis aries* as well as wild species including *Capreolus capreolus* and *Hydropotes inermis*). Our last objective was to benchmark the newly established protein databases by improving metaproteomic analysis using data from a recent study (Andersen et al., 2023). Overall, GutEuk and the new protein databases it creates open up new opportunities in analyzing and reanalyzing host-associated microbiomes for eukaryotic microbes, especially those within the rumen.

## Results

### The new GutEuk improves identification of eukaryotic microbes from metagenomes

Tiara (Karlicki et al., 2022) and EukRep (West et al., 2018) are tools for identifying eukaryotic sequences from metagenomic assemblies based on *k*-mer frequency. They were developed based on a limited selection of eukaryotic nuclear genomes (73 for Tiara and 70 for EukRep), encompassing those of algae and animals. The lack of representation of gut-associated eukaryotic microbes likely undermines their utility in analyzing gut-derived metagenomes. In addition, *k*-mer frequency merely represents metrics of simplified genomic contexts, while improved classification can be achieved using a convolutional neural network (CNN) with original DNA sequences as inputs, as illustrated by a recent study (Hou et al., 2023). We tested Tiara and EukRep for their ability to identify gut eukaryotic sequences using a dataset from 52 SAGs of rumen protozoa (Li et al., 2022) (see Methods). Tiara and EukRep correctly classify only 30,081 and 44,395 out of 57,253 eukaryotic contigs, achieving accuracies of 52.5% and 77.5%, respectively, which are much lower than their reported performance.

The aforementioned accuracies of Tiara and EukRep indicate that these two existing tools cannot achieve a generalizable performance for less-represented genomes of eukaryotes. We thus developed a specialized deep-learning classifier built upon diverse protozoal, fungal, and prokaryotic genomes to improve the classification of eukaryotic microbes in the rumen and gut. Leveraging thousands of prokaryotic, fungal, and protozoal genomes, we developed GutEuk, an ensembled deep-learning model combining CNN (one-hot encoded nucleotide bases as inputs) and feedforward neural network (FNN; *k*-mer frequencies as inputs) (Fig. 1A and 1B). GutEuk employs a two-stage classification with user-defined confidence levels at each stage: The first stage differentiates between prokaryotic and eukaryotic (fungal and protozoal) sequences, and the second stage further distinguishes fungal from protozoal sequences.




**Fig. 1:** Illustrative schematic of the two-stage classification process used by GutEuk (A), the pipeline to classify individual contigs and bins from metagenomes (B), the workflow to establish the ruminant foregut and hindgut eukaryotic protein databases (C).

We benchmarked the performance of GutEuk against EukRep and Tiara using independent datasets that were not used to train these tools at the contig level (see Methods). GutEuk outperformed Tiara and EukRep consistently at different contig lengths (Fig. 2A). GutEuk and Tiara exhibited comparable precision for eukaryotic contigs, while EukRep displayed a slightly lower precision, especially for contigs shorter than 20kb. However, for prokaryotic contigs, GutEuk achieved a consistent

precision of around 99%, even for contigs shorter than 15kb, whereas Tiara and EukRep achieved a precision of 95% and 91%, respectively, for contigs shorter than 10kb. The performance of Tiara and EukRep was poorer than that of GutEuk for contigs between 3 and 5kb. In terms of recall rate, all three tools performed similarly with prokaryotic contigs. However, for eukaryotic contigs, GutEuk achieved a recall rate of around 99% for contigs shorter than 15kb and still maintained a recall rate of around 98% for longer contigs. In comparison, EukRep and Tiara recorded lower recall rates, with EukRep achieving at best 95% and dropping below 90% with contigs shorter than 10kb, while Tiara achieved only 82% at best with contigs between 5 and 10kb, with lower recall rates with longer contigs. Regarding computing time, GutEuk processed the test dataset of 965,741 sequences (totaling 4.6 Gb) in 9.3 hours using 20 threads. In comparison, Tiara and EukRep completed the same task with the same resources in 0.3 hours and 2 hours, respectively.


We evaluated GutEuk for its ability to further differentiate protozoal and fungal contigs (Fig. 2B). Since the maximum number of classifiable fragments is two for contigs shorter than 15kb, the threshold used will not affect its performance (both fragments must provide the same prediction). For contigs exceeding 15kb, we observed the expected increase in precision and decrease in recall as the confidence level ( $pt_2$ ) rose. Precision exceeded 90% even at a  $pt_2$  of 0.5 and reached 95% at a  $pt_2$  above 0.8. In terms of recall rate, GutEuk achieved over 70% and 85% for protozoa and fungi, respectively, at a  $pt_2$  of 0.8. Therefore,  $pt_2$  can be adjusted to optimize GutEuk for precision and recall based on specific research objectives. Nevertheless, a balance between precision and recall was seen at a  $pt_2$  of 0.8, and the corresponding precision of above 95% was considered conservative.



**Fig. 2: Benchmarking the performance of GutEuk, EukRep, and Tiara for contigs.**

**A**, The performance of GutEuk, EukRep, and Tiara in differentiating prokaryotic and eukaryotic contigs of different lengths. EukRep and Tiara were used with their default settings. GutEuk was run at a  $pt_1$  of 0.5. **B**, The performance of GutEuk in further differentiating fungal and protozoal contigs at different confidence levels ( $pt_2$ ) and contig lengths.

In addition to evaluating the performance of GutEuk in identifying microbial eukaryotic contigs, we further assessed whether it could achieve consistent results across genomes of diverse phylogenetic lineages, including aquatic fungi and protozoa included in the database (Fig. 3A-D). All three tools



**Fig. 3: Benchmarking the performance of GutEuk, EukRep, and Tiara with genomic bins.**


The percentage of genomes being correctly classified as prokaryotic or eukaryotic by Tiara (A), EukRep (B), and GutEuk (C). The percentage of genomes being correctly classified as prokaryotic, protozoal, or fungal by GutEuk (D).

achieved above 90% accuracy. However, Guteuk, leveraging an expanded training set and a more complex model, achieved over 90% accuracy in classifying DNA from most eukaryotic microbes, especially the underrepresented rumen ciliate protozoa. The only eukaryotic genome whose fragments were identified with a lower than 70% accuracy (about 45%) was that of *Stentor coeruleus* (accession id: GCA\_001970955.1), a freshwater protozoan. In contrast, Tiara and EukRep each had 14 eukaryotic genomes classified with an accuracy lower than 70%. These results indicate that at a  $pt_1$  of 0.5, GutEuk predicts eukaryotic microbial genomes/bins more accurately than Tiara and EukRep. More importantly, GutEuk can also further classify eukaryotic genomes as fungal or protozoal with high accuracy, except for a few fungi and protozoa not found in the mammalian GI tract. Specifically, GutEuk identified the following fungal genomes with lower than 50% accuracy: *Smittium mucronatum* (GCA\_001953115.1, an insect gut symbiont), *Vavraia culicis* (GCA\_000192795.1, a microsporidian parasite), and *Enterospora cankeri* (GCA\_002087915.1, infecting mosquito and European shore crab). Similarly, protozoal genomes identified with less than 50% accuracy are not found in the GI tract either: *Perkinsus marinus* (GCA\_000006405.1) and *Perkinsus* sp. *BL\_2016* (GCA\_004369235.1), two parasitic flagellates infecting mollusks, as well as *Stentor coeruleus* (GCA\_001970955.1) and *Lenisia limosa* (GCA\_001655205.1), two aquatic ciliates. Given the narrow phylogenetic range of gut eukaryotes, GutEuk is expected to perform well with gut metagenomic sequences (see the next section). A higher  $pt_2$  threshold (e.g., 0.8) can be used if a lower false positive rate is required. Furthermore, GutEuk correctly identified 697 out of 708 (98.4%) newly sequenced human gut fungal genomes (Yan et al., 2024) as fungal (Supplementary Table S1).

## Diverse rumen protozoa are identified from underrepresented ruminant species


To date, 53 genomes of rumen ciliates have been sequenced, including 52 SAGs (Li et al., 2022) and one mono-species genome (Park et al., 2021). These genomes represent 19 species of rumen ciliates prevalent across various ruminant species. Using GutEuk, we systematically analyzed 15,000 bins (each exceeding 3 MB) assembled from 1,093 previously sequenced rumen metagenomes (Supplementary Table S2, see Methods) for eukaryotic bins. These samples were collected from 13 different ruminant species, encompassing both domestic and wild species with varying feeding regimes. We identified hundreds of protozoal bins and one fungal bin. The scarcity of bins identified as fungal likely reflects their low abundance and low fungal DNA yield from rumen fluid samples because rumen fungi strongly adhere to digesta particles. We assessed the genome completeness of the identified eukaryotic bins based on the presence of lineage-resolved single-copy marker genes, finding that most bins were largely incomplete, consistent with findings from a recent study using EukRep for eukaryotic bin identification (Saraiva et al., 2023).

Despite the majority of the eukaryotic bins identified from the rumen metagenomes being incomplete, we identified 21 protozoal bins with more than 50% completeness and less than 10% contamination. We further conducted a phylogenetic analysis on this set of protozoal bins and the rumen protozoa SAGs (Fig. 4). Examination of the phylogenetic tree, constructed from 30 concatenated single-copy marker proteins, reveals that the newly discovered rumen protozoal bins cluster with the SAGs within the Vestibuliferida order. In contrast, all Entodiniomorphida SAGs form a distinct cluster, suggesting a potentially greater diversity of Vestibuliferida species yet to be uncovered. Notably, the phylogenetic tree constructed with the 30 single-copy marker proteins exhibited the same topology as the original phylogenetic tree of the SAGs, which was based on 113 single-copy marker proteins. This consistency demonstrates the robustness of our phylogenetic analysis. It is also noteworthy that some of the newly identified protozoal bins were derived from metagenomes of ruminant species, including *Bison bison* and *Bos indicus*, from which no protozoal genomes have been previously sequenced. Therefore, future protozoal research should expand to include less-studied ruminant species to facilitate a more comprehensive genomic representation of rumen eukaryotes.



**Fig. 4: A phylogenomic tree of the identified rumen protozoal bins and 49 rumen protozoa single-cells assembled genomes.** All nodes with a greater than 85% bootstrap support are indicated with dots in the branch.

**Protein databases of ruminant gut eukaryotes and the repertoire of genes encoding glycoside hydrolases and peptidases**



**Fig. 5: Diverse ruminant gut eukaryotic proteins identified in the databases.**

**A**, The number of protein clusters identified in foregut protozoal, foregut fungal, and hindgut eukaryotic protein databases. **B**, The most prevalent foregut protozoal glycoside hydrolases (GHs) identified. **C**, The most prevalent foregut protozoal secretory (with identified signal peptides) peptidases and peptidase inhibitors identified. **D**, the number of unique foregut protozoal secretory peptidases or peptidase inhibitors identified based on catalytic sites.

From the eukaryotic sequences (bins and contigs) identified from the gut metagenomes of various ruminants, predominantly cattle, sheep, and goats, we predicted and translated their protein sequences, subsequently constructing several protein databases (Fig. 1C). These gut eukaryotic microbial protein databases include one each for foregut protozoa (both newly identified and from the 52 SAGs), foregut fungi, and hindgut eukaryotes (Fig. 5A). The foregut protozoal database consisted of 25,498,129 proteins, which were clustered at 50% amino acid identity into 5,265,119 protein clusters, 54% of which are newly


identified. Among these clusters, only 10% were annotated. The foregut fungal database is much smaller, containing 11,314 proteins, which formed 7,293 protein clusters. Around half of the fungal protein clusters could be annotated. Notably, the proportion of the annotated hindgut eukaryotic protein clusters exceeded 92%. The markedly smaller proportion of annotated protozoan proteins relative to those of fungal and hindgut eukaryotes highlights the need for further genomic investigations into rumen protozoa. This need is also underscored by their complex functions and implications, such as predation on other rumen microbes, enzymatic breakdown of plant material, and symbiotic interactions with bacteria and archaea.

Previously, bacteria and fungi were credited for most of the cellulolytic activity within the rumen microbiome. However, a recent genomic study (Li et al., 2022) and a metaproteomic study (Andersen et al., 2023) have revealed a substantial proportion of genes encoding glycoside hydrolases (GHs), including cellulase and hemicellulases, within rumen protozoa (Andersen et al., 2023). In the present study, we annotated the foregut protozoal database for GHs. We found that amylases (e.g., GH13) and cellulases (e.g., GH5 and GH9) are among the most prevalent CAZymes identified (Fig. 5b), underscoring the significant fibrolytic and amylolytic capability of rumen protozoa.

Rumen protozoa are known for their ability to degrade protein, especially microbial protein. We thus also annotated the foregut protozoal database for peptidases. Among the peptidases identified, metallo, aspartic, cysteine, and serine peptidases were predominant, with M08, C01, A01, and S09 peptidases being among the most prevalent secretory peptidases (Fig. 5C). *Prevotella*, the most predominant genus of rumen bacteria with broad protease activity (Hartinger et al., 2018), exhibits 48 different families of secretory peptidases (Patra & Yu, 2022). In comparison, 58 different secretory peptidase families were identified from the rumen protozoal proteins (Fig. 5D), highlighting their substantial proteolytic potential. Furthermore, the identification of ferredoxin hydrogenase of rumen protozoa (Supplementary Table S4) corroborates their ability to produce hydrogen.

## **Ruminant gut eukaryotic protein databases improve protein identification in proteomics**

Recently, a metaproteomic study demonstrated the utility of protozoal SAGs and genomes as a reference search database in annotating rumen protozoal proteins and assessing the significance of rumen eukaryotes in rumen functions (Andersen et al., 2023). Given that reference search databases are a major bottleneck in metaproteomic studies of complex microbiomes, and considering the previously demonstrated high genomic and functional diversity of the rumen eukaryotes, we evaluated the potential of expanding protozoal genome databases with ruminant eukaryotic proteins identified with GutEuk to enhance protein detection from rumen metaproteomic data. Using the same pipeline and threshold as the original study (Andersen et al., 2023), we compared the number of protein groups identified in each sample using (i) the original genome database (consisting of one *Entodinium caudatum* genome and 18 SAGs of rumen protozoa) and (ii) an expanded database combining the original genome database with the GutEuk-identified rumen eukaryotic proteins. The expanded database enabled substantially more protozoal protein groups to be identified in both cow and goat rumen samples, with an average increase of 64% for cow samples and 20% for goat samples (Fig. 6). In some samples, an increase of around 200% was seen for protozoal protein groups, suggesting that previously unknown protozoa are likely dominant in certain samples. Similarly, the expanded database greatly increased the number of fungal proteins identified, with a 118% increase in cow samples and 70% increase in goat samples. Notably, on average, only 10 fungal protein groups were identified in the original study. It is also important to note that rumen samples were obtained for this particular metaproteomic study via esophageal tubing, which primarily collects rumen fluid and significantly lowers fungal biomass.



**Fig. 6: Increases in the number of rumen protozoal and fungal protein groups identified with the new protein databases compared with the previous rumen proteomic study.**

The numbers represent the percentage ratio between newly identified protein groups and those identified in the original study. Box plots indicate the median (middle line), 25th and 75th percentile (box), as well as individual observations (dots).

## Discussion

Historically, gut eukaryotes have been studied primarily from a parasitological perspective. However, eukaryotic populations are also ubiquitous components of the gut microbiome (Parfrey et al., 2014; Ramayo-Caldas et al., 2020), and they contribute significantly to the overall microbial assembly, host nutrition, and physiology (Gerrick et al., 2023; Hoffmann et al., 2013; Laforest-Lapointe & Arrieta, 2018). Their role is particularly significant in the rumen, the foregut of ruminants, where they can constitute up to 50% of the microbial biomass—a proportion much higher than in other ecosystems. These microbes play direct roles in rumen fermentation and fiber degradation (Huws et al., 2018) and directly influence several important host characteristics, including feed efficiency and methane emissions (Firkins et al., 2020). Despite recent efforts in the cultivation and genomic sequencing of rumen protozoa and fungi, which is a highly labor-intensive process, we still lack a robust bioinformatic tool to analyze the eukaryotic microbes represented in shotgun metagenomes generated from rumen samples. Therefore,

it is crucial to develop bioinformatic tools tailored to gut eukaryotic microbes, particularly those in the rumen, to obtain a more comprehensive representation of their genomes and proteomes. Such new tools will enable a shift from taxonomic profiling to functional profiling, allowing for a deeper understanding of gut eukaryotic microbes as well as whole gut microbiomes.

Tiara and EukRep have been utilized for eukaryotic sequence identification; however, they fail to accurately identify rumen eukaryotes. To facilitate metagenomic and functional analysis of gut eukaryotic microbes, especially those in the rumen, we developed GutEuk. This tool outperforms Tiara and EukRep in both precision and recall when identifying microbial eukaryotic genome fragments. The ensemble model implemented in GutEuk (Fig. 1), which combines CNN and FNN, enables consistent performance for both short sequences (shorter than 5 kb) and long sequences (longer than 50 kb). The ability of GutEuk to further differentiate fungal and protozoal sequences is especially valuable for analyses of metagenomes derived from microbiomes that contain both types of eukaryotic microbes, such as the rumen microbiome. Although GutEuk was designed for metagenomes from the GI ecosystem, it also demonstrates consistent performance across genomes of diverse taxonomic lineages by accurately differentiating fungi and protozoa from other ecosystems. Since eukaryotic communities are less diverse in the gut than in aquatic and terrestrial environments, and there is minimal overlap in eukaryotic microbial diversity among these ecosystems (Parfrey et al., 2014), GutEuk is well suited for gut metagenomic studies, as demonstrated by its use with the newly sequenced human gut fungi genomes (Supplementary Table S1). In addition to prokaryotes and eukaryotes, metagenomes can contain significant amounts of mobile genetic elements, such as plasmids and viruses. If these elements are of particular interest, we recommend using specialized tools like geNomad (Camargo et al., 2023) to filter the input sequences.

Using GutEuk, we screened thousands of metagenomic bins assembled from the rumen and identified hundreds of individual eukaryotic bins. Our phylogenetic analysis of the dozens of identified protozoal bins with at least medium quality demonstrated a great genomic diversity of rumen protozoa, particularly those in less-studied ruminant species. The relative sparsity of the recovered eukaryotic bins

with high quality may be attributable to multiple factors, including the low abundance of eukaryotes in the rumen, their large and complex genomes, high genetic heterogeneity, and the presence of highly repetitive genomic sequences (Cho et al., 2023). Additionally, there is a risk of chimeric bins due to the presence of mobile elements and paralogous genes (West et al., 2018). Annotation of eukaryotic bins should, therefore, be performed with caution. Binning of contigs into high-quality MAGs of eukaryotic microbes from metagenomic sequences, especially short sequencing reads, is probably challenging. Sequencing repetitive regions contiguously (Cho et al., 2023), long-read sequencing, such as PacBio HiFi and Oxford Nanopore sequencing, may facilitate assembling and binning of eukaryotic sequences from metagenomes, especially for protozoa that have highly fragmented small chromosomes, such as the *Entodinium* genus (Park et al., 2021). With the growing use of single-cell sequencing, GutEuk can be applied to filter out symbiotic prokaryotic sequences prior to assembly, enhancing genome quality due to its high accuracy in distinguishing between prokaryotic and eukaryotic sequences. Additionally, the recently developed bin refinement tool ACR (Seong et al., 2023) can enhance genome-centric analyses of eukaryotes through metagenomic approaches. Furthermore, similar to the assessment of prokaryotic genomes and MAGs, the completeness and contamination rates of eukaryotic bins should be determined based on the presence of lineage-specific marker genes and corresponding databases as implemented in BUSCO (Simão et al., 2015) and EukCC (Saary et al., 2020). However, both tools show limited performance when applied to eukaryotic clades with sparse reference genomes (Jauhal & Newcomb, 2021; Saary et al., 2020). With the expansion of reference genomes and advanced quality assessment using newly developed tools like ContScout (Bálint et al., 2024), we can better assess the quality of the obtained eukaryotic microbial genomes.

Several genomic structural features of eukaryotes, such as the exon-intron architecture and the lack of conserved splicing signatures, complicate the gene-calling process in eukaryotic genome annotations. To overcome these challenges, a reference- and homology-based pipeline, MetaEuk (Levy Karin et al., 2020), with high precision (>99.9%) has been developed (Levy Karin et al., 2020). Utilizing this pipeline and a database combining single-cell protozoal transcriptomes and poly(A) enriched rumen

metatranscriptomes, we screened the contigs assembled from around one thousand ruminant GI metagenomes (see methods). This analysis led to the identification of millions of ruminant eukaryotic proteins, further highlighting the great metabolic potential of rumen protozoa, consistent with the previous studies (Andersen et al., 2023; Li et al., 2022).

Leveraging the databases we compiled, we tested their resource applicability by reanalyzing the detected proteomic profiles of the rumen of cows and goats (Andersen et al., 2023). We substantially increased the number of identified protein groups from both fungi and protozoa. The additional proteins identified are primarily from the ruminant eukaryotic sequences identified by GutEuk, validating the genomic identification and protein prediction pipeline we used. The potential for chimeric binning is unlikely to confound the protein prediction as all the genomic sequences are predicted to be either fungal or protozoal. With the assistance of GutEuk and the derived protein database, we could better understand the distinct metabolic contributions of fungi and protozoa in the rumen, particularly under varying dietary and microbial interventions. However, protein prediction will require further genome curation, as described previously (West et al., 2018), to attribute expressed protein to specific species.

In summary, the new GutEuk tool, in conjunction with its associated databases, enables comprehensive analyses of GI eukaryotes in metagenomes. The integration of the proteins identified from rumen metagenomes into reference genome databases markedly enhances metaproteomic studies of the rumen microbiome. Additionally, the expanded databases enhance read-based ecological analyses of gut eukaryotic microbes with tools such as EukDetect (Lind & Pollard, 2021), offering finer taxonomic resolution than marker-based metataxonomic approaches. Collectively, GutEuk and the associated databases can help provide new insights into the functional contributions of ruminant fungi and protozoa to the rumen ecosystem, as well as their interactions with diets and other microbes.

## Methods

### Evaluating existing tools in identifying and classifying rumen protozoa

Currently, Tiara (Karlicki et al., 2022) and EukRep (West et al., 2018) are tools available for identifying eukaryotic contigs assembled from metagenomes. To evaluate their performance in identifying sequences of eukaryotes, we utilized the 52 rumen protozoa SAGs reported recently by Li et al. (2022). The telomere sequences were trimmed off from the telomere-capped contigs with a custom Python script (Supplementary Codes). Since both tools were trained with contigs of 5kb length, the contigs of each SAG were chopped into 5kb fragments with a 1kb sliding window and used for benchmarking (see below).

### Genome dataset preparation

To specifically identify eukaryotic sequences in metagenomes derived from the gut, we compiled a dataset comprising genomes of prokaryotes, fungi, and protozoa. Briefly, 85,205 representative genomes of prokaryotes were downloaded from GTDB release 214.0 (<https://data.gtdb.ecogenomic.org/releases/release214/214.0/>). These genomes represent 80,789 species of bacteria and 4,416 species of archaea. Fungal genomes were downloaded from the MycoCosm genome portal (<https://mycocosm.jgi.doe.gov/mycocosm/home>). When multiple genomes were available for a fungal species, we only selected one genome with the highest quality. We also filtered out the genomes of mushrooms and soil fungi, reducing the number of fungal genomes not relevant to gut microbiomes. A total of 679 fungal genomes were retained, representing 639 genera. Protozoal genomes were downloaded from the NCBI GenBank database on September 2<sup>nd</sup>, 2023, and we selected the most complete genomes per species. We combined these protozoal genomes with the 52 rumen protozoal SAGs representing 19 species across 13 genera (Li et al., 2022). To remove potential sequence contamination from symbiotic prokaryotes, we retained only the contigs flanked by the identified telomere sequences of protozoa. Additionally, we included only one representative species per genus from the fungal and prokaryotic genomes to minimize classification bias toward overrepresented lineages and to prevent data leakage. The

dataset includes 20,739 prokaryotic genomes/contigs, 639 fungal genomes/contigs, and 380 protozoal genomes/contigs.

The genomes/contigs from each class were randomly assigned to one of the following datasets: training (70%), validation (15%), and testing (15%) to be used to train the classifier (see below). Within each dataset, genomes or contigs were randomly fragmented into 5 kb fragments with a sliding window of 5 kb. Then, we randomly selected 28.5% of the prokaryotic and fungal fragments and 75% of the protozoal fragments. This selection ensured a dataset size balanced with prokaryotic and gut eukaryotic sequences. The genomes/contigs used for training, validation, and testing are detailed in Supplementary Table S3.

### **Development of GutEuk and training**

We developed GutEuk, a two-stage classifier designed to differentiate prokaryotic and eukaryotic (fungal and protozoal) sequences in the first stage and then further differentiate fungal and protozoal sequences in the second stage. GutEuk employs an ensemble model comprising a feedforward neural network (FNN) and a convolutional neural network (CNN) in each stage (Fig. 1A) to classify individual input sequences after they were fragmented into 5kb length. For each input sequence, its 4-, 5-, and 6-mer frequencies were calculated, and its nucleotide bases (A, C, G, T) were one-hot encoded as [1, 0, 0, 0], [0, 1, 0, 0], [0, 0, 1, 0], and [0, 0, 0, 1], respectively). The *k*-mer frequencies and the one-hot encoded DNA sequences served as the input of FNN and CNN, respectively. In the first stage, the probability of a sequence being eukaryotic origin is calculated by FNN and CNN. The probabilities calculated by the two networks are then averaged. If the average probability of an input sequence exceeded 0.5, it was subjected to another round of FNN and CNN classification to be fungal or protozoal. Input sequences identified with a probability less than 0.5 were considered prokaryotic. For input sequences shorter than 5kb but longer than the user-defined minimal length, they were zero-padded to the right to reach 5kb, and their *k*-mer frequencies are calculated based on the original input sequences (Fig. 1B). For sequences longer than 5kb, the remainder fragments shorter than 5kb after the 5kb fragmentation were also zero-padded on the

right and processed as described above. The origin of the input sequences (before fragmentation) was determined based on the predicted origin of their fragmented sequences. A user-defined confidence level was pre-set at the first and second stages ( $pt_1$  and  $pt_2$ , respectively). If the percentage of fragments of the input sequence classified as either prokaryote or eukaryote was less than the threshold  $pt_1$ , it was classified as not available (NA). Otherwise, they were classified as either prokaryotic or eukaryotic based on the majority rule. Similarly, input sequences were further classified in the second stage if the percentage of their fragments classified as either fungal or protozoal exceeded the threshold  $pt_2$ .

The FNN architecture consists of four sequential layers with decreasing dimensionality, accepting an initial 5,376-dimensional input vector of  $k$ -mer frequencies. The first layer is a fully connected layer with 5,376 input units and 1,024 output units, followed by ReLU activation and a dropout rate of 0.5. The second layer has 1,024 input units and 256 output units, with similar activation and dropout. The third layer features 256 input units and 48 output units, again followed by ReLU and a dropout rate of 0.5. The final output layer has 48 input units and 2 output units. The CNN architecture begins with an input layer accepting a one-hot encoded DNA as input. The first convolutional layer has 1 input channel and produces 32 output channels with a kernel size of (6, 4), followed by a ReLU activation. The output is then flattened starting from dimension 2 for subsequent 1D convolutions. The second layer transforms 32 input channels into 64 output channels with a kernel size of 3, also followed by ReLU activation. This is followed by max pooling with a pool size of 3 and batch normalization for 64 channels. The architecture continues with two more convolutional layers that increase the output channels to 128 and then 256, each followed by ReLU activation. A second max pooling layer is applied with ceil mode enabled, along with batch normalization for 256 channels. The feature maps are processed through an adaptive average pooling layer, which is then flattened. The output from the convolutional layers feeds into fully connected layers, beginning with a layer that transforms 256 input features to 64 output features, followed by ReLU activation and dropout with a probability of 0.5. The second fully connected layer reduces this to 8 output features, again with ReLU activation and dropout. Finally, the output layer consists of 8 input features and 2 output features. The CNN and FNN models at each stage were trained separately in PyTorch (Paszke et

al., 2019) with the cross-entropy loss and AdamW optimizer applied. Hyperparameter optimization involved manually adjusting and training the models across various configurations until the validation loss no longer improved over ten consecutive epochs. The model with the best performance (in terms of validation loss) was used to predict sequence origins.

### **Benchmarking EukRep, Tiara and GutEuk**

To benchmark the performance of GutEuk against Tiara and EukRep, we filtered out the genomes originally used for training Tiara and EukRep in the testing set (2995, 93, and 43 genomes for prokaryotes, fungi, and protozoa, respectively, of diverse taxonomic origins). We first evaluated the performance of the three classifiers at the contig level. Because contigs vary substantially in length, the first 50 kb of those longer than 50 kb was chopped into fragments of 3, 4, 5, 8, 10, and 20 kb. Then, the performance of each tool was compared with the different lengths. For GutEuk, performance was also determined at different confidence levels. We further evaluated the performance of each tool at the genome level, with the criterion being the percentage of 5kb fragments of the genomes correctly classified. A recent compendium of 706 genomes of cultured human gut fungi, encompassing 206 species across 48 families (Yan et al., 2024), was downloaded from NCBI (project ID: PRJNA835516) and used to benchmark GutEuk's performance with human gut eukaryotes at a confidence level of 0.5.

### **Identifying and functionally annotating microbial eukaryotic sequences from rumen metagenomes with GutEuk**

We systematically analyzed 1,093 previously sequenced rumen metagenomes (Supplementary Table S2) to identify microbial eukaryotic sequences (Fig. 1C). Specifically, the raw reads downloaded from NCBI SRA were quality-filtered using fastp (Chen et al., 2018) and assembled (individually for each metagenome) with MEGAHIT (Li et al., 2015) (v.1.2.1) with the default setting. The quality-filtered reads were mapped to the resultant contigs with Bowtie 2 (Langmead & Salzberg, 2012), and the mapping results were used for binning with MetaBAT2 (Kang et al., 2019). Then, GutEuk was used to identify eukaryotic sequences from the resultant contigs. The bins greater than 3 Mb were also screened with

GutEuk with the parameter “-bin”. Based on the benchmarking result, we determined conservative thresholds to classify the contigs and bins. Specifically, in the first stage, bins were classified as either prokaryotes or eukaryotes if 80% of the individual contigs within a bin were consistently classified as such and if the contigs were longer than 20 kb. In the second stage, eukaryotic bins were classified as fungi or protozoa if more than 95% of the genomes were classified into the same category.

For functional annotations, the protein sequences from the previously sequenced polyadenylated-enriched rumen metatranscriptomics (Li et al., 2022) were first assembled using Plass (Steinegger et al., 2019). The resultant protein sequences were clustered at 50% amino acid identity with MMseqs2 (Steinegger & Soding, 2017) to compile a rumen eukaryote protein sequence database. UniProtKB/Swiss-Prot release 2024\_01 (The UniProt Consortium, 2023) was downloaded and used as the target database to analyze the sequences derived from ruminant hindgut metagenomes. Then, genes were predicted from the identified eukaryotic contigs and bins from the foregut and hindgut using MetaEuk (Levy Karin et al., 2020) with the clustered rumen eukaryotic protein sequences and UniProtKB/Swiss-Prot target database, respectively. The obtained protein sequences were functionally annotated using eggNOG-mapper v2 (Cantalapiedra et al., 2021). Peptidases were also annotated from the identified protein sequences using DIAMOND (Buchfink et al., 2015) with the e-value of  $1 \times 10^{-5}$  as the threshold against the MEROPS database (Rawlings et al., 2010). Signal peptides were identified from the protozoal peptidases using SignalP 6.0 (Teufel et al., 2022), and those with a signal peptide were considered secretory.

### **Phylogenomic analysis of recovered protozoa genomes**

The completeness of the identified protozoa genomes/bins was assessed with BUSCO v5.6.1 by identifying lineage-specific single copy marker genes with the parameter “-l alveolata\_odb10” applied. Only those with completeness exceeding 50% were subjected to further phylogenomic analysis. Briefly, the multiple sequence alignment of 30 BUSCO proteins of the identified protozoal genomes/bins and the 52 rumen SAGs (Li et al., 2022) were concatenated. Genomes/SAGs with less than 70% of these markers

were excluded to ensure robust phylogenomic analysis. Specifically, the amino acid sequences of individual BUSCO proteins of the collected genomes were aligned with MAFFT v7.505 (Katoh & Standley, 2013) and trimmed with trimAl v1.4 (Capella-Gutiérrez et al., 2009). The trimmed alignments of all the BUSCO proteins were then concatenated together with catfasta2phyml.pl (<https://github.com/nylander/catfasta2phyml>). The concatenated alignment was used to build a phylogenetic tree using IQ-TREE (Nguyen et al., 2015) with the parameters “-redo -bb 1000 -m MFP -mr E,I,G,I+G -mfreq FU -wbtl -nt AUTO” applied. The obtained tree was visualized with iTOL (Letunic & Bork, 2019).

## **Evaluation of the new protein databases by reanalyzing the proteomics profiles from a recent rumen proteomic study**

To benchmark the performance of the new protein databases in improving rumen eukaryotic protein identification and their applicability as a database resource for future analyses, we reanalyzed the metaproteomic data generated from both goat and cattle under two different dietary treatments using the same method described previously (Andersen et al., 2023). Briefly, FragPipe version 19 (Yu et al., 2023) was employed to analyze the raw data obtained from mass spectrometry (MS). The raw data were then analyzed with (i) the original sample-specific databases used in the study or (ii) the original databases supplemented with the newly established protozoal (no additional SAGs included) and fungal protein databases (described in the previous section) using MSFragger (Kong et al., 2017). To estimate false discovery rates (FDR), the databases were supplemented with contaminant protein entries such as human keratin, trypsin, and bovine serum albumin, alongside the reversed sequences of all protein entries. Variable modifications, including methionine oxidation and protein N-terminal acetylation, were accounted for, while carbamidomethylation of cysteine residues was held as a fixed modification. The choice of trypsin as the digestive enzyme allowed for a single missed cleavage, with matching tolerance levels set at 20 ppm for both MS and MS/MS. Filtering was executed to ensure a 1% or lower FDR

threshold. Quantitative analysis was conducted using IonQuant (Yu et al., 2021). Only the proteins identified in the majority of the replicates in at least one of the treatments were considered present, as described in the original study.

#### **Data access**

The established ruminant eukaryotic protein databases and identified eukaryotic bins have been submitted to <https://doi.org/10.6084/m9.figshare.c.7457755.v1>. GutEuk, along with the codes used for visualization, are available at [https://github.com/yan1365/rumen\\_eukaryotes](https://github.com/yan1365/rumen_eukaryotes), and in [Supplementary Code](#).

#### **Competing interest statement**

The authors declare that they have no competing interests.

#### **Acknowledgments**

This work is supported in part by the USDA National Institute of Food and Agriculture (Award number: 2021-67015-33393). We also thank the Ohio Supercomputer Center for providing the computing resources.

#### **Author contributions**

M. Y. conceived this work, built the package, and performed bioinformatics analysis. T. A. performed the proteomics analysis. M. Y. and Z. Y. wrote the manuscript. All authors revised and approved the final manuscript.

## References


- Alexander, H., Hu, S. K., Krinos, A. I., Pachiadaki, M., Tully, B. J., Neely, C. J., & Reiter, T. (2023). Eukaryotic genomes from a global metagenomic data set illuminate trophic modes and biogeography of ocean plankton. *MBio*, e01676-01623.
- Andersen, T. O., Altshuler, I., Vera-Ponce de León, A., Walter, J. M., McGovern, E., Keogh, K., Martin, C., Bernard, L., Morgavi, D. P., & Park, T. (2023). Metabolic influence of core ciliates within the rumen microbiome. *ISME J*, 17(7), 1128-1140.
- Bálint, B., Merényi, Z., Hegedüs, B., Grigoriev, I. V., Hou, Z., Földi, C., & Nagy, L. G. (2024). ContScout: sensitive detection and removal of contamination from annotated genomes. *Nat Commun*, 15(1), 936.
- Bergman, E. (1990). Energy contributions of volatile fatty acids from the gastrointestinal tract in various species. *Physiol Rev*, 70(2), 567-590.
- Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND. *Nat methods*, 12(1), 59-60.
- Camargo, A. P., Roux, S., Schulz, F., Babinski, M., Xu, Y., Hu, B., Chain, P. S., Nayfach, S., & Kyrpides, N. C. (2023). Identification of mobile genetic elements with geNomad. *Nat Biotechnol*, 1-10.
- Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., & Huerta-Cepas, J. (2021). eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Mol Biol Evol*, 38(12), 5825-5829.
- Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*, 25(15), 1972-1973.
- Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics*, 34(17), i884-i890.
- Cho, C. H., Park, S. I., Huang, T.-Y., Lee, Y., Ciniglia, C., Yadavalli, H. C., Yang, S. W., Bhattacharya, D., & Yoon, H. S. (2023). Genome-wide signatures of adaptation to extreme environments in red algae. *Nat Commun*, 14(1), 10.
- Clemmons, B. A., Shin, S. B., Smith, T. P., Embree, M. M., Voy, B. H., Schneider, L. G., Donohoe, D. R., McLean, K. J., & Myer, P. R. (2021). Ruminant protozoal populations of angus steers differing in feed efficiency. *Animals*, 11(6), 1561.
- The UniProt Consortium (2023). UniProt: the universal protein knowledgebase in 2023. *Nucleic Acids Res*, 51(D1), D523-D531.
- Duncan, A., Barry, K., Daum, C., Elae-Fadros, E., Roux, S., Schmidt, K., Tringe, S. G., Valentin, K. U., Varghese, N., & Salamov, A. (2022). Metagenome-assembled genomes of phytoplankton microbiomes from the Arctic and Atlantic Oceans. *Microbiome*, 10(1), 67.
- Edwards, J. E., Forster, R. J., Callaghan, T. M., Dollhofer, V., Dagar, S. S., Cheng, Y., Chang, J., Kittelmann, S., Fliegerova, K., & Puniya, A. K. (2017). PCR and omics based techniques to study the diversity, ecology and biology of anaerobic fungi: insights, challenges and opportunities. *Front Microbiol*, 8, 1657.
- Firkins, J. L., Yu, Z., Park, T., & Plank, J. E. (2020). Extending Burk Dehority's Perspectives on the Role of Ciliate Protozoa in the Rumen. *Front Microbiol*, 11, 123.
- Fliegerova, K., Kaerger, K., Kirk, P., & Voigt, K. (2015). Rumen fungi. *Rumen microbiology: from evolution to revolution*, 97-112.
- Fu, Y., Zhang, K., Yang, M., Li, X., Chen, Y., Li, J., Xu, H., Dhakal, P., & Zhang, L. (2023). Metagenomic analysis reveals the relationship between intestinal protozoan parasites and the intestinal microecological balance in calves. *Parasit Vectors*, 16(1), 257.
- Gerber, P. J., Henderson, B., & Makkar, H. P. (2013). *Mitigation of greenhouse gas emissions in livestock production: a review of technical options for non-CO2 emissions*. Food and Agriculture Organization of the United Nations (FAO).
- Gerrick, E. R., Zlitni, S., West, P. T., Carter, M. M., Mechler, C. M., Olm, M. R., Caffrey, E. B., Li, J. A., Higginbottom, S. K., Severyn, C. J., Kracke, F., Spormann, A. M., Sonnenburg, J. L., Bhatt, A. S.,

- & Howitt, M. R. (2023). Metabolic diversity in commensal protists regulates intestinal immunity and trans-kingdom competition. *Cell*.
- Guyader, J., Eugène, M., Nozière, P., Morgavi, D. P., Doreau, M., & Martin, C. (2014). Influence of rumen protozoa on methane emission in ruminants: a meta-analysis approach 1. *Animal*, 8(11), 1816-1825.
- Hagen, L. H., Brooke, C. G., Shaw, C. A., Norbeck, A. D., Piao, H., Arntzen, M. Ø., Olson, H. M., Copeland, A., Isern, N., & Shukla, A. (2020). Proteome specialization of anaerobic fungi during ruminal degradation of recalcitrant plant fiber. *ISME J*, 15(2), 421-434.
- Hartinger, T., Gresner, N., & Südekum, K.-H. (2018). Does intra-ruminal nitrogen recycling waste valuable resources? A review of major players and their manipulation. *J Anim Sci Biotechnol*, 9(1), 33.
- Hoffmann, C., Dollive, S., Grunberg, S., Chen, J., Li, H., Wu, G. D., Lewis, J. D., & Bushman, F. D. (2013). Archaea and fungi of the human gut microbiome: correlations with diet and bacterial residents. *PLoS One*, 8(6), e66019.
- Hou, S., Tang, T., Cheng, S., Liu, Y., Xia, T., Chen, T., ... & Sun, F. (2024). DeepMicroClass sorts metagenomic contigs into prokaryotes, eukaryotes and viruses. *NAR genom. bioinform*, 6(2), lqae044.
- Huffnagle, G. B., & Noverr, M. C. (2013). The emerging world of the fungal microbiome. *Trends in microbiol*, 21(7), 334-341.
- Huws, S. A., Creevey, C. J., Oyama, L. B., Mizrahi, I., Denman, S. E., Popova, M., Muñoz-Tamayo, R., Forano, E., Waters, S. M., & Hess, M. (2018). Addressing global ruminant agricultural challenges through understanding the rumen microbiome: past, present, and future. *Front Microbiol*, 9, 2161.
- Jauhal, A. A., & Newcomb, R. D. (2021). Assessing genome assembly quality prior to downstream analysis: N50 versus BUSCO. *Mol. Ecol. Resour.*, 21(5), 1416-1421.
- Jin, W., Cheng, Y.-F., Mao, S.-Y., & Zhu, W.-Y. (2011). Isolation of natural cultures of anaerobic fungi and indigenously associated methanogens from herbivores and their bioconversion of lignocellulosic materials to methane. *Bioresour Technol*, 102(17), 7925-7931.
- Kang, D. D., Li, F., Kirton, E., Thomas, A., Egan, R., An, H., & Wang, Z. (2019). MetaBAT 2: an adaptive binning algorithm for robust and efficient genome reconstruction from metagenome assemblies. *PeerJ*, 7, e7359.
- Karlicki, M., Antonowicz, S., & Karnkowska, A. (2022). Tiara: deep learning-based classification system for eukaryotic sequences. *Bioinformatics*, 38(2), 344-350.
- Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*, 30(4), 772-780.
- Kavagutti, V. S., Bulzu, P.-A., Chiriac, C. M., Salcher, M. M., Mukherjee, I., Shabarova, T., Grujčić, V., Mehrshad, M., Kasalický, V., & Andrei, A.-S. (2023). High-resolution metagenomic reconstruction of the freshwater spring bloom. *Microbiome*, 11(1), 15.
- Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D., & Nesvizhskii, A. I. (2017). MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry-based proteomics. *Nat methods*, 14(5), 513-520.
- Laforest-Lapointe, I., & Arrieta, M.-C. (2018). Microbial eukaryotes: a missing link in gut microbiome studies. *mSystems*, 3(2), e00201-00217.
- Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat methods*, 9(4), 357.
- Letunic, I., & Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res*, 47(W1), W256-W259.
- Levy Karin, E., Mirdita, M., & Söding, J. (2020). MetaEuk—sensitive, high-throughput gene discovery, and annotation for large-scale eukaryotic metagenomics. *Microbiome*, 8, 1-15.
- Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics*, 31(10), 1674-1676.


- Li, Z., Wang, X., Zhang, Y., Yu, Z., Zhang, T., Dai, X., Pan, X., Jing, R., Yan, Y., Liu, Y., Gao, S., Li, F., Huang, Y., Tian, J., Yao, J., Xing, X., Shi, T., Ning, J., Yao, B., . . . Jiang, Y. (2022). Genomic insights into the phylogeny and biomass-degrading enzymes of rumen ciliates. *ISME J*, 16(12), 2775-2787.
- Lind, A. L., & Pollard, K. S. (2021). Accurate and sensitive detection of microbial eukaryotes from whole metagenome shotgun sequencing. *Microbiome*, 9, 1-18.
- Nguyen, L.-T., Schmidt, H. A., Von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*, 32(1), 268-274.
- Parfrey, L. W., Walters, W. A., Lauber, C. L., Clemente, J. C., Berg-Lyons, D., Teiling, C., Kodira, C., Mohiuddin, M., Brunelle, J., & Driscoll, M. (2014). Communities of microbial eukaryotes in the mammalian gut within the context of environmental eukaryotic diversity. *Front Microbiol*, 5, 298.
- Park, T., Mao, H., & Yu, Z. (2019). Inhibition of Rumen Protozoa by Specific Inhibitors of Lysozyme and Peptidases in vitro. *Front Microbiol*, 10, 2822.
- Park, T., Wijeratne, S., Meulia, T., Firkins, J. L., & Yu, Z. (2021). The macronuclear genome of anaerobic ciliate *Entodinium caudatum* reveals its biological features adapted to the distinct rumen environment. *Genomics*, 113(3), 1416-1427.
- Park, T., Yang, C., & Yu, Z. (2019). Specific inhibitors of lysozyme and peptidases inhibit the growth of the rumen protozoan *Entodinium caudatum* without decreasing feed digestion or fermentation in vitro. *J. Appl. Microbiol*, 127(3), 670-682.
- Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., Killeen, T., Lin, Z., Gimelshein, N., & Antiga, L. (2019). Pytorch: An imperative style, high-performance deep learning library. *Advances in neural information processing systems*, 32.
- Patra, A. K., & Yu, Z. (2022). Genomic Insights into the Distribution of Peptidases and Proteolytic Capacity among *Prevotella* and *Paraprevotella* Species. *Microbiol Spectr*, 10(2), e02185-02121.
- Peng, X., Wilken, S. E., Lankiewicz, T. S., Gilmore, S. P., Brown, J. L., Henske, J. K., Swift, C. L., Salamov, A., Barry, K., & Grigoriev, I. V. (2021). Genomic and functional analyses of fungal and bacterial consortia that enable lignocellulose breakdown in goat gut microbiomes. *Nat Microbiol*, 6(4), 499-511.
- Pronk, L. J., & Medema, M. H. (2022). Whokaryote: distinguishing eukaryotic and prokaryotic contigs in metagenomes based on gene structure. *Microb Genom*, 8(5).
- Ramayo-Caldas, Y., Prenafeta-Boldú, F., Zingaretti, L. M., Gonzalez-Rodriguez, O., Dalmau, A., Quintanilla, R., & Ballester, M. (2020). Gut eukaryotic communities in pigs: diversity, composition and host genetics contribution. *Anim. Microbiome*, 2, 1-12.
- Rawlings, N. D., Barrett, A. J., & Bateman, A. (2010). MEROPS: the peptidase database. *Nucleic Acids Res*, 38(suppl\_1), D227-D233.
- Saary, P., Mitchell, A. L., & Finn, R. D. (2020). Estimating the quality of eukaryotic genomes recovered from metagenomic analysis with EukCC. *Genome Biol*, 21(1), 1-21.
- Saraiva, J. P., Bartholomäus, A., Toscan, R. B., Baldrian, P., & Nunes da Rocha, U. (2023). Recovery of 197 eukaryotic bins reveals major challenges for eukaryote genome reconstruction from terrestrial metagenomes. *Mol. Ecol. Resour*, 23(5), 1066-1076.
- Seong, H. J., Kim, J. J., & Sul, W. J. (2023). ACR: metagenome-assembled prokaryotic and eukaryotic genome refinement tool. *Brief Bioinform*, 24(6), bbad381.
- Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, 31(19), 3210-3212.
- Steinegger, M., Mirdita, M., & Söding, J. (2019). Protein-level assembly increases protein sequence recovery from metagenomic samples manyfold. *Nat methods*, 16(7), 603-606.
- Steinegger, M., & Söding, J. (2017). MMseqs2 enables sensitive protein sequence searching for the analysis of massive data sets. *Nat Biotechnol*, 35(11), 1026-1028.

- Swift, C. L., Louie, K. B., Bowen, B. P., Olson, H. M., Purvine, S. O., Salamov, A., Mondo, S. J., Solomon, K. V., Wright, A. T., & Northen, T. R. (2021). Anaerobic gut fungi are an untapped reservoir of natural products. *Proc Natl Acad Sci*, *118*(18), e2019855118.
- Teufel, F., Almagro Armenteros, J. J., Johansen, A. R., Gíslason, M. H., Pihl, S. I., Tsirigos, K. D., Winther, O., Brunak, S., von Heijne, G., & Nielsen, H. (2022). SignalP 6.0 predicts all five types of signal peptides using protein language models. *Nat Biotechnol*, *40*(7), 1023-1025.
- Wallace, R. J., Onodera, R., & Cotta, M. A. (1997). Metabolism of nitrogen-containing compounds. In *The rumen microbial ecosystem* (pp. 283-328). Springer.
- West, P. T., Probst, A. J., Grigoriev, I. V., Thomas, B. C., & Banfield, J. F. (2018). Genome-reconstruction for eukaryotes from complex natural microbial communities. *Genome Res*, *28*(4), 569-580.
- Yan, Q., Li, S., Yan, Q., Huo, X., Wang, C., Wang, X., Sun, Y., Zhao, W., Yu, Z., & Zhang, Y. (2024). A genomic compendium of cultivated human gut fungi characterizes the gut mycobiome and its relevance to common diseases. *Cell*, *187*(12), 2969-2989. e2924.
- Yu, F., Haynes, S. E., & Nesvizhskii, A. I. (2021). IonQuant enables accurate and sensitive label-free quantification with FDR-controlled match-between-runs. *Mol. Cell Proteomics*, *20*.
- Yu, F., Teo, G. C., Kong, A. T., Fröhlich, K., Li, G. X., Demichev, V., & Nesvizhskii, A. I. (2023). Analysis of DIA proteomics data using MSFragger-DIA and FragPipe computational platform. *Nat Commun*, *14*(1), 4154.
- Yu, Z., Yan, M., & Somasundaram, S. (2024). Rumen protozoa and viruses: the predators within and their functions—A mini-review. *JDS Commun*. *5*, 236–240 (2024).


A





B




C







A




B




C




D






A


Foregut protozoa protein DB  
(5,265,119 protein clusters)




Foregut fungi protein DB  
(7,293 protein clusters)




Hindgut eukaryotic protein DB  
(9,405 protein clusters)



B



C



D

