



Discovery of a Novel, Paternally Expressed Ubiquitin-specific Processing Protease Gene through Comparative Analysis of an Imprinted Region of Mouse Chromosome 7 and Human Chromosome 19q13.4

Joomyeong Kim, Vladimir N. Noskov, Xiaochen Lu, et al.

Genome Res. 2000 10: 1138-1147

Access the most recent version at doi:[10.1101/gr.10.8.1138](https://doi.org/10.1101/gr.10.8.1138)

References

This article cites 36 articles, 8 of which can be accessed free at:
<http://genome.cshlp.org/content/10/8/1138.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Cold Spring Harbor Laboratory Press

Discovery of a Novel, Paternally Expressed Ubiquitin-specific Processing Protease Gene through Comparative Analysis of an Imprinted Region of Mouse Chromosome 7 and Human Chromosome 19q13.4

Joomyeong Kim,^{1,2} Vladimir N. Noskov,³ Xiaochen Lu,¹ Anne Bergmann,¹ Xiaojia Ren,¹ Tiffany Warth,¹ Paul Richardson,² Natalya Kouprina,³ and Lisa Stubbs^{1,2,4}

¹Human Genome Center, Biology and Biotechnology Research Program, L-441, Lawrence Livermore National Laboratory, Livermore, California 94551 USA; ²DOE Joint Genome Institute, Walnut Creek, California 94598 USA; ³Laboratory of Molecular Genetics, National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina 27709 USA

Using mouse BAC clones spanning an imprinted interval of proximal mouse chromosome 7 and the genomic sequence of the related interval of human chromosome 19q13.4, we have identified a novel mouse gene, *Usp29* (ubiquitin-specific processing protease 29), near two known imprinted genes, *Peg3* and *Zim1*. Gene *Usp29* is located directly adjacent to *Peg3* in a “head-to-head” orientation, and comprises exons distributed over a genomic distance of at least 400 kb. A similar human gene is also found in the homologous location in human chromosome 19q13.4. The mouse *Usp29* gene is also imprinted and is transcribed mainly from the paternal allele with highest expression levels in adult brain, especially in the cerebral cortex and hippocampus, and in the forebrain, face, and limb buds of midgestation mouse embryos. Analysis of a full-length 7.6-kb cDNA clone revealed that *Usp29* encodes an 869-amino-acid protein that displays significant homology with yeast and nematode ubiquitin carboxyl-terminal hydrolases. These data suggest that, like the candidate Angelman syndrome gene *Ube3a* (ubiquitin ligase), *Usp29* may represent another imprinted gene involved in the ubiquitination pathway. This identification of a third imprinted gene, *Usp29*, from the *Peg3/Zim1*-region confirms the presence of a conserved imprinted domain spanning at least 500 kb in the proximal portion of mouse chromosome 7 (Mmu7).

[The sequence data described in this paper have been submitted to the GenBank data library under accession nos. AF229257 and AF229438.]

Although the two alleles of most mammalian autosomal genes are functionally equivalent, the maternal and paternal copies of a subset of genes are distinguished by genomic imprinting, a mechanism by which one allele is epigenetically modified and repressed, depending upon parental origin. The epigenetic modification that determines imprinting is yet to be defined, but several lines of evidence strongly suggest that differences in DNA methylation and chromatin structure may be part of the molecular mechanism (Tilghman 1999). About 30 different imprinted genes have been isolated from human and mouse DNA, and more imprinted genes are predicted to be present in mammalian genomes (Barlow 1997; Morrison and Reeve 1998). The imprinted genes that have been described to date are clustered in discrete chromosomal

regions, and the clustering of those genes is conserved in mammals (Barlow 1995; Bartolomei and Tilghman 1997). These data have been interpreted to imply that genome imprinting is a long-range mechanism that controls allele-specific transcription of multiple genes within a defined chromosomal region (Nicholls 1994; Dittrich et al. 1996; Tilghman et al. 1998, 1999).

Imprinted genes exhibit several unique features in addition to parent-of-origin-specific monoallelic expression and differential methylation. The known imprinted genes generally are expressed at early stages of development, and many encode proteins with functions related to embryonic growth (Tilghman 1999). Recent studies also suggest the involvement of imprinted genes in controlling the parental caring behavior of mammals (Lefebvre et al. 1998; Li et al. 1999). Several imprinted genes such as *UBE3A* (Rougeulle et al. 1998), *IGF2* (Moore et al. 1997), *ZNF127* (Jong et al.

⁴Corresponding author.
E-MAIL: stubbs5@llnl.gov; FAX (925) 422-2282.

1999), and *IGF2R* (Wutz et al. 1997) display bidirectional transcription, producing both sense and antisense transcripts. A number of imprinted genes including *H19* and *IPW* are expressed without any coding capability; the final products of these genes are RNAs rather than proteins (Bartolomei and Tilghman 1997). Imprinted genes also tend to be composed of relatively small numbers of exons and to contain smaller introns than nonimprinted genes (Hurst et al. 1996), although there are some notable exceptions (e.g., *KvLQT1*; Lee et al. 1997).

Early mouse genetic studies with translocation mutant mice predicted the presence of nine different imprinted domains distributed onto seven chromosomes (Beechey and Cattanaach 1996). The chromosomal locations of most imprinted genes, in fact, are consistent with this initial prediction (Imprinting Map; <http://www.mgu.har.mrc.ac.uk/imprinting/imptables.html#impmaps>). Mouse chromosome 7 (*Mmu7*) was predicted in these early studies to contain three different imprinted domains, located in centromeric, central, and distal regions (Searle and Beechey 1990). The central and distal regions of *Mmu7* are syntenically homologous to intervals of human chromosome 15q13-11 and 11p15, respectively, that contain genes associated with two imprinting genetic disorders—Prader–Willi/Angelman (Nicholls et al. 1998) and Beckwith–Wiedemann syndromes (Reik and Maher 1997). Each of these domains contains several imprinted genes whose order, sequence, and imprinting status are highly conserved in humans and mice. Until recently, only one imprinted gene, *Peg3* (paternally expressed gene 3), had been localized to the centromeric imprinted domain of *Mmu7* (Kuroiwa et al. 1996) and the homologous region of human chromosome 19q13.4 (Kim et al. 1997a). We exploited the well-developed physical map surrounding human *PEG3* and the known conservation of this interval in humans and mice to identify a second mouse-imprinted gene, *Zim1* (imprinted zinc-finger gene 1), located downstream of *Peg3* (Kim et al. 1999). More recent studies identified a related and homologously positioned human zinc-finger gene, *ZIM2*, and showed that the gene shares seven 5'-exons with *PEG3*, a feature that distinguishes this pair of human genes from their closest murine counterparts (Kim et al. 2000). The significant divergence of *Zim1* and *ZIM2* gene organization and amino acid sequence suggests that the region surrounding *PEG3* has undergone significant changes during the course of mammalian evolution.

Through analysis of clone contigs spanning homologous segments of *Mmu7* and sequenced regions of human 19q13.4, we have isolated and characterized another novel mouse gene, *Usp29* (ubiquitin-specific processing protease 29) and have identified its human counterpart. Our analyses show that *Usp29* is also im-

printed and is expressed mainly from the paternal allele during embryogenesis and in adult brain. The sequence of a full-length 7.6-kb mouse cDNA clone and human coding sequences indicate that *Usp29* encodes a novel ubiquitin-specific processing protease, and together with *Ube3a* may represent a second imprinted gene involved in the ubiquitination pathway.

RESULTS

Isolation of Mouse *Usp29*

The recent isolation of the maternally expressed gene *Zim1* adjacent to *Peg3* (Kim et al. 1999), suggested the presence of a larger imprinted domain in the proximal *Mmu7* region. To identify additional genes from this conserved region, we analyzed the recently completed human 19q13.4 genomic sequence and information derived from large-insert mouse clones spanning the homologous *Mmu7* region. We constructed a physical map of a region spanning approximately 1 Mb around the *Peg3* gene from the 129/Sv inbred strain by isolating a series of BAC and PAC clones through chromosome walking; a portion of this physical map is shown in Figure 1. We were unable to clone the genomic region between *BAC247N19* and *PAC95D23* by chromosome walking because sequences from this region are underrepresented in existing 129/Sv commercial genomic libraries. To obtain genomic clones spanning this region, we utilized the transformation-associated recombination (TAR) technique (Larionov et al. 1996). One 220-kb clone (*tar17-21*) was successfully obtained. The position of *tar17-21* within the *Peg3* region was confirmed by verifying that sequences from each end of the TAR clone were present in BACs in the flanking clone contigs. Using low-pass sequencing data from the TAR clone and the surrounding BACs, we developed a series of overgo probes that permitted a complete, parallel set of overlapping BACs to be isolated from the C57BL/6 (RPCI-23) library (Fig. 1). Comparison between the two clone maps provided additional confirmation of the TAR clone's integrity and the length and organization of the 600-kb region.

Genomic subclones derived from the BAC and PAC clones identified several EST matches representing genes derived from this region. In particular, one of the end sequences of BAC402H23, 1.2 kb in length, identified two ESTs (GenBank accession nos. AA549132 and AI503662), indicating the presence of an unknown gene at this site. Database searches with these EST sequences indicated a significant level of homology with a hypothetical yeast protein, a putative ubiquitin carboxyl-terminal hydrolase (GenBank accession no. NP 009614). The novel mouse gene was named *Usp29* (ubiquitin-specific processing protease 29). Sequence analysis of the genomic region immediately upstream of mouse *Peg3* also identified one EST match (GenBank




Figure 1 Genomic organization of mouse and human *USP29*. (A) A physical map of a 600-kb region of proximal *Mmu7* shows the relative locations and transcription direction of *Zim1*, *Peg3*, and *Usp29*. The exons of *Usp29* are indicated by black boxes, and the coding region of mouse *Usp29* is indicated by a gray box. The genomic clones used for 129/Sv map construction include BAC (588F20, 247N19, 402H23), PAC (95D23), and TAR (17-21). A separate clone contig map using C57BL/6 genomic libraries (RPC1-23) is also shown with a subset of BAC clones. (B) The relative physical locations of human *PEG3*, *ZIM2*, and *USP29* are indicated by arrows.

accession no. W90868). The genomic sequence matching this EST is located about 1 kb upstream of the first exon of *Peg3*, and the transcriptional direction of the EST sequence is opposite to that of *Peg3*. These data suggested the presence of an unidentified gene with a transcription start located very near that of the *Peg3* transcription unit.

The two distinct groups of ESTs were initially thought to be derived from different genes because of the 400-kb genomic distance that lies between the end of BAC clone 402H23 and the 5'-end of *Peg3* (Fig. 1). Separate screens of a large-insert mouse cDNA library with probes derived from each EST set, however, identified identical sets of cDNA clones. Sequence analysis of these cDNA clones and RT-PCR experiments using primers from the two EST clusters (data not shown) demonstrated that sequences located immediately upstream of *Peg3* and the *Usp29* region, located 400 kb downstream, represent 5'- and 3'-portions of a single, large transcription unit. One clone isolated from a mouse cerebellum cDNA library, representing a full-length 7.6-kb transcript, was sequenced to provide the complete open reading frame and 5'- and 3'-UTR se-

quences of the mouse *Usp29* gene (GenBank accession no. AF229257).

Analysis of this cDNA sequence identified one open reading frame encoding a protein 869 amino acids in length (Fig. 2). Database searches with this predicted protein sequence retrieved a large number of protein entries, most of which display ubiquitin carboxyl-terminal hydrolase activity. Although database searches with the *Usp29* cDNA sequence derived a long list of similar proteins, the relatively low level of homology to known proteins suggests that *Usp29* is a new member of this mammalian protein family. In fact, the highest homology identified was found with the yeast hypothetical protein (GenBank accession no. NP 009614; 21% sequence identity over 550 amino acid residues), originally identified in BLAST searches with the end sequence of BAC402H23. Comparison of different ubiquitin carboxyl-terminal hydrolases identified two conserved domains that typify the type-2 ubiquitin carboxyl-terminal hydrolases, also known as ubiquitin-specific processing protease (Wilkinson 1997). These two domains, the cysteine box and the histidine box, are also present in the predicted sequence of *Usp29*,




Figure 2 Comparison of mouse and human *USP29* amino acid sequences. The upper sequence represents the mouse *Usp29* and the lower the human *USP29* amino acid sequence. Two conserved domains of the type-2 ubiquitin carboxyl-terminal hydrolase family are indicated by asterisks under the amino acid sequence. The consensus sequences of these two domains constructed from the multiple alignment of yeast sequences are also presented and compared with the sequence alignment of human and mouse *USP29*. Sequence identity (black background) and conservative substitution with similar properties (gray background) are indicated. The two sequences are aligned for maximal match by the SIM program (<http://dot.imgen.bcm.tmc.edu:9331/seq-search/alignment.html>).

indicating that *Usp29* belongs to the type-2 ubiquitin carboxyl-terminal hydrolase family (Fig. 2).

The exon structure of *Usp29* was determined through comparing the cDNA sequence with a draft sequence derived from the 129/Sv genomic clones covering the mouse *Peg3*-region (Table 1; GenBank accession no. AC020961 and AC020966). *Usp29* is composed of 5 exons distributed throughout an ~400-kb genomic region, with the 5'-end of *Usp29* positioned within 150 bp of the 5'-end of *Peg3* (Fig. 1). The first and second exons of *Usp29* share a low level of homology with the MER repeat family (Smit 1996). The distance between the first and second exons in mouse is estimated to be 7 kb, and the distance between exons 4 and 5 is 900 bp. The relative location of exon 3 is presently unknown because of uncertainty of spacing of contigs in the draft sequence. The entire ORF is located in exon 5, the last exon of mouse *Usp29* (Fig. 1).

Table 1. Position and Size of Exons and Introns of the Mouse *Usp29* Gene

Exon	Position	Length (bp)	Intron	Length (bp)
1	1–722	722	1	~7,000 ^a
2	723–943	221	2	unknown
3	944–1264	321	3	unknown
4	1265–1452	188	4	949 ^b
5	1453–7550	6098		

^aThe length was determined by PCR.

^bThe length was determined by DNA sequencing.

Isolation of Human *USP29*

To identify the human homolog of mouse *Usp29*, the 1.2-kb end sequence of 402H23 corresponding to portions of mouse *Usp29* coding sequence was hybridized to a Southern blot containing human and other mammalian genomic DNAs. At low stringency, this mouse probe detected positive fragments only from the genomic DNAs of closely related rodent species, such as mouse and rat, but not from human and other mammals (data not shown). These data provided the first suggestion that *Usp29* sequences might not be well conserved among mammals. Comparison of the cDNA sequence of mouse *Usp29* with the complete, finished genomic sequence of the region surrounding human *PEG3* (GenBank accession no. AC025588), however, enabled us to locate a single set of human sequences related to the mouse *Usp29* coding region, approximately 400 kb upstream of *PEG3* (Fig. 1). The genomic sequence containing part of the potential coding region of human *USP29* displays 65%–70% sequence identity at the nucleotide level with the mouse coding sequence. The human *USP29* sequence identified a perfect match with one human EST (GenBank accession no. AA909508), confirming the transcription of this sequence. With RT-PCR and RACE, about 3.2 kb of human *USP29* transcribed sequence was obtained. Analysis of this sequence identified a single open reading frame, encoding a predicted protein 922 amino acids in length (GenBank accession no. AF229438). Despite the relatively low level of overall sequence conservation between human and mouse predicted proteins

(42.5% amino acid sequence identity), the two conserved domains that are unique to the type-2 family, and which define the active sites of these enzymes, are well conserved between human and mouse *USP29* (Fig. 2). No other sequence with significant match to mouse *Usp29* coding sequence was found in the completed sequence of the human region.

The *USP29* cDNA sequence we obtained by RACE is derived from a contiguous 3.2-kb stretch of human genomic sequence, indicating that the coding region of human *USP29* is also located in a single exon. Alignment of the 5'-UTR of mouse *Usp29*, including exons 1–4 (1.3 kb), with the human genomic sequence also revealed the presence of sequences related to the first two exons of mouse *Usp29* immediately upstream of human *PEG3*. As in mouse, this conserved sequence identified two human EST matches (GenBank accession nos. H80201 and H79292), suggesting that this region is also transcribed. But repeated attempts using RT-PCR or RACE did not succeed in linking these upstream sequences to the human *USP29* ORF (data not shown). In fact, the human 5'-EST sequences and the *USP29* coding exon probe detected different transcripts in human Northern blots; in contrast, the two sets of related mouse sequences detected similar sets of transcripts (see below). These data suggested that the two clusters of cDNA sequences might represent different transcription units in the human genome.

Overlap between cDNA sequences we had isolated and a partial gene recently deposited sequence (GenBank accession no. AF124433) identified the 5'-end of human *USP29*. RT-PCR using human testis RNA as template confirmed the joining of this 5'-sequence and the *USP29* ORF. Assembly of these sequences permitted derivation of a complete 3.5-kb *USP29* transcript. Alignment of *USP29* cDNA and human genomic sequences identified three additional exons, and also localized the potential 5'-end of the human transcript to a position 9 kb upstream of the *USP29* ORF (Fig. 1).

Expression Pattern of Mouse *Usp29*

Northern Analyses

To examine the expression pattern of mouse *Usp29*, we hybridized two different cDNA probes, derived from the first exon and 3'-UTR of *Usp29*, respectively, to Northern blots containing poly(A)⁺ RNAs isolated from different adult tissues and embryos at various developmental stages. The highest levels of adult mouse *Usp29* expression were detected in brain. The size of the major transcript detected by both sets of probes was estimated to be about 7.5 kb, consistent with the length of the cDNA clone and sequence we obtained (Fig. 3A). Two minor transcripts, 4.5 and 5 kb in length, respectively, were also detected in brain, and several minor transcripts of different lengths were also detected in




Figure 3 Northern blot analyses of mouse and human *USP29*. (A) The expression pattern of mouse *Usp29* was analyzed using poly(A)⁺ RNAs derived from adult tissues and different-stage embryos. A 500-bp DNA fragment corresponding to the 3'-UTR of *Usp29* was used as a probe. A major transcript of mouse *Usp29*, 7.5 kb in length, and also two minor forms, 5.0 kb and 4.4 kb, were detected in adult brain as well as in embryos. (B) Human *USP29* transcripts were detected at a low level only in the adult testis (*upper*). The estimated size of the transcript is about 3.5 kb. The ESTs matching sequence located in the 5'-side of *PEG3* detected a 9-kb mRNA species only in ovary (*lower*).

testis with the 3'-UTR probe. In contrast, the probe derived from the first exon of mouse *Usp29* detected only the 7.5-kb major transcript in brain. Although the significance of these smaller transcripts cannot be established from present data, alternative splicing or alternative promoter usage might provide an explanation for their differential detection. The expression of *Usp29* was detected in RNA isolated from mouse embryos as early as 11 days postcoitum (dpc), and the consistent levels of *Usp29* expression appear to be maintained throughout the latter half of the gestation period (Fig. 3A).

In Situ Hybridization

The spatial pattern of *Usp29* transcription was analyzed by in situ hybridization of whole-mount midgestation mouse embryos and sectioned adult mouse brain. Both antisense and sense RNA probes of mouse *Usp29* were used for this experiment, in which only the antisense probe detected significant expression levels of mouse *Usp29*. At 10.5 dpc, *Usp29* is expressed at highest levels in the forebrain and limb buds, with significant levels of transcript also detected in the hindbrain and in developing structures of the face (Fig. 4A). In adult brain, *Usp29* transcripts were detected in many regions, but most notably in the cerebral cortex and hippocampus. These data are consistent with embryonic expression data, since the cerebral cortex and hippocampus are




Figure 4 Localization of *Usp29* transcripts in mouse embryo and adult brain. (A) Whole mount in situ hybridization of *Usp29* to 10.5-dpc embryo. *Usp29* expression (stained in dark blue) was detected at high levels in the face, forebrain (fb), hindbrain (hb), forelimb (fl), and hindlimb (hl). (B) Sagittal section of whole adult mouse brain hybridized with *Usp29* probe. Cells expressing the *Usp29* transcript are stained in red; the tissue has been counterstained with hematoxylin, which stains nuclei and therefore regions that are dense in neuronal cell bodies blue. Regions expressing *Usp29* at significant levels include cerebellum (cb), hippocampus (hc), habenula (hn), hypothalamus (hy), ependyma (ep), and cerebral cortex (cc). High-power magnification of selected *Usp29*-expressing regions, showing cell types expressing the gene. *Left panel (cb)*: Section of the cerebellum, showing high levels of *Usp29* expression in Purkinje cells (pc) but not in the densely packed granule cells (gc); *Center left panel (cc)*: section of the cerebral cortex, showing high levels of expression in pyramidal cells throughout the cerebral cortex; *Center right panel (ep)*: expression in ependymal cells; and *Right panel (hc)*: showing high levels of *Usp29* expression in the granule layer of the hippocampus.

derived from forebrain structures (Fig. 4B). Examination of sections under high-power magnification revealed that *Usp29* is transcribed in specific cell types within each region. For example, *Usp29* transcripts are present at very high levels in cerebellar Purkinje cells, but not in the underlying, densely packed granule cells (Fig. 4C, left panel). In the cerebral cortex, *Usp29* is expressed in the numerous pyramidal cells (Fig. 4C, center left panel). *Usp29* expression is also highly concentrated in ependymal cells (Fig. 4C, center right panel) and in the granule layer of the hippocampus (Fig. 4C, right panel).

Expression Pattern of Human *USP29*

The expression of human *USP29* was also examined with two cDNA probes derived from the 3'-UTR and coding region of the human gene. As shown in Figure 3B, significant amounts of transcript were detected only in adult testis with these two probes. Although Northern blots carrying poly(A)⁺ RNA derived from several different human fetal tissues were hybridized, we were unable to detect any significant levels of human *USP29* gene expression at these early developmental stages (data not shown). Because the mouse brain *Usp29* transcript initiates immediately upstream of *Peg3* and the related human upstream region is also transcribed, we hybridized probes derived from matching human EST sequences to a separate Northern blot. Unlike probes from related mouse regions, human "upstream" (adjacent to 5'-*PEG3*) and *USP29* coding sequence probes did not detect the same transcripts. The upstream probes detected a single 9-kb transcript,

which is expressed at high levels only in ovary (Fig. 3B). Taken together with the results of RT-PCR and RACE attempts and the different lengths and sites of expression of human and mouse *USP29* transcripts, these data suggest that human and mouse genes are transcribed differently.

Imprinting of Mouse *Usp29*

To determine the imprinting status of *Usp29*, we used hybrid animals derived from matings between *Mus musculus* and *Mus spretus* mice (Kim et al. 1999). Several sequence polymorphisms that differentiate alleles of each parental species were identified by comparing the sequences of *Usp29* cDNA segments derived from *M. musculus* and from *M. spretus*. Among several polymorphisms, two were selected and

used for the imprinting test: (1) a base substitution located in the first exon, and (2) a 19-bp insertion/deletion polymorphism located in the 3'-UTR of the gene. Experiments conducted with the two polymorphisms yielded identical results; in both cases, *Usp29* transcripts were derived primarily from the paternal allele in the hybrid mice. Data obtained using the insertion/deletion polymorphism located in the 3'-UTR are presented in Figure 5. To confirm these




Figure 5 Paternal expression of mouse *Usp29*. PCR amplification was performed across the 19-bp insertion/deletion polymorphism located in the 3' untranslated region of mouse *Usp29*. RNAs derived from the F1 hybrid (male *M. spretus* × female *M. musculus*) and the F2 hybrid (male *M. musculus* × female F1) were used for this imprinting test. Three genomic DNAs—*M. spretus* (lane 1), *M. musculus* (lane 2), and F1 hybrid (lane 3)—are also included in this test. Exclusive paternal expression of mouse *Usp29* is observed when using RNAs derived from embryos (lanes 4–6), neonatal (lane 7), and adult (lane 8) brains of F1 offspring, and also the neonatal brain (lane 9) of F2 animals.

results, a reciprocal analysis was performed with interspecific backcross progeny (the offspring of an interspecific hybrid female and a C3Hf male; Fig. 5, lane 9). The result of this test confirmed results obtained with F1 hybrid mice, demonstrating that *Usp29* transcripts are derived primarily from the paternal allele.

Analysis of *Usp29* transcripts derived from mouse embryos at different stages of gestation confirmed monoallelic expression. *Usp29* transcripts detected in 9.5-dpc and 14.5-dpc embryos are derived mainly from the paternal allele (Fig. 5, lanes 4–6). Since the major site of mouse *Usp29* expression is brain, we focused primarily on analysis of cDNA prepared from this tissue. *Usp29* transcripts in neonatal (7 day old) and adult (1 month old) mouse brains were also derived mainly from the paternal allele (Fig. 5, lanes 7 and 8), indicating that *Usp29* expression is stably imprinted throughout development.

DISCUSSION

Data presented here identify a new imprinted mouse gene, *Usp29*, located adjacent to mouse *Peg3* and describe the location, sequence, and expression patterns of the homologous human sequence. Mouse *Usp29* is expressed primarily from the paternal allele in the mouse brain from at least midgestation into adulthood; significant levels of embryonic expression were also detected in the developing limbs. The predicted protein is likely to function as a ubiquitin carboxyl-terminal hydrolase based on sequence similarity to other known genes of this class. Because it does not share significant amino acid sequence homology with identified mammalian proteins, *Usp29* appears to represent a novel member of this diverse family. Along with *Peg3* and *Zim1*, *Usp29* represents the third imprinted gene identified in a conserved 500-kb interval of proximal Mmu7. Our results indicate that this relatively unexplored imprinted domain may occupy a large genomic segment containing a number of imprinted genes, as has been observed for other known imprinted regions.

Although the actual biochemical role of *Usp29* is currently unknown, the sequence similarity of *Usp29* with members of the type-2 ubiquitin carboxyl-terminal hydrolase family suggest that *Usp29* might be a deubiquitinating enzyme (Wilkinson 1997). This raises the possibility that the gene might be involved in regulating the turnover rate of many proteins. As shown for *Ube3a*, an imprinted gene that is also thought to be involved in the ubiquitin pathway (Kishino et al. 1997), the absence of *Usp29* might be expected to be associated with pleiotropic effects on growth, development, and fitness with significant detriment to the animal. Although the biochemical functions of both proteins must be demonstrated formally, it is tempting to speculate that these two oppositely

imprinted genes—one functioning as a ligase and the other as a hydrolase of ubiquitin–protein complexes—might serve opposing functions that must be properly balanced for successful development of the mammalian nervous system. The high level of *Usp29* expression in mouse brain, a feature that is shared with *Ube3a*, suggests that the *Usp29* gene might play an especially significant role in mouse neurological development. However, although both mouse and human genes are expressed in testis, human *USP29* is not transcribed at significant levels in brain. Although Northern blots are notoriously insensitive, Northern data were supported by the fact that we could not amplify *USP29* coding region sequences from human brain cDNA templates by RT-PCR (data not shown). Low levels of human *USP29* transcription or expression in very limited brain regions cannot be ruled out at this point. But these data may also indicate that the mouse and human *USP29* genes have diverged significantly in biological function.

The different expression patterns of human and mouse *USP29* genes is not entirely unprecedented. For example, although human and mouse *PEG3* genes are both expressed in adult brain, the human gene is expressed at highest levels in ovary. By contrast, trace levels of mouse *Peg3* were detected in ovary only by sensitive methods such as RNase protection (Kim et al. 1997a). Significant differences in tissue-specific expression of orthologs have also been noted by other groups, including differences linked to the species-specific usage of different promoters (Wang et al. 1996; Kissinger and Raff 1998; Fougousse et al. 2000). Our data also suggest that exon structures of human and mouse *USP29* differ significantly. Although human and mouse *USP29* transcripts are both expressed in testis, human *USP29* mRNA is not present in significant levels in brain—the primary site of mouse expression. Our data indicate that the human and mouse major transcripts, at least, originate at significantly different genomic sites and most likely use different promoters. Whether these differences reflect real change in gene function is difficult to predict at this time, and must await further study.

Comparison of cDNA and genomic sequence shows that the 5'-ends of *Peg3* and *Usp29* genes are located less than 150 bp apart. It therefore seems likely that *Peg3* and *Usp29* share regulatory elements that control sites and levels of each gene's transcription, and possibly allele-specific expression. Consistent with this notion, the expression patterns of mouse *Usp29* and *Peg3* are very similar: both genes are expressed in embryos and at significant levels only in adult brain. Recent studies also indicate that the region surrounding the transcription start sites of the two genes is associated with a CpG-rich region that is differentially methylated in parent-of-origin-specific manner (Li et

al. 2000; J. Kim et al., unpub.). These observations support the idea that the transcription and imprinting of *Peg3* and *Usp29* might be jointly controlled through a small, shared genomic region. Further insights to the regulation and function of this newly defined imprinted domain should soon be forthcoming, as the boundaries of the domain, additional resident genes, and candidate regulatory elements are discovered through comparative sequence analysis in the upcoming year.

METHODS

Mouse Genomic Clone Isolation and Analysis

For the construction of a physical map of the *Peg3/Zim1*-region, several genomic clones were identified and characterized. The two BAC clones 588F20 and 247N19 were obtained through screening high-density mouse 129/Sv BAC library filters (Research Genetics, Huntsville, Alabama) with a pool of three genomic DNA fragments corresponding to the transcribed region of *Peg3* (Kuroiwa et al. 1996). The BAC clone 402H23 was isolated by screening the same library with two evolutionarily conserved probes (25670krab and ZNF134 finger probes; Kim et al. 1997a, 1999) derived from the coding region of human ZNFs located in H19q13.4. Because of the lack of clones covering the genomic region between the two groups of BAC clones described in the above, we utilized the TAR (Transformation-Associated Recombination) cloning procedure (Larionov et al. 1996) with one of the end sequences of BAC247N19 and a 200-bp B2 sequence as bites using 129/Sv genomic DNA. This technique derived one genomic clone, tar17-21. The PAC clone 95D23, connecting the tar 17-21 and BAC 402H23 clones, was obtained by screening mouse 129/Sv PAC library filters with two end clones of tar17-21 and 402H23. A separate screening of genomic clones covering the *Peg3/Zim1*-region was performed using C57BL/6 BAC library filters (RPC1-23 library, obtained from Roswell Park Cancer Institute, Buffalo, New York). About 30 BAC clones were obtained using six pooled overgo probes spanning the region (Fig. 1). The probes were designed and labeled as described previously (Ross et al. 1999). All BAC-end clones were isolated using a single-primer PCR approach (Kim et al. 1997b). The DNA samples of BAC, PAC, and TAR clones were prepared with the alkaline lysis protocol (Sambrook et al. 1989), digested with rare-cutting restriction enzymes including *NotI*, *EagI*, and *ClaI*, separated on pulse-field gels (Chef Mapper instrument, BioRad). DNAs were transferred to nylon membranes (Hybond, Amersham), and hybridized according to standard protocols (Stubbs et al. 1990).

cDNA Isolation and 5'- and 3'-RACE

Mouse *Usp29* cDNA clones were isolated from a cerebellum cDNA library (Doyle et al. 1997) with two probes corresponding to the first exon and the 3'-UTR of *Usp29*. Six of the 14 cDNA plaques positive with the two probes were randomly selected, plaque-purified, and analyzed further. To extend further the 5'-end of *Usp29*, we performed the RACE (Rapid Amplification of cDNA End) technique (Frohman et al. 1988). A cDNA template derived from adult brain RNA was used with the following two oligonucleotides: *Mim2* (5'-ATGG-GAAGGTGCGACATGAA-3') and *Mim5* (5'-CTACAACTTCG-GCAACGGGTT-3'). Human *USP29* cDNA clones were also ob-

tained through 5'- and 3'-RACE. A cDNA template derived from adult testis RNA was used with the following two sets of oligonucleotides: *HUSP29-1* (5'-ACACCTAGATCCCAGAACT-CAG-3') and *HUSP29-3* (5'-CCCTGCAAGATTAGAATGGTGC-3') for 3'-RACE, and *HUSP29-2* (5'-GATACTCCACACATTAC-CAGTG-3') and *HUSP29-4* (5'-CATGAGCACTAGAATCTC-CATC-3') for 5'-RACE. Amplified RACE products were separated on 0.8% agarose gel, and the major fragments in terms of mass were isolated from the gels using a gel extraction column (QIAquick gel extraction kit, QIAGEN). The fragments were subcloned into the TA cloning vector (TA cloning kit, Invitrogen).

Imprinting Tests

The imprinting (or monoallelic expression) of mouse *Usp29* was tested with hybrid offspring (F1) produced by crossing *M. musculus* (C3Hf) females with *M. spretus* male mice. The reciprocal imprinting test was performed with the offspring (F2) produced by backcrossing (*M. musculus* × *M. spretus*) hybrid females with C3Hf males. Embryos of 2 different stages (E9.5, E14.5) and tissues from 7-day-old and 1-month-old animals were collected from each cross. RNA was isolated using a commercially available kit (Rapid total RNA isolation kit; 5'-3', Inc, Boulder, CO). RNA samples were treated with RNase-free DNase I (Stratagene) for 30 min at 37°C, and 50 µg were used for the synthesis of cDNA (using the cDNA synthesis module, Amersham). The final volume of each reverse transcription reaction was 40 µl, and 1 µl of this material was taken for each PCR reaction. For imprinting tests of mouse *Usp29*, the 3'-UTR of mouse *Usp29* was amplified with the following two oligonucleotides: *mUsp29-7* (5'-TGCAAGGTCCTTAGCACATC-3') and *mUsp29-8* (5'-GTGCATAGCCTGCATCAGCA-3'). PCR amplification of the 3'-UTR of *Usp29* was performed using the following program in an MJ Research PTC-250 instrument: 95°C, 30 sec; 60°C, 1 min; 72°C, 1 min for 30 cycles, and 72°C, 5 min for 1 cycle. PCR reactions were carried out in a 50 µl reaction mixture containing 300 ng of each primer, 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, 1 mM dNTPs, 1% Triton X-100, and 1.25 U *Taq* DNA polymerase. Ten µl of each PCR reaction mixture were separated on 1.8% agarose gels to visualize the amplified PCR products.

Sequencing and Sequence Analysis

Subcloned genomic and cDNA fragments were sequenced from both directions using a fluorescence-based cycle-sequencing DNA sequencing kit (Dye terminator sequencing core kit, PE Applied Biosystems), and reactions were analyzed on an ABI 377 automated sequencer. Sequence alignments and database search were performed using BLAST (<http://www.ncbi.nlm.nih.gov/BLAST/>).

Northern Blot Analysis

A set of commercially available poly(A)⁺ RNA blots were used for expression analyses of human and mouse *USP29*: mouse multiple northern blot, mouse embryo blot, human multiple tissue northern blot, and human fetal northern blot (Clontech). For mouse *Usp29*, two probes representing different *Usp29* transcribed regions were used: the first exon and 3'-UTR of *Usp29*. For human *USP29*, two probes corresponding to the coding region and 3'-UTR of the gene, respectively, were used. For the human ESTs derived from the 5'-side of *PEG3*, a 1-kb DNA fragment isolated from the EST clone was used. Procedures and conditions for generating probes and for

performing hybridizations were as described previously (Stubbs et al. 1990).

Whole Mount and Section in Situ Hybridization

To generate an antisense and a sense riboprobe for the *Usp29* in situ hybridization, the transcribed region of mouse *Usp29* (Genbank accession no. AF229257, nt 2979-3638) was first amplified with the following two oligonucleotides: *mUsp29-1* (5'-CCAGGGTCTCATTGTTTCAT-3') and *mUsp29-2* (5'-TTGCAAGGGATGCCGGTAGA-3'). The initial PCR product was further amplified with the following two sets of oligonucleotides: *mUsp29-1* and *mUsp29-2-T7* (5'-TAAT-ACGACTCACTATAGGGTTGCAAGGGATGCCGGTAGA-3') for an antisense probe and *mUsp29-1-T7* (5'-TAAT-ACGACTCACTATAGGGCCAGGGTCTCATTGTTTCAT-3') and *mUsp29-2* for a sense probe. Reamplified PCR products were treated once with phenol/chloroform, washed with TE on a microcon-100 (Amicon), and concentrated to 1 µg/µl concentration. One µg of each template DNA was used for each in vitro transcription reaction with T7 polymerase. For whole mount in situ hybridizations, antisense and sense probes were labeled by digoxigenin-UTP with an RNA labeling kit (DIG RNA labeling kit, Boehringer Mannheim), anti-DIG-AP (alkaline phosphatase) from sheep, and BM Purple AP substrate (Boehringer Mannheim), which were used according to the manufacturers instructions. The hybridization experiments were performed according to a standard protocol (Hogan et al. 1994) with minimum modification. For section in situ hybridizations, brain tissues derived from 2-month-old mice were fixed by cardiac perfusion with 4% paraformaldehyde. Serial sections, in sagittal planes, were cut at 5 µm. The probes used for the section in situ hybridization were labeled with biotin-16-UTP (Boehringer Mannheim) and were detected by the DAKO GenPoint kit (DAKO). After color development of the probes, the sectioned tissues were also counterstained with hematoxylin according to the manufacturer's instructions (DAKO).

ACKNOWLEDGMENTS

We thank Dr. Elbert Branscomb and Mark Shannon for helpful discussion and critical reading of the manuscript. This work was performed under the auspices of the U.S. Department of Energy by LLNL under contract no. W-7405-ENG-48.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

REFERENCES

- Barlow, D.P. 1995. Gametic imprinting in mammals. *Science* **270**: 1610-1613.
- Barlow, D.P. 1997. Competition—A common motif for the imprinting mechanism? *EMBO J.* **16**: 6899-6905.
- Bartolomei, M.S. and Tilghman, S.M. 1997. Genomic imprinting in mammals. *Annu. Rev. Genet.* **31**: 493-525.
- Beechey, C.V. and Cattanach, B.M. 1996. Genetic imprinting map. *Mouse Genet.* **94**: 96-99.
- Dittrich, B., Buiting, K., Korn, B., Rickard, S., Buxton, J., Saitoh, S., Nicholls, R.D., Poustka, A., Winterpacht, A., Zabel, B. et al. 1996. Imprint switching on human chromosome 15 may involve alternative transcripts of the *SNRPN* gene. *Nature Genet.* **14**: 163-170.
- Doyle, J., Ren, X., Lennon, G., and Stubbs, L. 1997. Mutations in the *Cacn11a4* calcium channel gene are associated with seizures, cerebellar degeneration, and ataxia in tottering and leaner mutant mice. *Mamm. Genome* **8**: 113-120.
- Fougerousse, F., Bullen, P., Herasse, M., Lindsay, S., Richard, I., Wilson, D., Suel, L., Durand, M., Robson, S., Abitbol, M. et al. 2000. Human-mouse differences in the embryonic expression patterns of developmental control genes and disease genes. *Hum. Mol. Genet.* **9**: 165-173.
- Frohman, M.A., Dush, M.K., and Martin, G.R. 1988. Rapid production of full-length cDNAs from rare transcripts: Amplification using a single gene-specific oligonucleotide primer. *Proc. Natl. Acad. Sci. USA* **85**: 8998-9002.
- Hogan, B., Beddington, R., Costantini, F., and Lacy, E. 1994. *Manipulating the Mouse Embryo: A Laboratory Manual*. 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Hurst, L.D., McVean, G., and Moore, T. 1996. Imprinted genes have few and small introns. *Nature Genet.* **12**: 234-237.
- Jong, M.T.C., Gray, T.A., Ji, Y., Glenn, C.C., Saitoh, S., Driscoll, D.J., and Nicholls, R.D. 1999. A novel imprinted gene, encoding a RING zinc-finger protein, and overlapping antisense transcript in the Prader-Willi syndrome critical region. *Hum. Mol. Genet.* **8**: 783-793.
- Kim, J., Ashworth, L., Branscomb, E., and Stubbs, L. 1997a. The human homolog of a mouse-imprinted gene, *Peg3*, maps to a zinc finger gene-rich region of human chromosome 19q13.4. *Genome Res.* **7**: 532-540.
- Kim, J., Bergmann, A., and Stubbs, L. 2000. Exon sharing of a novel human zinc-finger gene, *ZIM2*, and paternally expressed gene 3 (*PEG3*). *Genomics* **64**: 114-118.
- Kim, J., Carver, E.A., and Stubbs, L. 1997b. Amplification and sequencing of end fragments from bacterial artificial chromosome clones by single-primer polymerase chain reaction. *Anal. Biochem.* **253**: 272-275.
- Kim, J., Lu, X., and Stubbs, L. 1999. *Zim1*, a maternally expressed mouse Kruppel-type zinc-finger gene located in proximal chromosome 7. *Hum. Mol. Genet.* **8**: 847-854.
- Kishino, T., Lalonde, M., and Wagstaff, J. 1997. *UBE3A/E6-AP* mutations cause Angelman syndrome. *Nature Genet.* **15**: 70-73.
- Kissinger, J.C. and Raff, R.A. 1998. Evolutionary changes in sites and timing of actin gene expression in embryos of the direct- and indirect-developing sea urchins, *Heliocidaris erythrogramma* and *H. tuberculata*. *Dev. Genes Evol.* **208**: 82-93.
- Kuroiwa, Y., Kaneko-Ishino, T., Kagitani, F., Kohda, T., Li, L.-L., Tada, M., Suzuki, R., Yokoyama, M., Shiroishi, T., Wakana, S. et al. 1996. *Peg3* imprinted gene on proximal chromosome 7 encodes for a zinc finger protein. *Nature Genet.* **12**: 186-190.
- Larionov, V., Kouprina, N., Graves, J., Chen, X.N., Korenberg, J.R., and Resnick, M.A. 1996. Specific cloning of human DNA as yeast artificial chromosomes by transformation-associated recombination. *Proc. Natl. Acad. Sci. USA* **93**: 491-496.
- Lee, M.P., Hu, R.J., Johnson, L.A., and Feinberg, A.P. 1997. Human *KVLQT1* gene shows tissue-specific imprinting and encompasses Beckwith-Wiedemann syndrome chromosomal rearrangements. *Nature Genet.* **15**: 181-185.
- Lefebvre, L., Viville, S., Barton, S.C., Ishino, F., Keverne, E.B., and Surani, M.A. 1998. Abnormal maternal behaviour and growth retardation associated with loss of the imprinted gene *Mest*. *Nature Genet.* **20**: 163-169.
- Li, L., Keverne, E.B., Aparicio, S.A., Ishino, F., Barton, S.C., and Surani, M.A. 1999. Regulation of maternal behavior and offspring growth by paternally expressed *Peg3*. *Science* **284**: 330-333.
- Li, L., Szeto, I.Y., Cattanach, B.M., Ishino, F., and Surani, A.M. 2000. Organization and Parent-of-origin-specific methylation of imprinted *peg3* gene on mouse proximal chromosome 7. *Genomics* **63**: 333-340.
- Moore, T., Constancia, M., Zubair, M., Bailleul, B., Feil, R., Sasaki, H., and Reik, W. 1997. Multiple imprinted sense and antisense transcripts, differential methylation and tandem repeats in a putative imprinting control region upstream of mouse *Igf2*. *Proc. Natl. Acad. Sci. USA* **94**: 12509-12514.
- Morrison, I.M. and Reeve, A.E. 1998. A catalogue of imprinted genes and parent-of-origin effects in humans and animals. *Hum. Mol. Genet.*

- Genet.* **7**: 1599–1609.
- Nicholls, R.D. 1994. New insights reveal complex mechanisms involved in genomic imprinting. *Am. J. Hum. Genet.* **54**: 733–740.
- Nicholls, R.D., Saitoh, S., and Horsthemke, B. 1998. Imprinting in Prader-Willi and Angelman syndromes. *Trends Genet.* **14**: 194–200.
- Reik, W. and Maher, E.R. 1997. Imprinting in clusters: Lessons from Beckwith-Wiedemann syndrome. *Trends Genet.* **13**: 330–334.
- Ross, M.T., LaBrie, S., McPherson, J., and Stanton, V.P., Jr. 1999. Screening large-insert libraries by hybridization. In *Current Protocols in Human Genetics* (ed. Ann Boyl). Wiley Press, New York.
- Rougeulle, C., Cardoso, C., Fontes, M., Colleaux, L., and Lalande, M. 1998. An imprinted antisense RNA overlaps *UBE3A* and a second maternally expressed transcript. *Nature Genet.* **19**: 15–16.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. 1989. *Molecular Cloning. A Laboratory Manual*. 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Searle, A.G. and Beechey, C.V. 1990. Genome imprinting phenomena on mouse chromosome 7. *Genet. Res.* **56**: 237–244.
- Smit, A.F. 1996. The origin of interspersed repeats in the human genome. *Curr. Opin. Genet. Dev.* **6**: 743–748.
- Stubbs, L., Huxley, C., Hogan, B., Evans, T., Fried, M., Duboule, D., and Lehrach, H. 1990. The *hox-5* and *surfeit* gene clusters are linked in the proximal portion of mouse chromosome 2. *Genomics* **6**: 645–650.
- Tilghman, S.M., Caspary, T., and Ingram, R.S. 1998. Competitive edge at the imprinted Prader-Willi/Angelman region? *Nature Genet.* **18**: 206–208.
- Tilghman, S.M. 1999. The sins of the fathers and mothers: Genomic imprinting in mammalian development. *Cell* **96**: 185–193.
- Wang, D., Marsh, J.L., and Ayala, F.J. 1996. Evolutionary changes in the expression pattern of a developmentally essential gene in three *Drosophila* species. *Proc. Natl. Acad. Sci. USA* **93**: 7103–7107.
- Wilkinson, K.D. 1997. Regulation of ubiquitin-dependent processes by deubiquitinating enzymes. *FASEB J.* **11**: 1245–1256.
- Wutz, A., Smrzka, O., Schweifer, N., Schellander, K., Wagner, E.F., and Barlow, D.P. 1997. Imprinted expression of the *Igf2r* gene depends on an intronic CpG island. *Nature* **389**: 745–749.

Received March 16, 2000; accepted in revised form June 2, 2000.