



## Gene Expression Changes Triggered by Exposure of *Haemophilus influenzae* to Novobiocin or Ciprofloxacin: Combined Transcription and Translation Analysis

Hans Gmuender, Karin Kuratli, Karin Di Padova, et al.

*Genome Res.* 2001 11: 28-42

Access the most recent version at doi:[10.1101/gr.157701](https://doi.org/10.1101/gr.157701)

---

**References** This article cites 44 articles, 15 of which can be accessed free at:  
<http://genome.cshlp.org/content/11/1/28.full.html#ref-list-1>

### License

**Email Alerting Service** Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).



---

To subscribe to *Genome Research* go to:  
<https://genome.cshlp.org/subscriptions>

---

Cold Spring Harbor Laboratory Press

# Gene Expression Changes Triggered by Exposure of *Haemophilus influenzae* to Novobiocin or Ciprofloxacin: Combined Transcription and Translation Analysis

Hans Gmuender,<sup>1</sup> Karin Kuratli, Karin Di Padova, Christopher P. Gray, Wolfgang Keck, and Stefan Evers

*F. Hoffmann-La Roche Ltd, Pharmaceuticals Division, CH-4070 Basel, Switzerland*

The responses of *Haemophilus influenzae* to DNA gyrase inhibitors were analyzed at the transcriptional and the translational level. High-density microarrays based on the genomic sequence were used to monitor the expression levels of >80% of the genes in this bacterium. In parallel the proteins were analyzed by two-dimensional electrophoresis. DNA gyrase inhibitors of two different functional classes were used. Novobiocin, as a representative of one class, inhibits the ATPase activity of the enzyme, thereby indirectly changing the degree of DNA supercoiling. Ciprofloxacin, a representative of the second class, obstructs supercoiling by inhibiting the DNA cleavage-resealing reaction. Our results clearly show that different responses can be observed. Treatment with the ATPase inhibitor Novobiocin changed the expression rates of many genes, reflecting the fact that the initiation of transcription for many genes is sensitive to DNA supercoiling. Ciprofloxacin mainly stimulated the expression of DNA repair systems as a response to the DNA damage caused by the stable ternary complexes. In addition, changed expression levels were also observed for some genes coding for proteins either annotated as “unknown function” or “hypothetical” or for proteins not directly involved in DNA topology or repair.

[The sequence data described in this paper have been submitted to the EMBL data library under accession nos. AJ297131 and AL135960.]

In the last few years, the total genomic sequences of many prokaryotes were determined. In parallel, techniques were developed that allow the monitoring of the expression levels of thousands of genes simultaneously. One of these techniques, developed by Affymetrix, is based on the principle of photolithography and conventional oligonucleotide synthesis, which allows the synthesis of short oligonucleotides in high-density arrays directly on a solid surface (Fodor et al. 1991, 1993; Chee et al. 1996; Lockhart et al. 1996; Wodicka et al. 1997). Total genomic DNA sequences are used to select sets of unique oligonucleotides to represent each open reading frame (ORF). To further increase the sensitivity and specificity of detection, a mismatch partner, which is identical except for a single base difference at the central position, is synthesized for each perfect-match oligonucleotide. These mismatch oligonucleotides serve as internal controls for the specificity of the probes.

For the gene expression analysis described here, a high-density microarray containing selected oligo-

nucleotides for ~2000 genes from the bacterium *Streptococcus pneumoniae* and for ~1800 genes from *Haemophilus influenzae* was used (Fleischmann et al. 1995). In addition, the microarray contains many control genes, sequence information from intergenic regions, and genes coding for ribosomal and transfer RNA. A set of 25-mer oligonucleotides for a specific gene usually includes 25 probe pairs (a probe pair consists of the perfect match and the corresponding mismatch oligonucleotide) and at least 20 probe pairs for very short genes. This microarray was used to simultaneously determine the changes in RNA levels for all the genes transcribed by *H. influenzae* following the addition of DNA gyrase inhibitors.

Regulation at the transcriptional level is only one possibility for a cell to respond to changing growth conditions. Other regulatory mechanisms act at the level of mRNA translation. Therefore, two-dimensional polyacrylamide gel electrophoresis (2D-PAGE), another tool for expression analysis that has been facilitated by the availability of whole-genome sequences and new developments in mass spectrometry, was performed in parallel. This allowed the comparison of the changes at the protein level with those of the transcriptional pattern.

<sup>1</sup>Corresponding author.

E-MAIL [hans.gmuender@roche.com](mailto:hans.gmuender@roche.com); FAX 41-61-688 27 29.

Article and publication are at [www.genome.org/cgi/doi/10.1101/gr.157701](http://www.genome.org/cgi/doi/10.1101/gr.157701).

DNA gyrase (E.C. 5.99.1.3.), a prokaryotic topoisomerase II enzyme essential for viability, consists of two subunits, A and B, the active enzyme being an A<sub>2</sub>B<sub>2</sub> tetrameric complex (for reviews, see Reece and Maxwell 1991; Menzel and Gellert 1994; Luttinger 1995; Roca 1995; Sharma and Mondragon 1995). The enzyme has no direct mammalian counterpart and is the only enzyme known to be able to introduce negative supercoils into DNA by using the energy derived from ATP hydrolysis. A key step in this supercoiling reaction is the DNA gyrase mediated cleavage of DNA. It has been shown that the class of subunit A inhibitors, the quinolones and the pyrimido[1,6-*a*]benzimidazoles, interrupt the cleavage and resealing cycle at the cleavage step (for reviews, see Hooper and Wolfson 1991; Hubschwerlen et al. 1992; Hooper 1993; Gmuender et al. 1995, 1997). However, there is evidence that quinolones acting *in vivo* have effects beyond the inhibition of DNA gyrase. They induce the formation of a stable ternary complex consisting of the enzyme, DNA, and the inhibitor, resulting in DNA damage, which in turn blocks replication and transcription (for reviews, see Drlica and Zhao 1997; Maxwell 1999). As a consequence, the expression of DNA repair systems, mainly the SOS system, is induced (Pidcock and Wise 1987; Walters et al. 1989; Pidcock et al. 1990). Another class of DNA gyrase inhibitors, the cyclothialidines and the coumarins, bind to the ATP binding site located in the subunit B, thereby inhibiting the supercoiling activity of the enzyme but leaving the DNA otherwise intact (Contreras and Maxwell 1992; Ali et al. 1993; Goetschi et al. 1993; Maxwell 1993; Nakada et al. 1994; Ali et al. 1995; Nakada et al. 1995; Gormley et al. 1996; Lewis et al. 1996; Oram et al. 1996; Tingey and Maxwell 1996; Kampranis et al. 1999). The initiation of transcription of many genes is sensitive to DNA supercoiling, often exhibiting an optimum with respect to the degree of supercoiling (Jovanovich and Lebowitz 1987; Steck et al. 1993; Wang and Lynch 1993).

The goals of this study were (1) to investigate if both classes of antibiotics, although inhibiting the same enzyme but through different mechanisms, induce different mechanism-related expression and translational patterns; (2) to cross-validate the two technologies; and (3) to test to what extent the use of both technologies in conjunction enhances the power of expression analysis.

## RESULTS AND DISCUSSION

### Methodological Studies

#### *Sensitivity*

The Affymetrix chip analysis detected transcripts for typically 70%–85% (in some experiments, even 90%) of the genes represented on the chips, indicating that

most of the genes are transcribed independently of the growth conditions and that the sensitivity of the system is high enough to detect even low abundance transcripts. Cross-hybridization with *S. pneumoniae* oligonucleotides present on the same chip was negligible in all experiments.

Two-dimensional gel electrophoresis followed by the detection of radioactive spots by using a Phosphorimager and the PDQuest program reproducibly detected 560 spots, which corresponds to ~30% of all theoretical gene products. Of these spots, 274 can be assigned to the corresponding gene by comparison with a 2D protein map (Langen et al. 2000). Fifty-one proteins are present as isoforms in more than one spot, so that the expression of 223 genes (12% of all genes) could be simultaneously quantified in 2D-gels.

#### *Reproducibility: Transcriptional Imaging*

To estimate the variation between expression patterns derived from RNA isolated from the same culture or from different cultures but grown under the same conditions, we hybridized reverse transcribed RNA from control cultures to the microarrays and normalized and analyzed the results by using the Affymetrix GeneChip software. RNA isolated from the same culture but reverse transcribed and hybridized independently shows a highly reproducible gene expression pattern and almost identical results after hybridization (Table 1; Fig. 1). About 90% of the ratios between the average differences of two normalized experiments lay between 0.66 and 1.5 with only a few transcripts being identified as increased or decreased. No significant changes, that is fold changes of the corresponding gene transcripts >2 or <-2, could be observed.

Samples of RNA from the same culture but isolated, reverse transcribed, and hybridized independently showed slightly more variation, but, again, little deviation from the ideal 1:1 ratio was observed. The transcripts identified as increased or decreased represented <1% of all genes present on the microarray (Table 1; Fig. 1). Ninety-three percent of the ratios between the average differences were between 0.66 and 1.5. These results show that the reproducibility of the procedures for RNA extraction, reverse transcription, labeling, and hybridization is very high and that detected variations are due to real differences in the mRNA concentrations in the cultures.

RNA samples from different cultures grown under the same conditions showed clearly more variability, even though the sample preparation procedure was kept as constant. For 68% of all genes, the ratios between the average differences were between 0.66 and 1.5. Because the reproducibility of the sample preparation and hybridization is very high, this apparent variability can only be due to small differences in the culture conditions that may have been introduced by us-

**Table 1.** Reproducibility of the mRNA Quantification

Sample preparation	Experiment	NF <sup>a</sup>	Increased or decreased <sup>b</sup>	Fold change >2 or <-2 <sup>c</sup>
Same culture, same RNA preparation, independent labeling and hybridization	1	0.98	1	0
	2	1.27	29	6 <sup>c</sup>
	3	0.66	7	0
Same culture, independent RNA preparation, labeling, and hybridization	1	0.88	32	7
	2	1.12	3	1
	3	1.34	13	1
	4	1.70	3	0
Different cultures		1.22	3.33	61

<sup>a</sup>(NF) Normalization factor.

<sup>b</sup>Transcripts called increased or decreased according to the Affymetrix GeneChip software.

<sup>c</sup>Fold changes between -2.1 and -2.5.

Genes total = 1961.

ing different batches of media, slightly different inoculum sizes, or different optical densities (ODs) at the time of cell collection. However, the 61 transcripts from two independently grown cultures that were called increased or decreased by the Affymetrix software and that showed a fold change of >2 or <-2 represented only about 3% of the genes present on the microarray.

#### Reproducibility: Protein Quantification


To estimate the gel-to-gel reproducibility, we resolved a protein extract from a control culture on two 2D gels in parallel and matched the resulting gel images. A sample from a second culture obtained under the same conditions was also analyzed and the gel image was compared to the parallel gels. Of the 560 spots detected, the calculated change factor was >2 or <-2 for 185 spots (32%) when gels from the same sample were compared and 220 (39%) when comparing different samples. A comparison for the 274 spots for which an identity has been assigned resulted in values of 68

(25%) for the comparison of two gels from the same sample and 140 (50%) for the comparison of two different samples. These figures indicate that the reproducibility for the annotated spots is higher, most likely reflecting the stronger intensity of these spots, which makes their quantification more accurate. The reproducibility of the 2D-gel-based protein quantification is clearly inferior to that observed for transcriptional imaging by using Affymetrix chips.

To take account of these findings, we chose to carry out five parallel experiments and to analyze two on Affymetrix chips and all five on 2D gels. Only parallel samples were compared to avoid misinterpretations resulting from culture-to-culture variability.

#### Quantitative Comparison between Total mRNA Levels and Protein Synthesis Levels

In previous studies comparing mRNA and protein levels, samples were not prepared in parallel so that only an approximation of the reproducibility of mRNA



**Figure 1** (A) The correlation between the hybridization results obtained from the same RNA but independently reverse transcribed and hybridized. (B) The correlation between the hybridization results obtained from the RNA isolated, reverse transcribed, and hybridized independently. (C) The correlation between the hybridization results obtained from the RNA isolated from different cultures grown under the same conditions. Averaged intensities from 100 to 10,000 units are plotted. (Solid line) the ideal 1:1 ratio; (dashed line) a factor of two; (broken line) a factor of three between the two measurements.

quantification and quantification of proteins could be made (Anderson and Seilhamer 1997; Gygi et al. 1999). One goal of this work was to be able to distinguish between technological and biological variability. Therefore, care was taken to prepare samples for transcription and 2D gel analysis in a strictly parallel fashion.

The intensity of the signals obtained on the Affymetrix chips was compared to that of the protein spots detected on 2D gels that could be assigned an identity. For this purpose, the spot quantities were normalized for their methionine content to make both values comparable. The correlation coefficient was calculated between both sets of values and was found to be 0.5. This means that the degree of correlation between the levels of mRNA and those of newly synthesized proteins was relatively low. However, neither technology is suited to measure absolute quantities and a bias may have been introduced at one of the following steps: (1) The features on the chips are not totally uniform for their annealing temperature and consequently for hybridization intensity; (2) cDNA synthesis could introduce a bias; and (3) proteins have different solubilities in the sample solution for IEF and the equilibration solution for the second dimension so that only a portion of the synthesized protein is detected on the gel.

#### *Comparison between Differential RNA and Protein Synthesis*

To estimate the degree of consistency between the results obtained by RNA and protein quantification, we calculated the average fold changes and compared for the Novobiocin treatment at the higher concentration. Genes were selected that showed significant changes by using either one of both detection methods (143 genes). The values were grouped into induced (fold change  $>2$ ), repressed (fold change  $<-2$ ), and unchanged genes/proteins. In 55% of all cases, the results fell into the same category. About 40% of all changes were detected by only one of the technologies. In most of these cases, however, this finding is due to the thresholds set for categorization, and the sign of the fold changes correlated in both techniques. In a small number of cases (3.5%), a clear discrepancy was found. In most of these cases, the proteins were represented in more than one spot and only one of these showed results conflicting with the mRNA measurements. The spots may therefore correspond to degradation products or to covalent reaction intermediates of enzymes. Only one protein, phosphoglycerate kinase (HI0525), was clearly detected as repressed by 2D-gel comparison and induced by transcriptional imaging.

Statistical analysis of the value pairs was performed and yielded correlation coefficients between 0.02 and 0.52 for the three time points, indicating a

very weak degree of relatedness between the obtained values. These observations suggest that, in *H. influenzae*, the gene expression changes observed on the mRNA and protein level may be qualitatively similar, but that the magnitude of the change detected differs significantly between the two technologies.

#### *Proteins Represented by Multiple Spots*

Forty-three proteins were represented by more than one spot on the 2D gels. These protein isoforms could reflect protein posttranslational modifications, degradation products, covalent reaction intermediates, or artifacts such as acrylamide adducts. Many of these proteins are enzymes known to form covalent intermediates with the respective reaction educt (glyceraldehyde-3-phosphate dehydrogenase, threonine synthase, malonyl transacylase, fructose-1,6-bisphosphate aldolase, transaldolase). Not surprisingly, in most of these cases the expression of only one of these spots followed the mRNA levels, whereas the intensity of the other one, presumably the covalent reaction intermediate, remained unchanged. Three proteins present as isoforms were tRNA synthetases (Asp, Gly, Lys), suggestive of a covalent reaction intermediate or of proteolytic cleavage. Other proteins have not been reported to undergo covalent modification or to act by an unknown mechanism (e.g., hslUV, aspartase, phosphoenolpyruvate carboxykinase). The study of these protein isoforms, although not performed in detail in this work, could yield important information about protein modifications that affect protein function.

#### **The Effect of DNA Gyrase Inhibitors on the Gene Expression Pattern**

We addressed the question of whether two different classes of DNA gyrase inhibitors would induce different responses at the level of transcription or of translation or of both. As described earlier, a key step in the DNA supercoiling reaction is the DNA gyrase-mediated double-stranded cleavage and resealing of DNA. It has been shown that the two classes of subunit A inhibitors, the quinolones and the pyrimido[1,6-*a*]benzimidazoles, interrupt this cleavage and resealing cycle at the cleavage step (Hooper and Wolfson 1991; Hubschwerlen et al. 1992; Hooper 1993; Gmuender et al. 1995). Two other classes of gyrase inhibitors, cyclothialidines and coumarins, bind to the ATP binding site located in the subunit B and inhibit the ATPase activity (Ali et al. 1993; Maxwell 1993; Ali et al. 1995; Nakada et al. 1995; Lewis et al. 1996; Oram et al. 1996). These compounds therefore inactivate the enzyme without introducing DNA strand breaks. Novobiocin (a coumarin) and Ciprofloxacin (a quinolone) were chosen as well-characterized DNA gyrase inhibitors representing these two functional classes. To elucidate the

concentration- and time-dependence of the response, we used low inhibitor concentrations, approximating the minimal inhibitory concentration (MIC, values derived from conventional agar plate techniques), and 10-fold higher concentrations. Cells were collected after 10, 30, and 60 min. Both inhibitors were added at an OD of 0.4 (mid log phase). Control cells grown from the same cultures in the absence of inhibitors were collected at the same time. RNA from treated and from control cells was isolated, reverse transcribed, and hybridized to microarrays. The expression patterns from the inhibitor treated cells were compared with the corresponding control cells after normalization by using the Affymetrix GeneChip software. At the time of harvesting, an aliquot was removed from the culture and the bacteria were pulse labeled with radioactive methionine. The protein extract from this culture was used for the quantification of protein synthesis rates by 2D-PAGE followed by computerized spot quantification.

#### Novobiocin

Novobiocin is not a bactericidal DNA gyrase inhibitor but indirectly influences the optimal supercoiling. The degree of supercoiling can influence promoter activity (Jovanovich and Lebowitz 1987; Steck et al. 1993; Wang and Lynch 1993). In addition, a nonspecific inhibitory activity on ATPases with similar binding cannot be excluded and may have an effect on gene expression. In fact, about 640 genes (37% of all genes) showed, at least at one concentration and time point, increased or decreased expression rates. At the lower concentration however, even after 60 min only a few genes showed a change in expression level. The expression pattern observed at this concentration was not significantly different between the three time points. The most prominent genes in this group, that is, those showing a clearly changed expression pattern at all time points, are listed in Table 2. All of these genes were also detected as increased or decreased at the higher Novobiocin concentration.

About 40 other genes also showed a changed ex-

pression rate, but whether these are primary or secondary effects of the addition of the inhibitor remains to be determined. It is very unlikely that these differences are artifacts derived from the culture conditions, because the cultures were grown under identical conditions apart from the addition of the inhibitor. The observation that all of these genes were also clearly called increased or decreased at the higher Novobiocin concentration confirms the significance of the findings.

At a higher Novobiocin concentration, ~140 genes were detected as changed according to the stringent criteria described earlier. They were ranked by the amplitude of the change and the most pronounced of these genes are shown in Table 3.

Not listed but noteworthy is that about 50 ORFs coding for hypothetical proteins also reproducibly show changed expression patterns.

The results demonstrate that with the low inhibitor concentration and after a short incubation period only a few genes showed a clear change in the expression pattern, namely, DNA gyrase subunit B and the ribosome releasing factor with an approximate twofold increase, topoisomerase I and a hypothetical protein with a twofold decrease. The ribosome releasing factor is responsible for the release of ribosomes from messenger RNA at the termination of protein biosynthesis (Kaji et al. 1998). The increased DNA gyrase expression and the decreased topoisomerase I expression can be interpreted as the attempt to maintain an optimal supercoiling degree to compensate for the enzyme inhibition. The hypothetical proteins for which a function is unknown are noteworthy. Their role is subject for detailed functional genomic studies.

At the higher inhibitor concentration many more genes showed an increased or decreased expression. They belong to various functional classes and their function appears unrelated to DNA topology. Possibly the higher concentration is too high to determine the primary effect on the gene expression and the observations are mostly secondary effects. However, the genes with a changed expression pattern after incubation with the lower concentration were also detected at

**Table 2.** Genes with Changed Expression Levels after Addition of 12 ng/mL of Novobiocin

No. <sup>a</sup>	Gene product	Fold change mRNA <sup>b</sup> (protein)		
		10 min	30 min	60 min
hi0567	DNA gyrase, subunit B	2.0	2.2	1.8
hi0808	Ribosome releasing factor	-2.4 (NC)	-2.1 (NC)	-2.0 (-3.1)
hi1365	Topoisomerase I	-1.9	-2.2	-1.8
hi1048	Hypothetical protein	-2.6	-3.2	-1.8
hi1507	Hypothetical protein	NC (-3.7)	NC (-2.6)	NC (-2.0)

<sup>a</sup>Numbering according to Fleischmann et al. (1995).

<sup>b</sup>Standard deviations were below 20%.

**Table 3.** Genes Showing Major Expression Changes after Addition of 125 ng/mL of Novobiocin

No. <sup>a</sup>	Gene product	Fold change mRNA (protein)		
		10 min	30 min	60 min
hi0944	Riboflavin biosynthesis protein	2.9	2.9	3.6
hi0873	UDP-N-acetylglucosamine epimerase	4.6	4.7	3.4
hi1722	Methionine aminopeptidase	3.5 (NC)	3.9 (NC)	3.4 (3.2)
hi1051	Leukotoxin secretion ATP-binding protein	3.1	3.9	3.3
hi1623	Mercury resistance regulatory protein	3.2	2.7	3.2
hi0991	DNA/ATP binding protein	2.5	2.7	3.1
hi0872	Undecaprenyl-phosphate galactosephosphotransferase	3.6	4.3	3.0
hi1305	Hydrogenase gene region	2.7	3.2	2.9
hi1573	Pyruvate kinase type II	2.3 (NC)	2.3 (NC)	2.7 (NC)
hi0394	Peptidyl-tRNA hydrolase	2.6	2.4	2.6
hi0533	RNA polymerase sigma-70 factor	2.4	2.1	2.6
hi1247	Excinuclease ABC subunit B	2.2	2.4	2.5
hi1304	N utilization substance protein B	2.3	2.9	2.5
hi1441	Stringent starvation protein A	2.3 (NC)	2.9 (2.2)	2.5 (NC)
hi0567	DNA gyrase, subunit B	2.1	2.9	2.4
hi0603	Uroporphyrinogen III methylase	2.7	2.1	2.4
hi1307	Virulence plasmid protein	2.4	2.5	2.3
hi0769	Cell division ATP-binding protein	3.0	2.5	2.3
hi0970	3-Dehydroquinase	2.3	2.2	2.2
hi1264	DNA gyrase, subunit A	2.5	2.1	2.2
hi0992	DNA polymerase III beta-subunit	2.2 (NC)	2.5 (NC)	2.2 (NC)
hi0969	O-Succinylbenzoate-CoA synthase	2.1	2.4	2.2
hi0623	Methionyl-tRNA formyltransferase	3.0	2.8	2.1
hi0625	TRK system potassium uptake protein	3.1	3.0	2.1
hi0236	ArsC protein	-2.4	-2.3	-2.1
hi1734	Enoyl-reductase <sup>b</sup>	-2.6 (-8.8)	-2.6 (-7.2)	-2.1 (-6.2)
hi0422	ATP-dependent RNA helicase	-2.8	-2.6	-2.1
hi0537	Urease accessory protein	-2.3	-4.0	-2.1
hi1661	Dihydrolipoamide succinyltransferase	-4.7 (NC)	-2.6 (NC)	-2.2 (NC)
hi0425	Phosphatidylserine synthase	-2.9	-2.1	-2.2
hi0863	Pyridoxamine phosphate oxidase	-3.0	-3.4	-2.2
hi1434.1	Cold shock-like protein	-3.0	-2.1	-2.2
hi1186	Dipeptide transport system permease protein	-2.5	-2.5	-2.2
hi1676	Molybdenum cofactor biosynthesis protein A	-3.9	-3.5	-2.2
hi1040	Restriction enzyme	-3.0	-3.4	-2.2
hi0503	High affinity ribose transport protein	-3.4	-2.9	-2.3
hi1152	PmbA protein homolog	-3.0	-2.2	-2.3
hi1193	Branched-chain-amino-acid transaminase	-2.3 (NC)	-2.5 (-1.5)	-2.3 (NC)
hi0538	Urease accessory protein	-2.9	-2.9	-2.3
hi1033	Phosphoserine phosphatase	-5.0	-2.8	-2.3
hi1632	Lysine-sensitive aspartokinase III	-3.3	-2.4	-2.3
hi1227	Uracil permease	-2.3	-2.8	-2.4
hi1041	Modification methylase HgiDI	-2.8	-2.4	-2.5
hi1163	D-Lactate dehydrogenase	-3.2	-3.2	-2.5
hi0117	Membrane-bound lytic murein transglycosylase A	-2.5	-2.8	-2.6
hi1022	Biotin synthetase	-4.7	-3.4	-2.6
hi0563	Regulatory protein AsnC	-2.7	-2.5	-2.6
hi0539	Urease, alpha subunit	-3.2 (NC)	-3.8 (NC)	-2.6 (NC)
hi1201	Adenine specific methylase, putative	-2.7	-2.9	-2.7
hi1682	Putative protease	-2.9	-2.3	-2.7
hi1065	MukB suppressor protein	-2.5	-2.5	-2.9
hi1545	C4-Dicarboxylate transport protein	-5.0	-3.9	-2.9
hi1383	Periplasmic phosphate-binding protein	-2.1	-4.0	-3.0
hi0078	Cys-tRNA synthetase	-2.6	-2.5	-3.0
hi1263	Homoserine acetyltransferase	-2.7	-2.6	-3.1
hi0419	Collagenase	-4.3	-3.2	-3.2
hi1106	Xylose operon regulatory protein	-2.8	-2.7	-3.2
hi1507	Hypothetical protein	NC (-4.1)	-3.5 (-3.2)	-3.2 (-2.9)
hi1068	Formate-dependent nitrite reductase	-3.0	-2.7	-3.3
hi1735	Peptide-chain-release factor 3	-4.3	-3.3	-3.4
hi1112	Xylose isomerase	-2.3	-3.2	-3.4
hi1019	Thiamin-binding periplasmic protein	-5.0	-4.7	-3.4
hi0445	Protein translocation protein, low temperature	-3.6	-4.6	-3.8
hi0548	Initiation factor IF-1	-7.7	-5.8	-4.0

**Table 3.** (Continued)

No. <sup>a</sup>	Gene product	Fold change mRNA (protein)		
		10 min	30 min	60 min
hi1181	Phosphoheptose isomerase	-4.2	-5.2	-4.0
hi1172	S-Adenosylmethionine synthetase 2 <sup>c</sup>	-3.9 (NC)	-4.1 (3.0)	-4.0 (NC)
hi1166	Histidinol-phosphate aminotransferase	-6.5	-4.1	-4.1
hi1171	Anthranilate synthase component II	-3.8	-3.4	-4.1
hi0501	High affinity ribose transport protein	-2.7	-2.4	-4.1
hi0182	Glucose kinase	-3.1	-5.2	-4.3
hi1728	Branched chain amino acid transport system II carrier protein	-9.5	-7.5	-4.5
hi1170	Anthranilate synthase component I	-5.7	-4.2	-4.8
hi0549	16S rRNA-dimethyltransferase	-7.6	-5.0	-4.9
hi0808	Ribosome releasing factor	-4.5 (-4.7)	-5.3 (-4.4)	-4.9 (-4.7)
hi0214	Oligopeptidase A	-4.8	-6.8	-5.0
hi1009	Glycerol-3-phosphate regulon repressor	-7.4	-9.8	-5.0
hi0179	Pyruvate formate-lyase activating enzyme	-6.9	-6.1	-5.1
hi0211	Phosphatidylglycerophosphatase B	-4.5	-7.6	-5.7
hi1365	Topoisomerase I	-4.4	-5.8	-6.1
hi1167	Phosphoserine aminotransferase	-9.1	-9.6	-6.2
hi1154	Proton glutamate symport protein	-4.2	-4.1	-6.3
hi1729	Lactam utilization protein	-6.9	-9.3	-6.5
hi1048	Hypothetical protein	-9.1	-14.1	-9.6
hi1035	Magnesium and cobalt transport protein	-9.2	-10.5	-12.2

<sup>a</sup>Numbering according to Fleischmann et al. (1995).

<sup>b</sup>Present as two spots; other spot decreased to a lesser extent.

<sup>c</sup>Present as two spots; other spot increased in intensity at the 30-min time point.

the higher concentration, confirming those results, and the fold changes after incubation with the higher inhibitor concentration were, in most cases, more pronounced. With Novobiocin, the incubation time did not play an important role, indicating that the response to the inhibitor is rather fast and remains constant over the time measured, perhaps reflecting the bacteriostatic effect of this type of inhibitor.

The rates of synthesis of proteins largely followed that of the corresponding mRNA (Table 4). For some proteins the rates were in full agreement (e.g., heat shock protein GroES, ribosome release factor), indicating that their expression is transcriptionally regulated. The changes of other protein synthesis rates were much more pronounced than that of their corresponding mRNAs (e.g., ribosomal proteins, elongation factor Ts), indicative of a translational mode of regulation. This is well documented in *Escherichia coli* for the components of the translational apparatus (Zengel and Lindahl 1994). The synthesis of phosphoglycerate kinase was detected as down-regulated at the protein level but as up-regulated at the mRNA level. Because the *pgk* gene is transcriptionally regulated in *E. coli* (Charpentier et al. 1998), it is unlikely that regulation of this gene takes place entirely on the translational level in *H. influenzae*. The existence of a second, unidentified spot corresponding to this protein or a quantification artifact due to spot overlapping are two possible explanations for this finding.

#### Ciprofloxacin

Quinolones are bactericidal and provoke the so-called RecA (SOS) DNA repair system (Piddock and Wise 1987; Walters et al. 1989; Piddock et al. 1990). They form a stable ternary complex with DNA gyrase and DNA and thereby inhibit replication and transcription. Probably because of this different mode of action, incubation of the *H. influenzae* with Ciprofloxacin resulted in a somewhat different change in the expression pattern. At a low inhibitor concentration, only a few genes showed changed expression, most of them being involved in SOS repair (*recA*, *uvrA*, *lexA*) (Table 5).

The first genes showing changed expression after addition of Ciprofloxacin belong or may belong to DNA repair systems. After 10 min incubation in the presence of the higher Ciprofloxacin concentration, ~15 genes showed a change in expression and after 60 min ~140 genes were called increased or decreased (Table 6). In addition, about 40 ORFs coding for hypothetical proteins also reproducibly show changed expression patterns. At the higher concentration (300 ng/mL), the mRNA analyses for the 30-min time point did not yield consistent results and were therefore not included in the analysis.

At the higher Ciprofloxacin concentration many genes showed changed expression. Unlike Novobiocin, this inhibitor triggered a time-dependent response. After 60 min about 10-fold more genes showed changed expression than after 10 min. This may indicate that,

**Table 4.** Changes at the Protein Level after Addition of 125 ng/mL of Novobiocin

No.	Gene product	Fold change protein (mRNA)		
		10 min	30 min	60 min
hi0048	Oxidoreductase	-3.1 (NC)	-3.5 (NC)	-1.7 (NC)
hi0089	Aspartokinase I/homoserine dehydrogenase I	-2.6 (-3.4)	-1.3 (NC)	-2.0 (-2.0)
hi0119	Adhesin B precursor	-261.8 (NC)	-12.8 (-2.8)	-34.1 (-2.7)
hi0124	Inorganic pyrophosphatase	-2.1 (1.2)	-3.3 (NC)	-5.4 (NC)
hi0504	Periplasmic ribose-binding protein	-1.8 (-2.2)	-2.5 (-3.9)	-3.5 (-1.9)
hi0524	Fructose-bisphosphate aldolase	2.0 (1.8)	2.1 (1.6)	1.0 (1.9)
hi0525	Phosphoglycerate kinase	-2.0 (2.2)	-2.1 (2.1)	-2.9 (2.0)
hi0542	Heat shock protein GroES	-1.9 (-2.3)	-4.4 (-4.6)	-3.5 (-3.7)
hi0544	Ribosomal protein L9	-3.5 (-1.5)	-2.0 (-1.7)	-3.6 (NC)
hi0547	Ribosomal protein S6	-4.2 (NC)	-2.1 (NC)	-2.7 (NC)
hi0553	6-Phosphogluconate dehydrogenase, decarboxylating	-5.5 (NC)	-4.4 (NC)	-3.2 (NC)
hi0574	Conserved hypothetical protein	-3.8 (-2.1)	-3.8 (-2.9)	-4.0 (-2.3)
hi0808	Ribosome releasing factor	-4.7 (-4.5)	-4.4 (-5.4)	-4.7 (-4.9)
hi0846	Disulfide oxidoreductase	-2.6 (NC)	-2.0 (-1.9)	-2.4 (-2.1)
hi0913	Ribosomal protein S2	-2.3 (NC)	-3.8 (-2.4)	-5.2 (-2.2)
hi0914	Elongation factor EF-Ts	-4.2 (NC)	-5.5 (-3.0)	-2.9 (-2.6)
hi0927	Glycyl-tRNA synthetase alpha chain	2.5 (NC)	2.9 (1.5)	-1.2 (1.6)
hi1228	Uracil phosphoribosyltransferase	-3.0 (-2.0)	-3.2 (-2.2)	-3.4 (-2.0)
hi1507	Hypothetical protein	-4.1 (1.3)	-3.2 (-3.5)	-2.9 (-3.2)
hi1647	Conserved hypothetical protein	-2.0 (-2.3)	-1.8 (-1.8)	-2.3 (-1.6)
hi1734	Enoyl-reductase	-8.8 (-2.6)	-7.2 (-2.6)	-6.2 (-2.1)

although only a few genes are involved in the primary response, many more gene products are involved in the following cellular processes. Not only the previously mentioned genes involved in the SOS repair mechanism but also others, presumably involved in DNA repair, showed an increased expression level (*ruvB*, *recO*, *recN*, *impA*, *recF*). Noteworthy is the fact that topoisomerase IV, which seems to be the primary target for quinolones in other species (see, for example, Fukuda and Hiramatsu 1999; Munoz and De La Campa 1996), is not drastically changed in *H. influenzae* (fold changes for ParE = 1.7 and for ParC = NC), suggesting that in this species DNA gyrase is the primary target for

quinolones. This is supported by studies that identified *gyrA* mutations as a cause for the development of Ciprofloxacin resistance in *H. influenzae* (Bootsma et al. 1997). As for Novobiocin, the lower Ciprofloxacin concentration and the early time points may be the conditions best suited for the analysis of the primary response.

At the proteome level, Ciprofloxacin changed the synthesis rates of fewer proteins than did Novobiocin and with a lower amplitude. This is in agreement with the weaker effect detected by mRNA analysis. For Novobiocin, the changes were, qualitatively, in good agreement with the mRNA data (Table 7).

**Table 5.** Genes with Changed Expression Levels after Addition of 30 ng/mL of Ciprofloxacin

No.	Gene product	Fold change mRNA <sup>a</sup> (protein)		
		10 min	30 min	60 min
hi1546	ImpA protein	NC	6.2	6.3 ± 4.2
hi0070	DNA repair protein	NC	3.0	5.9 ± 3.3
hi0249	Excinuclease ABC subunit A	NC	3.4	3.7 ± 0.8
hi0250	Single-stranded DNA binding protein	NC (NC)	2.0 (NC)	1.7 (NC)
hi0312	Holliday junction DNA helicase	3.2	2.9	4.7 ± 2.2
hi0313	Holliday junction DNA helicase	NC	2.6	3.1 ± 0.9
hi0600	RecA recombinase	NC	2.7	3.3 ± 0.9
hi0749	LexA repressor	NC	2.4	3.5 ± 2.6
hi0525	Phosphoglycerate kinase	NC (1.9)	NC (2.3)	NC (3.2)

<sup>a</sup>Standard deviations were below 20% if not otherwise indicated.

**Table 6.** Genes Showing Major Changes in Expression after Addition of 300 ng/mL of Ciprofloxacin

No.	Gene product	Fold change	
		10 min	60 min
hi0567	DNA gyrase, subunit B	2.1	1.9
hi1264	DNA gyrase, subunit A	1.8	1.8
hi1365	Topoisomerase I	NC	-3.1
hi0070	DNA repair protein RecN	4.8	13.0
hi0749	LexA repressor	3.9	7.0
hi1349	Neutrophil activating protein	NC (1.7)	6.3 (3.4)
hi0750	Diaminopimelate epimerase	2.2	5.4
hi0965	Ribosomal protein S20	1.7	5.3
hi1188	DNA helicase II	1.9	4.5
hi0249	Excinuclease ABC subunit A	2.3	4.4
hi0715	ATP-dependent protease ATPase subunit (Clpx)	NC (NC)	4.2 (NC)
hi0313	Holliday junction DNA helicase	1.5	3.9
hi0459	Pyrimidine operon regulatory protein	NC (NC)	3.6 (NC)
hi1392	HindIII modification methyltransferase	NC	3.5
hi0332	DNA repair protein (RecO)	NC	3.5
hi0623	Methionyl-tRNA formyltransferase	2.4	3.4
hi0330	Cell envelope protein	1.9	3.3
hi0312	Holliday junction DNA helicase	1.8	3.3
hi0713	Trigger factor	2.1 (NC)	3.3 (1.9)
hi0394	Peptidyl-tRNA hydrolase	NC	3.3
hi1125	Transaldolase B	2.1 (NC)	3.2 (2.1) <sup>a</sup>
hi0250	Single-stranded DNA binding protein	2.5 (2.0)	2.9 (1.5)
hi0113	Heme-hemopexin utilization protein C	-1.6	2.8
hi0256	Lipoprotein-34	NC	2.8
hi0079	Peptidyl-prolyl cis-trans isomerase B	NC (NC)	2.8 (NC)
hi0872	Undecaprenyl-phosphate galactosephosphotransferase	NC	2.8
hi0970	3-Dehydroquinase	NC	2.8
hi1384	Ferritin like protein	NC	2.7
hi0413	Ribonuclease E	2.4	2.7
hi1742	RNA polymerase omega subunit	1.5	2.7
hi0602	HemY protein homolog	1.6	2.6
hi0717	Transcription antitermination protein NusG	NC	2.6
hi1447	GTP cyclohydrolase I	NC (1.7)	2.6 (NC)
hi0411	Host factor-I	2.1	2.6
hi0287	Tryptophan-specific permease	NC	2.6
hi0331	Opacity associated protein	NC	2.5
hi0991	DNA/ATP binding protein	NC	2.5
hi0622	Polypeptide deformylase	NC	2.5
hi0206	5-Nucleotidase, putative	NC (NC)	2.5 (NC)
hi0603	Uroporphyrinogen III methylase	NC	2.4
hi1325	3-Hydroxydecanoyl-(acyl carrier-protein) dehydratase	NC	2.4
hi0993	Chromosomal replication initiator protein	NC	2.4
hi0865	Glutamine synthetase	1.6	2.4
hi1561	Polypeptide chain release factor 1	NC (NC)	2.3 (NC)
hi1304	N utilization substance protein B	1.8	2.3
hi1740	DNA recombinase RecG	NC	2.3
hi1305	Hydrogenase gene region	NC	2.2
hi0881	Octaprenyl-diphosphate synthase	NC (NC)	2.2 (NC)
hi1385	Ferritin like protein	NC (1.4)	2.2 (NC)
hi0927	Glycyl-tRNA synthetase alpha chain	NC (NC)	2.1 (1.7)
hi0921	Leucyl-tRNA synthetase	2.1	2.1
hi0718	VacJ lipoprotein	NC	2.1
hi0689	Glycerophosphoryl diester phosphodiesterase	NC	2.1
hi0768	Cell division protein FtsY	NC (NC)	2.1 (NC)
hi0624	Fmu/Fmv gene product	2.1	2.1
hi0600	DNA recombinase RecA	2.7	2.0
hi0567	DNA gyrase, subunit B	2.1	1.9
hi1220	Ribosomal protein S1	3.3	1.9
hi0785	Ribosomal protein L29	2.6	NC
hi0516	Ribosomal protein L1	2.0	NC
hi0782	Ribosomal protein L22	3.0	NC
hi0779	Ribosomal protein L23	2.2	NC
hi0571	Hydrogen peroxide-inducible activator	NC	-2.1
hi0150	Lambda cII stability-governing protein	NC	-2.1

**Table 6.** (Continued)

No.	Gene product	fold change	
		10 min	60 min
hi0629	Negative rpo regulator	NC	-2.1
hi1135	Phospho-N-acetylmuramoyl-pentapeptide-transferase E	-2.2	-2.1
hi1251	Virulence associated protein A	NC	-2.2
hi0508	Menaquinone biosynthesis protein	NC	-2.2
hi0538	Urease accessory protein	-1.6	-2.2
hi0506	<i>rbs</i> repressor	NC	-2.2
hi1661	Dihydroliipoamide succinyltransferase	NC (NC)	-2.2 (1.7)
hi0546	Primosomal protein N presursor	NC	-2.2
hi0239	Protein-export membrane protein SecF	-2.0	-2.3
hi0745	L-Asparaginase II	NC (NC)	-2.3 (-2.0)
hi0470	Histidinol-phosphate aminotransferase	NC	-2.3
hi0008	Formate dehydrogenase-O gamma subunit	NC	-2.3
hi1133	UDP N-acetylmuramyl-tripeptide synthetase	NC	-2.3
hi0140	N-Acetylglucosamine-6-phosphate deacetylase	NC	-2.4
hi0117	Membrane-bound lytic murein transglycosylase A	-1.8	-2.4
hi1154	Proton glutamate symport protein	-1.9	-2.4
hi1218	L-Lactate permease	NC	-2.4
hi0537	Urease accessory protein	NC	-2.4
hi0373	Heat shock cognate 66	NC	-2.5
hi0142	N-Acetylneuraminase lyase	NC	-2.5
hi1227	Uracil permease	-2.3	-2.5
hi0505	Ribokinase	NC	-2.6
hi0544	Ribosomal protein L9	NC (NC)	-2.6 (NC)
hi1167	Phosphoserine aminotransferase	NC	-2.6
hi0536	Urease accessory protein	NC	-2.6
hi0087	Threonine synthase	NC (-2.6) <sup>a</sup>	-2.7 (NC)
hi0089	Aspartokinase I/homoserine dehydrogenase I	NC (NC)	-2.8 (NC)
hi1367	Threonyl-tRNA synthetase	NC (NC)	-2.8 (NC)
hi1511	Sheath protein Gpl	NC	-2.8
hi0447	L-Phosphofructokinase	NC	-2.8
hi1172	S-Adenosylmethionine synthetase 2	NC (-1.3)	-2.8 (-1.9) <sup>b</sup>
hi0564	Asparatate ammonia ligase	NC (NC)	-2.8 (NC)
hi0088	Homoserine kinase	NC (NC)	-2.9 (NC)
hi1545	C4-dicarboxylate transport protein	-2.4	-2.9
hi0189	Glutamate dehydrogenase	NC	-3.0
hi0465	Phosphoglycerate dehydrogenase	NC (NC)	-3.1 (NC)
hi0446	Fructose-permease IIBC component	-2.3	-3.1
hi1727	Argininosuccinate synthetase	NC (NC)	-3.2 (NC)
hi1181	Phosphoheptose isomerase	NC	-3.3
hi1177	Arginine transport system permease protein	-2.9	-3.4
hi0548	Initiation factor IF-1	-2.7	-3.4
hi0141	Glucosamine-6-phosphate isomerase	NC	-3.7
hi0913	Ribosomal protein S2	NC (NC)	-3.9 (-4.3)
hi1179	Periplasmic arginine-binding protein	NC (NC)	-4.0 (NC)
hi0509	Menaquinone biosynthesis protein	NC	-4.0
hi0628	RNA polymerase sigma-E factor	NC	-4.0
hi1636	Phosphoenolpyruvate carboxylase	NC	-4.4
hi1178	Arginine transport system permease protein	-2.1	-4.6
hi1729	Lactam utilization protein	NC	-5.2
hi1728	Branched chain amino acid transport system II carrier protein	-2.6	-5.5
hi1180	Arginine transport ATP-binding protein	-2.2	-6.4

<sup>a</sup>Only one of two spots changed in intensity.<sup>b</sup>The second spot corresponding to this protein was increased.

#### Expression Changes Specific for Novobiocin or Ciprofloxacin

The response to Ciprofloxacin shows some clear differences when compared with the Novobiocin-induced response. As shown in Table 8, the induction of DNA repair systems distinguished the response of the cells to Ciprofloxacin from that to Novobiocin. Furthermore,

the onset of the response was delayed for Ciprofloxacin and immediate for Novobiocin. The induction of these proteins could therefore represent a molecular marker to distinguish between the response to quinolones compared with other DNA gyrase inhibitors. Novobiocin had little effect at a low concentration. At the

**Table 7.** Changes at the Protein Level after Addition of 300 ng/mL of Ciprofloxacin

No.	Gene product	Fold change protein (mRNA)		
		10 min	30 min	60 min
hi0250	Single-stranded DNA binding protein	2.0 (2.5)	2.8	1.5 (2.9)
hi0496	HslUV operon heat shock protein	3.6 (NC)	2.9	3.2 (NC)
hi0574	Conserved hypothetical protein	-2.5 (NC)	-2.6	-1.5 (-2.6)
hi0914	Elongation factor EF-Ts	-2.2 (NC)	-2.8	-3.2 (-5.6)
hi0924	Glycyl-tRNA synthetase beta chain	NC (NC)	-2.4	3.4 (1.6)
hi1116	Deoxyribose aldolase	1.4 (NC)	2.2	2.7 (1.8)
hi1172	S-Adenosylmethionine synthetase 2	-1.3 (NC)	-5.2	-1.9 (-2.8)
hi1303	Riboflavin synthase, beta chain	2.7 (NC)	3.0	1.7 (1.8)
hi1349	Neutrophil activating protein	1.7 (NC)	3.1	3.4 (6.3)

higher concentration, however, it affected many more genes than did Ciprofloxacin, suggesting that it exerts a stronger action on the cell at  $10\times$ MIC than does Ciprofloxacin. It is therefore difficult to determine genes that are specifically affected by Novobiocin. Detailed studies of concentration-dependent responses could clarify this issue.

#### Common Effects

To examine whether the commonly affected genes code for proteins belonging to a specific functional group, they were classified as increased, decreased, or not changed at the respective highest antibiotic concentration and at the 60-min time point and were ordered in functional groups according to the scheme proposed by Fleischmann et al. (1995) (Table 9). Overall, almost twice as many genes were down-regulated as opposed to up-regulated. Many amino acid biosynthesis enzymes and amino acid transporters were detected as down-regulated. In line with this finding, the expression of genes coding for ribosomal proteins and for enzymes indirectly linked to amino acid biosynthesis (urease components, molybdopterin biosynthesis enzymes) was also decreased. On the other hand, the synthesis rates for most tRNA synthetases were increased. This could suggest that exposure to gyrase inhibitors leads to a decrease in protein biosynthetic activity, although we did not observe a decreased methionine incorporation in the cultures treated with the antibiotics.

#### Conclusions

In summary, the results show that the high-density microarrays yield highly reproducible results and that the main difficulties for reproducible analysis of low abundance transcripts lie in biological sampling. Our results show that if conditions are kept as reproducible as possible, most of the transcripts can be detected even when present at low concentrations. Our studies analyzing the number of transcripts called present after hybridization showed that in all experiments, up to

85% of transcribed genes could be detected. Transcripts showing very low intensity values, only a few percent above background fluorescence, could also be reproducibly identified as present, although the uncertainties of the signals in this fluorescence range is higher. Not surprisingly, the sensitivity and reproducibility of the expression analysis by using oligonucleotide chip technology was clearly better than expression analysis by using 2D-PAGE followed by computerized image analysis. The dynamic ranges were  $10^3$  for transcript analysis and  $10^5$  for the quantification of proteins. However, although qualitatively similar, there are some quantitative differences in the response detected by protein quantification compared with mRNA quantification. This highlights the importance of combining both technologies to obtain important information on the level (transcriptional or translational) at which the regulatory mechanisms act. Moreover, although relatively rare in bacteria, posttranslational modifications constitute an important additional level of regulation and can only be studied by proteome investigations. The detection of proteins present as multiple spots underscores this point.

Expression analysis by using the bacterial microarray system or 2D gels can be used to profile the effect of an inhibitor on a cell, but the main initial challenge is to discover the appropriate concentration and time point. In the described experiments a low concentration, around the MIC values, and a rather short incubation period that is within minutes, seem to provide the best results for the detection of the genes that are primarily affected. Because of the overwhelming amount of data it is more difficult to analyze the responses after incubation with a higher inhibitor concentration and/or after longer incubation times but, on the other hand, these results may help us analyze and understand more complex response patterns. Profiling the response of a selected inhibitor class may also give indications for a classification of an unknown inhibitor because its profile can be compared with those from known inhibitors. The example of Ciprofloxacin and

**Table 8.** Expression Changes Specific for Ciprofloxacin or Novobiocin

No.	Gene product	Fold change mRNA			
		Cipro 10 min	Novo 10 min	Cipro 60 min	Novo 60 min
Specific for Ciprofloxacin					
hi0020	2-Oxoglutarate/malate translocator	-2.0	NC	-1.7	NC
hi1135	Phospho-N-acetylmuramoyl-pentapeptide-transferase	-2.2	NC	-2.1	NC
hi0070	DNA repair protein RecN	4.8	NC	13.0	NC
hi0162	Conserved hypothetical protein	1.5	NC	1.8	NC
hi0250	Single-stranded DNA binding protein	2.5	NC	2.9	NC
hi0312	Holliday junction DNA helicase	1.8	NC	3.3	NC
hi0600	Recombinase RecA	2.7	NC	2.0	-1.8
hi1159	Conserved hypothetical protein	1.5	-3.5	3.0	-1.8
hi1188	DNA helicase II	1.9	NC	4.5	NC
hi1546	ImpA protein	4.8	NC	15.0	NC
Specific for Novobiocin					
hi0343	Conserved hypothetical protein	NC	-2.7	NC	-2.5
hi0365	Conserved hypothetical protein	NC	-4.4	NC	-2.6
hi0434	Transformation protein	NC	-7.1	NC	-4.3
hi0440	Penicillin-binding protein	NC	-2.2	NC	-1.1
hi0498	Spermidine/putrescine-binding periplasmic protein precursor	NC	-2.4	NC	-1.4
hi0558	Glucose-6-phosphate 1-dehydrogenase	NC	-2.3	NC	-1.4
hi0838	Small protein a	NC	-2.8	NC	-1.7
hi0898	Multidrug resistance protein	NC	-2.5	NC	-1.5
hi0983	Conserved hypothetical protein	NC	-2.9	NC	-1.9
hi1151	Conserved hypothetical protein	NC	-3.9	NC	-2.8
hi1159	Conserved hypothetical protein	NC	-3.5	NC	-1.8
hi1163	D-Lactate dehydrogenase	NC	-3.2	3.6	-2.5
hi1190	Conserved hypothetical protein	NC	-4.1	NC	-9.8
hi1191	Conserved hypothetical protein	NC	-11.5	NC	-17.3
hi1388	Anthranilate synthase component ii	NC	-3.3	NC	0.3
hi1585	Acetolactate synthase iii large chain	NC	-2.5	NC	-1.4
hi0037	Rod shape-determining protein	NC	2.0	1.7	1.8
hi0690	Glycerol uptake facilitator protein	NC	1.6	NC	1.2
hi0691	Glycerol kinase	NC	1.7	NC	1.4
hi0971	Acetyl-con carboxylase, biotin carboxyl carrier protein	NC	1.7	NC	1.8
hi1308	Dihydrodipicolinate reductase	NC	2.0	NC	1.9
hi1623	Mercury resistance regulatory protein	NC	3.2	NC	3.2
hi1659	Ribonucleoside-diphosphate reductase, alpha chain	NC	1.7	3.6	2.0
hi1660	Ribonucleoside diphosphate reductase, beta chain	NC	2.3	NC	2.2
hi1717	Hypothetical protein	NC	1.8	NC	1.9

Novobiocin illustrates that the response to an antibiotic can yield important information as to its mode of action. Both compounds induce the expression of DNA gyrase and negatively affect topoisomerase I expression. Ciprofloxacin, in addition, induces the SOS response. The two different modes of action are thus clearly reflected in the cellular response. These genes may therefore be useful indicators for gyrase inhibition by either mechanism. More experiments with different gyrase inhibitors would be required to substantiate these findings. Expression analysis will prove to be an invaluable tool not only for the study of disease processes but also for the characterization of novel pharmaceuticals.

## METHODS

### Cell Growth

*H. influenzae* Rd KW 20 was used as the model organism for these experiments. Bacterial cultures were grown in minimal medium with a reduced methionine concentration (0.6  $\mu$ M) to an OD<sub>600</sub> of 0.4 (Barcak et al. 1991). The cultures were divided into aliquots of 300 mL and antibiotics were added to a final concentration of 30 and 300 ng/mL for Ciprofloxacin, and 12.5 and 125 ng/mL for Novobiocin, respectively. Controls without antibiotic were grown in parallel. After 10, 30, and 60 min, aliquots were taken for metabolic labeling and RNA extraction. To the first aliquot (1mL), 0.7 MBq of L-[<sup>35</sup>S]Methionine (>37 TBq/mmol, Amersham Radiochemicals) were added and incubation was continued for 2 min. The cells were then rapidly chilled on ice, harvested by centrifugation.

**Table 9.** Common Effects for Novobiocin or Ciprofloxacin

Functional Group <sup>a</sup>	Total no. of genes	Novobiocin				Ciprofloxacin			
		Increased	%	Decreased	%	Increased	%	Decreased	%
Amino acid biosynthesis	70	5	7%	29	41%	2	3%	16	23%
Aromatic amino acid family	16	1	6%	4	25%		0%	1	6%
Aspartate family	18	1	6%	7	39%	1	6%	8	44%
Branched chain family	10		0%	5	50%		0%		0%
Glutamate family	9	2	22%	1	11%	1	11%	4	44%
Histidine family	9		0%	9	100%		0%	1	11%
Serine family and pyruvate family	8	1	13%	3	38%		0%	2	25%
Biosynthesis of cofactors, prosthetic groups, and carriers	56	5	9%	16	29%		0%	2	4%
Cell envelope	79	12	15%	11	14%	2	3%	5	6%
Cellular processes	53	5	9%	12	23%	1	2%	5	9%
Cell division	17	3	18%	6	35%		0%		0%
Chaperones	6		0%	2	33%		0%	2	33%
Others	30	2	7%	4	13%	1	3%	3	10%
Central intermediary metabolism	30	1	3%	12	40%		0%	7	23%
Energy metabolism	109	8	10%	21	19%	4	6%	17	16%
Fatty acid and phospholipid metabolism	26	11	42%	5	19%	2	8%	2	8%
Hypothetical	339	4	76%	22	13%	11	3%	22	6%
Other categories	95	7	7%	14	15%	1	1%	5	5%
Purines, pyrimidines, nucleosides, and nucleotides	52	6	12%	5	10%	1	2%	5	10%
Regulatory functions	68	6	9%	11	16%	2	3%	8	12%
Replication, recombination, and DNA repair	84	18	21%	14	17%	13	15%	3	4%
Transcription	26	4	15%	4	15%	3	12%	1	4%
DNA-dependent RNA polymerase	17	3	18%	3	18%	2	12%	1	6%
Degradation of RNA	9	1	11%	1	11%	1	11%		0%
Translation	143	18	13%	23	16%	15	10%	17	12%
Amino acyl tRNA synthetases	34	9	26%	4	12%	6	18%	3	9%
Ribosomal proteins synthesis and modification	57	2	4%	3	5%	3	5%	8	14%
Others	52	7	13%	16	31%	6	12%	6	12%
Transport and binding proteins	123	10	8%	31	25%	2	2%	19	15%
Amino acids peptides and amines	38	3	8%	12	32%		0%	8	21%
Other transport and binding proteins	85	7	8%	19	11%	2	2%	11	13%
Total	1353	132	10%	284	21%	62	5%	134	10%

<sup>a</sup>As defined by Fleischmann et al. (1995).

gation, and frozen at  $-20^{\circ}\text{C}$ . For RNA preparation, a 35-mL aliquot was spun in a chilled centrifuge and the cell pellet was snap-frozen in liquid nitrogen and kept at  $-80^{\circ}\text{C}$ . Five individual experiments were performed, whereby aliquots for RNA extraction were only collected for two experiments.

### RNA Extraction, Preparation, Array Hybridization, and Scanning

Bacterial RNA was isolated, labeled, and hybridized to the chips essentially as described (de Saizieu et al. ). Before fragmentation of the biotin-labeled cDNA, an additional purification step was performed by using Chromaspin-100 columns (Clontech) and the fragmented cDNA was centrifuged quickly through 0.22- $\mu\text{m}$  filter units (Millipore).

### 2D-PAGE

The cell pellets were washed once in PBS buffer (Life Technologies). The cells were then lysed by resuspension in sample

buffer containing 8 M urea, 4% CHAPS, 40 mM of Tris base (Fluka), 65 mM of 1,4-dithioerythritol (Merck), and 2% ampholytes (Resolyte 3-10, BDH). The extracts were centrifuged at 100,000g and the supernatant recovered. The amount of incorporated radioactivity was determined in a Model 2500 TR liquid scintillation counter (Packard Instrument Co.).

Aliquots of the protein extracts containing  $4 \times 10^6$  cpm of radioactivity were loaded onto Immobiline 3–10 nonlinear pH gradient strips (Pharmacia) at the basic end and resolved according to the manufacturer's recommendations. The strips were equilibrated as described (Sanchez et al. 1995) and loaded onto 1-mm thick vertical 12% polyacrylamide slab gels. After electrophoresis, the gels were dried on 3 MM Whatman filter paper and exposed to PhosphorImager screens (Molecular Dynamics). Images were analyzed by using PDQuest software (BioRad). Parallel samples (one sample per time point and concentration and their corresponding controls) were run on parallel gels (same batch of strips, same isoelectric focussing run, same batch of gels for SDS-PAGE, and same

**Table 10.** Parameters Used for Data Analysis

Parameter	Description
Average difference <sup>a</sup>	Average of the differences in intensity values between the perfect match and the mismatch oligonucleotides over the entire probe set for one gene (arbitrary units)
Average difference change <sup>a</sup>	Average difference (induced)—average difference (control)
Difference call <sup>a</sup>	Five possible values: D = decreased; MD = marginally decreased; NC = not changed; MI = marginally increased; I = increased
Fold change <sup>a</sup>	Permits the symmetrical expression of the fold change as a positive number when the transcript has increased over its baseline state, and as a negative number when the transcript level declines. Corresponds to average difference (induction)/average difference (control) for induced and average difference (control)/average difference (induction) for repressed genes. For weakly expressed genes, values are corrected for noise.
Change factor	For increased intensity: fold change – 1; For decreased intensity: fold change + 1
Sort score <sup>a</sup>	Quality measure based on fold change and average difference change
Purity	(# pairs used – least (# of pairs called increased, # of pairs called decreased)*2)/pairs used

<sup>a</sup>Output from the Affymetrix GeneChip software.

SDS-PAGE run). Only pairs of gels that had been obtained under identical conditions were considered for analysis.

The 2D gels were matched and the data were stored in an Oracle database. The spot intensities were normalized so that the sum of all the spot intensities was equal for all gels. The spot intensity data were exported to Microsoft Excel for further analysis.

### Data Handling and Analysis

For transcriptional imaging, the hybridization intensities were processed by using Affymetrix GeneChip software. Pair-wise comparisons of hybridization intensities were performed and the results were exported to Microsoft Access for further analysis. The parameters that were used for analysis were partly adopted from and partly derived from the output of the GeneChip software (Table 10).

The fold change values were averaged for all experiments performed in duplicate or triplicate. Results were considered significant if the averaged fold change was >2 or <-2, the standard deviation of the averaged fold change <0.25, the difference call “D” or “I”, the purity factor >0.9, and the sort score >1 or <-1.

The spot intensities from the 2D gel experiments were exported to Excel and the significance of the results was estimated by using the *t*-test for paired samples. When the obtained *P* values were lower than .05, the changes were considered significant. For the calculation of average fold change values, the values were first converted to change factor format. The averages were then reconverted to fold change format to improve comparability and clarity.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked “advertisement” in accordance with 18 USC section 1734 solely to indicate this fact.

### REFERENCES

- Ali, J.A., Jackson, A.P., Howells, A.J., and Maxwell, A. 1993. The 43-kilodalton N-terminal fragment of the DNA gyrase B protein hydrolyzes ATP and binds coumarin drugs. *Biochemistry* **32**: 2717–2724.

Ali, J.A., Orphanides, G., and Maxwell, A. 1995. Nucleotide binding to the 43-kilodalton N-terminal fragment of the DNA gyrase B protein. *Biochemistry* **34**: 9801–9808.

Anderson, L. and Seilhamer, J. 1997. A comparison of selected mRNA and protein abundances in human liver. *Electrophoresis* **18**: 533–537.

Barcak, G.J., Chandler, M.S., Redfield, R.J., and Tomb, J.F. 1991. Genetic systems in *Haemophilus influenzae*. *Methods Enzymol.* **204**: 321–342.

Bootsma, H.J., Troelstra, A., van Veen-Rutgers, A., Mooi, F.R., de Neeling, A.J., and Overbeek, B.P. 1997. Isolation and characterization of a ciprofloxacin-resistant isolate of *Haemophilus influenzae* from the Netherlands. *J. Antimicrob. Chemother.* **39**: 292–293.

Charpentier, B., Bardey, V., Robas, N., and Branlant, C. 1998. The EII<sub>Glc</sub> protein is involved in glucose-mediated activation of *Escherichia coli* gapA and gapB-pgk transcription. *J. Bacteriol.* **180**: 6476–6483.

Chee, M., Yang, R., Hubbell, E., Berno, A., Huang, X.C., Stern, D., Winkler, J., Lockhart, D.J., Morris, M.S., and Fodor, S.P.A. 1996. Accessing genetic information with high-density DNA arrays. *Science* **274**: 610–614.

Contreras, A. and Maxwell, A. 1992. gyrB mutations which confer coumarin resistance also affect DNA supercoiling and ATP hydrolysis by *Escherichia coli* DNA gyrase. *Mol. Microbiol.* **6**: 1617–1624.

de Saizieu, A., Gardes, C., Flint, N., Wagner, C., Kamber, M., Mitchell, T., Keck, W., Amrein, K., and Lange, R. Microarray based identification of a novel *Streptococcus pneumoniae* regulon controlled by an autoinduced-peptide. *J. Bacteriol.* **182**: 4696–4703.

Drlica, K. and Zhao, X. 1997. DNA gyrase, topoisomerase IV, and the 4-quinolones. *Microbiol. Mol. Biol. Rev.* **61**: 377–392.

Fleischmann, R.D., Adams, M.D., White, O., Clayton, R.A., Kirkness, E.F., Kerlavage, A.R., Bult, C.J., Tomb, J.F., Dougherty, B.A., Merrick, J.M., et al. 1995. Whole-genome random sequencing and assembly of *Haemophilus influenzae* Rd. *Science* **269**: 496–512.

Fodor, S.P., Read, J.L., Pirrung, M.C., Stryer, L., Lu, A.T., and Solas, D. 1991. Light-directed, spatially addressable parallel chemical synthesis. *Science* **251**: 767–773.

Fodor, S.P., Rava, R.P., Huang, X.C., Pease, A.C., Holmes, C.P., and

- Adams, C.L. 1993. Multiplexed biochemical assays with biological chips. *Nature* **364**: 555–556.
- Fukuda, H. and Hiramatsu, K. 1999. Primary targets of fluoroquinolones in *Streptococcus pneumoniae*. *Antimicrob. Agents Chemother.* **43**: 410–412.
- Gmuender, H., Kuratli, K., and Keck, W. 1995. Effect of pyrimido[1,6-a]benzimidazoles, quinolones, and Ca<sup>2+</sup> on the DNA gyrase-mediated cleavage reaction. *Antimicrob. Agents Chemother.* **39**: 163–169.
- . 1997. In the presence of subunit A inhibitors DNA gyrase cleaves DNA fragments as short as 20 bp at specific sites. *Nucleic Acids Res.* **25**: 604–610.
- Goetschi, E., Angehrn, P., Gmuender, H., Hebeisen, P., Link, H., Masciadri, R., and Nielsen, J. 1993. Cyclothialidine and its congeners: A new class of DNA gyrase inhibitors. *Pharmacol. Ther.* **60**: 367–380.
- Gormley, N.A., Orphanides, G., Meyer, A., Cullis, P.M., and Maxwell, A. 1996. The interaction of coumarin antibiotics with fragments of the DNA gyrase B protein. *Biochemistry* **35**: 5083–5092.
- Gygi, S.P., Rochon, Y., Franza, B.R., and Aebersold, R. 1999. Correlation between protein and mRNA abundance in yeast. *Mol. Cell. Biol.* **19**: 1720–1730.
- Hooper, D.C. 1993. Quinolone mode of action—New aspects. *Drugs* **3**: 8–14.
- Hooper, D.C. and Wolfson, J.S. 1991. Mode of action of the new quinolones: New data. *Eur. J. Clin. Microbiol. Infect. Dis.* **10**: 223–231.
- Hubschwerlen, C., Pflieger, P., Specklin, J.L., Gubernator, K., Gmuender, H., Angehrn, P., and Kompis, I. 1992. Pyrimido[1,6-a]benzimidazoles: A new class of DNA gyrase inhibitors. *J. Med. Chem.* **35**: 1385–1392.
- Jovanovich, S.B. and Lebowitz, J. 1987. Estimation of the effect of coumermycin A1 on *Salmonella typhimurium* promoters by using random operon fusions. *J. Bacteriol.* **169**: 4431–4435.
- Kaji, A., Teyssier, E., and Hirokawa, G. 1998. Disassembly of the post-termination complex and reduction of translational error by ribosome recycling factor (RRF)—A possible new target for antibacterial agents. *Biochem. Biophys. Res. Commun.* **250**: 1–4.
- Kampranis, S.C., Gormley, N.A., Tranter, R., Orphanides, G., and Maxwell, A. 1999. Probing the binding of coumarins and cyclothialidines to DNA gyrase. *Biochemistry* **38**: 1967–1976.
- Langen, H., Takacs, B., Evers, S., Berndt, P., Lahm, H.P., Wipf, B., Gray, C., and Fountoulakis, M. 2000. Two-dimensional map of the proteome of *Haemophilus influenzae*. *Electrophoresis* **21**: 411–429.
- Lewis, R.J., Singh, O.M.P., Smith, C.V., Skarzynski, T., Maxwell, A., Wonacott, A.J., and Wigley, D.B. 1996. The nature of inhibition of DNA gyrase by the coumarins and the cyclothialidines revealed by X-ray crystallography. *EMBO J.* **15**: 1412–1420.
- Lockhart, D.J., Dong, H., Byrne, M.C., Follettie, M.T., Gallo, M.V., Chee, M.S., Mittmann, M., Wang, C., Kobayashi, M., Horton, H., et al. 1996. Expression monitoring by hybridization to high-density oligonucleotide arrays. *Nature* **381**: 1675–1680.
- Luttinger, A. 1995. The twisted 'life' of DNA in the cell: Bacterial topoisomerases. *Mol. Microbiol.* **15**: 601–606.
- Maxwell, A. 1993. The interaction between coumarin drugs and DNA gyrase. *Mol. Microbiol.* **9**: 681–686.
- Maxwell, A. 1999. DNA gyrase as a drug target. *Biochem. Soc. Trans.* **27**: 48–53.
- Menzel, R. and Gellert, M. 1994. The biochemistry and biology of DNA gyrase. In *Adv. Pharmacol.* (ed. L.F. Liu), pp. 39–69. Academic Press, San Diego, CA.
- Munoz, R. and De La Campa, A.G. 1996. ParC subunit of DNA topoisomerase IV of *Streptococcus pneumoniae* is a primary target of fluoroquinolones and cooperates with DNA gyrase A subunit in forming resistance phenotype. *Antimicrob. Agents Chemother.* **40**: 2252–2257.
- Nakada, N., Gmuender, H., Hirata, T., and Arisawa, M. 1994. Mechanism of inhibition of DNA gyrase by cyclothialidine, a novel DNA gyrase inhibitor. *Antimicrob. Agents Chemother.* **38**: 1966–1973.
- . 1995. Characterization of the binding site for cyclothialidine on the B subunit of DNA gyrase. *J. Biol. Chem.* **270**: 14286–14291.
- Oram, M., Dosanjh, B., Gormley, N.A., Smith, G.V., Fisher, L.M., Maxwell, A., and Duncan, K. 1996. Mode of action of GR122222X, a novel inhibitor of bacterial DNA gyrase. *Antimicrob. Agents Chemother.* **40**: 473–476.
- Piddock, L.J.V. and Wise, R. 1987. Induction of the SOS response in *Escherichia coli* by 4-quinolone antimicrobial agents. *FEMS Microbiol. Lett* **41**: 289–294.
- Piddock, L.J.V., Walters, R.N., and Diver, J.M. 1990. Correlation of quinolone MIC and inhibition of DNA, RNA, and protein synthesis and induction of the SOS response in *Escherichia coli*. *Antimicrob. Agents Chemother.* **34**: 2331–2336.
- Reece, R.J. and Maxwell, A. 1991. DNA gyrase: Structure and function. *Crit. Rev. Biochem. Mol. Biol.* **26**: 335–375.
- Roca, J. 1995. The mechanisms of DNA topoisomerases. *Trends Biochem. Sci.* **20**: 156–160.
- Sanchez, J.C., Appel, R.D., Golaz, O., Pasquali, C., Ravier, F., Bairoch, A., and Hochstrasser, D.F. 1995. Inside SWISS 2DPAGE database. *Electrophoresis* **16**: 1131–1151.
- Sharma, A. and Mondragon, A. 1995. DNA topoisomerases. *Curr. Opin. Struct. Biol.* **5**: 39–47.
- Steck, T.R., Franco, R.J., Wang, J.Y., and Drlica, K. 1993. Topoisomerase mutations affect the relative abundance of many *Escherichia coli* proteins. *Mol. Microbiol.* **10**: 473–481.
- Tingey, A.P. and Maxwell, A. 1996. Probing the role of the ATP-operated clamp in the strand-passage reaction of DNA gyrase. *Nucleic Acids Res.* **24**: 4868–4873.
- Walters, R.N., Piddock, L.J.V., and Wise, R. 1989. The effect of mutations in the SOS response on the kinetics of quinolone killing. *J. Antimicrob. Chemother.* **24**: 863–874.
- Wang, J.C. and Lynch, A.S. 1993. Transcription and DNA supercoiling. *Curr. Opin. Genet. Dev.* **3**: 764–768.
- Wodicka, L., Dong, H., Mittmann, M., Ho, M.H., and Lockhart, D.J. 1997. Genome-wide expression monitoring in *Saccharomyces cerevisiae*. *Nature* **381**: 1359–1367.
- Zengel, J.M. and Lindahl, L. 1994. Diverse mechanisms for regulating ribosomal protein synthesis in *Escherichia coli*. *Prog. Nucleic Acid Res. Mol. Biol.* **47**: 331–370.

Received August 1, 2000; accepted in revised form October 12, 2000.