



A Human cDNA Expression Library in Yeast Enriched for Open Reading Frames

Caterina Holz, Angelika Lueking, Lara Bovekamp, et al.

Genome Res. 2001 11: 1730-1735

Access the most recent version at doi:[10.1101/gr.181501](https://doi.org/10.1101/gr.181501)

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

An advertisement banner with a teal background. On the left, the text reads "CRISPR and RNAi Genetic Screening. Your new superpower." In the center, there is a white box with the words "LEARN MORE". On the right, there is a photograph of a woman wearing a red mask and a red cape, and the Cellecta logo, which consists of a cluster of green dots.

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Cold Spring Harbor Laboratory Press

Methods

A Human cDNA Expression Library in Yeast Enriched for Open Reading Frames

Caterina Holz,¹ Angelika Lueking,² Lara Bovekamp,² Claudia Gutjahr,²
Natalia Bolotina,¹ Hans Lehrach,² and Dolores J. Cahill^{2,4}

¹Technical University Berlin, Institute for Biotechnology, D-13355 Berlin, Germany; ²Max Planck Institute of Molecular Genetics, D-14195 Berlin, Germany; ³PROTAGEN, D-44801 Bochum, Germany

We developed a high-throughput technique for the generation of cDNA libraries in the yeast *Saccharomyces cerevisiae* which enables the selection of cloned cDNA inserts containing open reading frames (ORFs). For direct screening of random-primed cDNA libraries, we have constructed a yeast shuttle/expression vector, the so-called ORF vector pYEXTSH3, which allows the enriched growth of protein expression clones. The selection system is based on the *HIS3* marker gene fused to the C terminus of the cDNA insert. The cDNAs cloned in-frame result in histidine prototrophic yeast cells growing on minimal medium, whereas clones bearing the vector without insert or out-of-frame inserts should not grow on this medium. A randomly primed cDNA library from human fetal brain tissue was cloned in this novel vector, and using robot technology the selected clones were arrayed in microtiter plates and were analyzed by sequencing and for protein expression. In the constructed cDNA expression library, about 60% of clones bear an insert in the correct reading frame. In comparison to unselected libraries it was possible to increase the clones with inserts in the correct reading frame more than fourfold, from 14% to 60%. With the expression system described here, we could avoid time-consuming and costly techniques for identification of clones expressing protein by using antibody screening on high-density filters and subsequently rearranging the selected clones in a new "daughter" library. The advantage of this ORF vector is that, in a one-step screening procedure, it allows the generation of expression libraries enriched for clones with correct reading frames as sources of recombinant proteins.

In recent years, the human genome sequencing projects have generated enormous amounts of sequence data. With the human genome sequence now available, the challenge of understanding the function of the newly discovered genes has to be addressed. High-throughput technologies have been developed that allow the monitoring of gene activity on the transcriptional level by analysis of complex expression patterns of a specific tissue (Harrison et al. 1995; Schena et al. 1996, 1998; Perret et al. 1998). The next step is the profiling of protein products encoded by expressed cDNA clones to obtain more information on their regulation, biochemical function, and potential interaction partners. This requires the simultaneous expression of protein from a large number of cDNA clones, which has been performed in the *Escherichia coli* bacterial system (Büssow et al. 1998).

Using robot technology, a human fetal brain cDNA expression library was screened for clonal protein expression in high-throughput on automatically gridded high-density protein arrays. Because the unselected frequency of in-frame clones in libraries with random orientation is statistically only 1/6 of all clones, it would be useful to develop a vector that enables the direct selection of open reading frames (ORFs), improving the yield of clones expressing protein. We have developed such a system in which a C-terminally fused marker gene is expressed only if the cloned insert carries no internal stop sequences, which may result from frameshifts or

5' and 3' untranslated regions. Consequently, the use of random-primed cDNAs is required in this system instead of oligo(dT)-tailed cDNAs that carry their own C-terminal termination codon.

Previously, *E. coli* vectors have been constructed on the basis of the β -galactosidase coding sequence for the generation of in-frame fusion libraries (Gray et al. 1987). However, only DNA fragments in the range of 100–1000 bp could be enriched using this system. Moreover, expression of the marker gene was also observed when the LacZ gene was not in frame with the cDNA because of the polycistronic mRNA in prokaryotes and the reinitiation of translation. Davis and Benzer (1997) constructed a vector that confers kanamycin resistance to the host on translation of an insert in the correct reading frame. They made three size-fractionated cDNA libraries in *E. coli*, namely 100–200, 200–300, and 300–500 bp. Only the library that contained small cDNA fragments in the range of 100–300 bp could be enriched to 60%–80% for ORF clones.

For the improvement of the selection efficiency, especially of larger inserts, it would be advantageous to use a eukaryotic host, which, in contrast to *E. coli*, has a codon usage homologous to that of mammalian cells, thus avoiding translation problems and frameshifts. Therefore, we chose the yeast *Saccharomyces cerevisiae* as a eukaryotic host, which is able to produce soluble proteins in large amounts (Romanos et al. 1992). The ORF vector pYEXTSH3 carries the *HIS3* gene, coding imidazol-glycerol-phosphate-dehydrogenase, which enables the selection of ORFs based on histidine prototrophy. Here, we describe the construction of this vector, the development of the selection procedure, and the construction and characterization of a human fetal brain cDNA library in this expression system.

⁴Corresponding author.

E-MAIL cahill@molgen.mpg.de; FAX 49-30-8413-1128.

Article published on-line before print: *Genome Res.*, 10.1101/gr.181501.
Article and publication are at <http://www.genome.org/cgi/doi/10.1101/gr.181501>.




Figure 1 Restriction map of pYEXTSH3. The vector includes the *E. coli* *Amp^r* gene and the yeast selectable markers *URA3* and *HIS3*. The modified MCS (*SalI/NotI*) contains a stop codon in each reading frame. The epitope tags HA and RGS-His₆ were inserted at the N terminus of the MCS.

RESULTS AND DISCUSSION

Construction and Function of pYEXTSH3

The ORF vector pYEXTSH3 is a derivative of the commercially available expression vector pYEXbx (Clontech), including the Cu²⁺-inducible *CUP1* promoter from the yeast metallothionein gene, which induces expression of the encoded protein (Fig. 1). In addition, the vector contains the *E. coli* ampicillin resistance (*Amp^r*) gene for selection in *E. coli* and the yeast selectable markers *URA3* and *LEU2d*. For the improved expression of randomly primed cDNAs, we introduced a translation initiation sequence that was adapted to *S. cerevisiae*. For the detection and purification of expressed gene products, the plasmid includes a tandem epitope tag positioned N-terminal to the multiple cloning site (MCS), consisting of the hemagglutinine epitope (HA) and the oligo histidine domain (RGS-His₆). The modified MCS includes a stop codon in each reading frame and allows the directional cloning (*SalI/NotI*) of cDNAs. For the cleavage of HIS3p from the fusion protein we additionally inserted the recognition sequence (6P) of the PreScission protease. The yeast *HIS3* gene was introduced C-terminal to the MCS (Fig. 2). The translation of the DNA proceeds from the start codon ATG through the tag sequences into the *HIS3* gene. Only if an insert is cloned in the correct reading frame and if it does not contain any stop codon will complete fusion protein be produced (Fig. 2B). In this case, expression of the gene product leads to histidine prototrophy in a *his3* *S. cerevisiae* strain. Clones bearing the plasmid pYEXTSH3, but without inserts (Fig. 2A), remain auxotrophic, as no *HIS3* gene is expressed because of translation stops in all three

reading frames. The yeast cells are also expected to be auxotrophic if they contain a cDNA that is inserted in an incorrect reading frame, leading to internal stop codons and resulting in the termination of translation before reaching the *HIS3* sequence (Fig. 2C). In general, mRNA sequences carry many stop codons in reading frames other than the correct one.

The *S. cerevisiae* expression strain GRF18 (*his3, leu2*) was used for selection following transformation. To check the selection system, three different test inserts of human *GAPDH* were generated by PCR amplification. These *GAPDH* amplicons, which all contained the same N terminus but differed in their C-terminal regions, were cloned via *SalI/NotI* in-frame with the tag sequences and the start codon in pYEXTSH3. In the first case, the *GAPDH* insert was introduced in the correct reading frame relative to the *HIS3* marker (Fig. 2B). The second construct included the stop codon of the *GAPDH* gene (Fig. 2D), and the third amplicon was modified by inserting an additional base pair at the C-terminal end of *GAPDH*, leading to a frameshift (Fig. 2C). These plasmids were transformed into yeast and the expression clones were selected directly on minimal medium (SD-his) supplemented with CuSO₄ to induce protein expression. The vector without insert was used as a negative control (Fig. 2A), and a modified vector (pYEXT2HIS3), in which the stop codons between *SalI* and *NotI* were deleted and which also contains an N-terminally tagged *HIS3* gene, was transformed in yeast as a positive control. As expected, the insertion of the cDNA in-frame resulted in histidine-prototrophic yeast cells growing on minimal medium, whereas the clones containing the vector without insert, or with the complete gene including the stop codon, did not grow. However, we observed that 3% of the selected transformants with the out-of-frame *GAPDH* could grow on minimal medium. This phenomenon might be caused by a translational frameshift and was reduced by increasing selection stringency (from 0.1 to 0.5 mM CuSO₄). The survival ratio and the transformation efficiency of the in-frame construct were improved significantly by reducing the copper




Figure 2 Design and function of the ORF vector pYEXTSH3. (A) pYEXTSH3 includes a tandem epitope tag, a translation start and stop codon, and the *HIS3* marker gene, which is under the control of the *CUP1* promoter. In this plasmid, translation terminates before reaching *HIS3*. (B–D) The cloning of cDNAs insert, represented as XXX_n, results in the following outcomes at the C terminus: (B) If the size of the cDNA insert is a multiple of three and both N and C termini are cloned in the correct reading frame, *HIS3* is translated. (C) The translation terminates after translation of a few triplets in *HIS3*, if the insert is a multiple of 3 + 1, as the reading frame is shifted. (D) If the cloned cDNA includes an internal stop, there is no translation of *HIS3*.

Table 1. Transformation Efficiency of pYEXTSH3 and pYEXT2HIS3

Vector construct	Concentration of CuSO ₄				
	w/o CuSO ₄	0.05 mM	0.1 mM	0.3 mM	0.5 mM
pYEXT2HIS3	1.2×10^3	1.2×10^3	2×10^4	4.8×10^3	1.2×10^3
(A) pYEXTSH3	0	0	60	2.7×10^2	0
(B) pYEXTGH3 (RF ⁺)	7×10^2	4×10^2	5×10^4	2.8×10^4	2×10^2
(C) pYEXTGnH3 (RF ⁻)	10		3×10^2	10	0

¹Transformation efficiency of pYEXTSH3 constructs (A–C) and the control plasmid pYEXT2HIS3 in *Saccharomyces cerevisiae* GRF18 is shown to be dependent on concentration of CuSO₄. The efficiencies are given in cfu/μg DNA. (RF⁺), insert in frame with *HIS3*. (RF⁻), insert not in frame with *HIS3*.

concentration in the selection medium. These results are shown in Table 1.

Generation of the Yeast Expression Library

To subclone the cDNAs in this vector, two linkers containing *Sall* and *NotI* sites, respectively, were ligated to a randomly primed cDNA library from human fetal brain tissue, resulting in *Sall*–*NotI* overhangs. The library was cloned into pYEXTSH3 and transformed in *S. cerevisiae* strain GRF18. The clones were selected on SD-his agar supplemented with 0.1 mM CuSO₄. In an initial experiment to characterize the library, 2000 colonies were picked by a robot into 384-well plates. To determine whether the library was enriched for cloned sequences representing ORFs, 96 clones were selected randomly and analyzed by PCR according to the method described in Lueking et al. (2000), by sequencing and protein expression. The obtained sequences were used for BLASTN searches against the public databases (NCBI). Fifty-eight clones (60%) were found to contain genes that were expressed and contained an ORF fused in-phase with *HIS3*. A selection of analyzed clones is listed in Table 2. In contrast to unselected libraries, it was possible to increase the percentage of clones in the correct reading frame fourfold, from 14% (Büslow et al. 2000) to 60%.

Following PCR analysis of 96 clones, we determined the average insert size of expressed cDNAs in this library to be 1.6

kb. The size range of the inserts was 200–3000 bp. The frequency of sequences cloned in the correct reading frame seems not to be dependent on the size of the cloned cDNAs, as was seen to be the case for bacterial expression (Davis and Benzer 1997), in which the proportion of in-frame clones decreased with increasing size of the cloned cDNAs, and only inserts with a maximum size of 500 bp were expressed. However, we observed in our yeast expression system that the efficiency of selection for sequences cloned in ORFs is influenced by the selection conditions. On optimization of the selection stringency, we obtained 60% of clones with in-frame inserts. The remaining 40% had either no insert, an internal stop codon, or were in the incorrect reading frame but, nevertheless, showed a histidine prototrophic phenotype.

On sequence analysis of these 40%, 16% had no insert but resulted from in-frame religation of the vector to itself. Fourteen percent of the clones were shown to have internal stops and, in this case, the *HIS3* gene product should not have been produced. Perhaps its expression is the result of a translational frameshift or reinitiation of translation involving interactions between the cloned cDNA sequences and the yeast transcription/translation machinery. Although recognized as playing a key role in the control of a few eukaryotic mRNAs (Hinnebusch 1997), reinitiation has tended not to be regarded as a significant factor in eukaryotic gene expression. Also, the sequence context surrounding the stop codons has a strong

Table 2. ORF Library Clones with Sequence Database Matches in the Correct Reading Frame

Clone	Accession no.	Protein name	Cds/First matched amino acid in database sequence	Predicted protein size (kDa)
001-B01/3	D63997	Golgi autoantigene, GCP170	270-4862/590	63
001-F01/7	NM006031.1	Pericentrin mRNA	1-10018/8720	48
001-H01/9	HSU26396	α-spectrin mRNA	1-2159/1411	28
001-B02/11	AF155821	Protein serine/threonine kinase CPG16	160-1467/717	38
001-A01/2	AB002373	KIAA0375	84-4706/601	147
001-D02/14	Z26634	Akryn 2, neuronal mRNA	1->11775/9237	93
001-F02/16	AB007890	KIAA0430	1-3767/627	115
001-E03/23	NM018013	Protein FLJ10159	1-807/1	29
001-G03/25	NM014787	KIAA0076	87-5183/4023	40
001-D03/22	NM003128.1	Spectrin b, non-erythrocytic 1	311-7405/1893	202
001-G04/33	X79536	HnRNP core protein A1	27-989/1	35
002-D08/63	NM001272.1	Chromodomain DNA helicase	211-6331/5290	38
002-F09/38	HSM802205	DKFZ p434M2...	1-3043/2180	32
002-B12/14	NM007027.1	Topoisomerase II binding protein	341-4648/3103	57
002-D12/28	AB033044	KIAA1208	32-2626/108	92

influence on the efficiency of reinitiation. AU richness in the penultimate upstream codon and in the downstream region has been shown to be correlated with efficient reinitiation (Grant and Hinnebusch 1994). Internal initiation and leaky ribosomal scanning are alternative pathways used instead of the standard scanning mechanism and can be a basis for translational regulation in response to environmental stress (Rhoads and Lamphear 1995; Gan and Rhoads 1996).

The 10% of clones remaining contained inserts that were not in-frame with respect to the downstream fused *HIS3* gene but nevertheless led to a histidine prototrophic phenotype. We suggest that, in these clones, the translation of *HIS3* was caused by a ribosomal frameshift. Different types of frameshifting have been described in *S. cerevisiae* (Farabaugh 1996). They are found most commonly during the translation process, when the fusion of two protein subunits that are not in the same reading frame but belong to the same protein is required. It is not known which characteristics play a role in the frameshifting events in the present experiments. It would be interesting to test different host strains and further selection markers, instead of *HIS3*, to reduce the number of false-positive clones. The application of a dominant resistance marker could possibly improve the screening efficiency for in-frame clones because more stringent selection conditions could be used.

Antibody Screening on Protein Filters

The picked yeast clones were checked additionally for protein expression on high-density protein arrays. The cDNA clones were spotted onto polyvinylidene difluoride (PVDF) membranes that were processed subsequently for protein expression using the monoclonal antibody RGS(His₆) (QIAGEN), which recognizes the N-terminal sequence RGS-His₆ of recombinant expression products (Fig. 3). Approximately 60% of the clones expressed proteins, which were detected by the antibody screening on protein arrays, confirming the results from the sequence analysis of the selected cDNA. For example, the clone in position A2 in the spotting pattern shown in Figure 3 corresponds to the clone 001-F01/7 (Table 2).

Protein Expression

The gene products of some representative clones, which had been sequenced previously, were characterized by protein expression analysis. Whole-cell proteins separated by SDS-PAGE




Figure 3 Protein expression screening of the HFBr library in *S. cerevisiae* GRF18 on protein filters. The protein filters were screened for cDNA clones expressing recombinant fusion proteins using the RGS-His₆ antibody. The clone in position D6 is the negative control containing a vector without an insert, and E6 is the positive control expressing the human *GAPDH* gene.




Figure 4 Expression of different inserts from HFBr cDNA clones in the ORF-selecting vector pYEXTSH3. Proteins separated by SDS-PAGE (100 μ g each lane) were detected from crude cell extracts by immunoblotting with an anti-RGS-His₆ antibody. (Lanes 1–5) *HIS3* fusion proteins from selected cDNA expression clones 001-B2, 001-H1, 001-F1, 001-F2, and 001-E3; (lanes 6–9) *HIS3* from pYEXT2His3, pYEXTSH3 without insert (negative control), pYEXTGnH3 (*GAPDH* not in-frame), and pYEXTGH3 (*GAPDH* in-frame, positive control). The positions of the prestained Bio-Rad broad range marker are indicated on the left.

were transferred to PVDF membranes, followed by detection of expression products by Western immunoblotting with the antibody specific for the N-terminal RGS-His₆ epitope of the expressed fusion proteins. As shown in Figure 4, the *HIS3* gene product was detected at 36 kD (lane 6), whereas no protein was detectable from crude cell lysate of the negative control pYEXTSH3, which does not contain a cDNA insert (lane 7). The gene products of the RGS-His₆-tagged fusion genes, *GAPDH**His3* (construct with frameshift) and *GAPDH*-*HIS3*, are shown in lanes 8 and 9, respectively. The translation of the first construct terminates after 10 amino acids because of the stop codon that was introduced by the reading-frame shift, whereas the latter construct resulted in the expression of a full-length fusion protein with the expected molecular weight of 86 kD. The expression products of five cDNA clones are shown in lanes 1–5, in which all clones express a detectable fusion protein.

In Figure 5 the purification (lanes 2–8) and cleavage (lane 9) of the *GAPDH*-*HIS3* protein is shown as an example. The RGS-His₆ tagged fusion protein was affinity purified under native conditions using Ni-NTA agarose, and the C-terminally fused *HIS3* protein was cleaved after the 6P site by protease digestion. After 16 h incubation at 4°C, the fusion protein was cleaved completely without any degradation products (lane 9).

In summary, the screening system described here is a useful one-step screening procedure for the generation of expression libraries enriched with clones that produce authentic polypeptides. Using the eukaryotic host *S. cerevisiae*, we were able to produce soluble proteins in large amounts that can be used to generate protein arrays and for the analysis of antibody specificity and other protein–protein interaction or ligand–receptor systems (Cahill 2001). A further field of application for our system is to establish protein product catalogs




Figure 5 Purification and cleavage of the GAPDH-HIS3 fusion protein. The fusion protein was expressed in *S. cerevisiae* GRF18 and purified using Ni-NTA agarose. Proteins were visualized by Coomassie staining. (Lane 1) Prestained Bio-Rad broad range marker; (lane 2) cell lysate; (lane 3) flow-through; (lanes 4–6) wash 1–3; (lanes 7,8) eluate 1 and 2; (lane 9) cleavage of the C-terminally fused HIS3 protein by PreScission Protease.

from different tissues and development stages. As these proteins are expressed from arrayed cDNA clones their identity can be checked easily by commonly used DNA techniques or by mass spectrometry (Büssow et al. 2000; Cahill et al. 2000). The cDNA clones can also be used to generate high-density DNA arrays (Clark et al. 1999).

METHODS

Strains, Media, and Transformation

E. coli strains XL1-Blue (Stratagene) and DH10B (Life Technology) were used as host strains for cloning and amplification of plasmid DNA. The transformation of recombinant plasmids was performed by electroporation using the Gene Pulser (Bio-Rad). *E. coli* transformants were selected on LB (0.5% yeast extract, 1% NaCl, 15% bacto-tryptone) medium supplemented with 100 µg/mL ampicillin. The *S. cerevisiae* strain GRF18 (α , *leu2-3*, *leu2-2112*, *his3-15*, *can1*, *mal*) was used as the eukaryotic expression strain. The ORF vector constructs and the cDNA expression library were transformed in *S. cerevisiae* by the lithium acetate method (Gietz et al. 1992), and the resulting transformants were selected on SD-his induction medium (2% dextrose, 0.67% Yeast Nitrogen Base [YNB], 40 mg/L leucine, 0.1 mM CuSO₄). The *S. cerevisiae* clones were cultivated in microtiter plates in freezing medium (2% dextrose, 5% glycerol, 0.67% YNB, 40 mg/L histidine, 0.5 M betaine). For protein expression, the clones were transferred in SD-leu medium (2% dextrose, 0.67% YNB, 40 mg/L histidine) and the expression was induced by addition of 0.5 mM CuSO₄.

Construction of the Vector

The YEX system uses the Cu²⁺-inducible *CUP1* promoter to control expression of genes inserted in the MCS. The expression vector pYEXbx (Clontech) was modified by introduction of a *NotI* restriction site and a translation initiation sequence (CAAATGTCT) downstream of the promoter region, which allows the translation of cDNAs without their own start codon. For detection and purification of the gene product, we introduced a tandem epitope tag positioned N-terminal of the MCS consisting of the hemagglutinine epitope (HA) and the oligo histidine domain (RGS-His₆). A DNA fragment that carries the coding sequence of the *HIS3* gene (SWISS-PROT P06633) and the PreScission protease recognition (6P) sequence (CTGGAAGTTCTGTTCAGGGGCC) at the N terminus was generated by PCR amplification from the plasmid YdpH and cloned *NotI/EcoRI* in the expression vector. The resulting plasmid pYEXTH3 can be used for cloning of

cDNAs with *Sall/NotI* overhangs and for the expression of randomly primed cDNAs. The restriction map of pYEXTH3 is shown in Figure 1.

Cloning of the cDNA Expression Library

The cDNA inserts of a random-primed cDNA library from human fetal brain tissues in the vector pSPORT (provided by B. Korn; Resource Centre of the German Genome Project [RZPD], Heidelberg) were cleaved with *Sall/EagI* and electroporated through 1% agarose gel. Fragments in the size range 200 bp–2 kb were eluted from the gel (QIAGEN, QIAquick Gel extraction Kit) and ligated into the *Sall* and *NotI* sites of the vector. Following transformation in *E. coli* DH10B by electroporation, the library was plated onto square agar plates (230 mm × 230 mm, Genetics) and grown for 20 h at 37°C. Approximately 150,000 clones were scraped from the agar, resuspended in P1 buffer (50 mM Tris-HCl at pH 8.0, 10 mM EDTA, 100 µg RNase A), and plasmid DNA was isolated. The purified DNA was transformed in *S. cerevisiae* GRF18.

Generation of the Yeast Expression Library

Following transformation the yeast cells were plated on SD-his induction medium at a density of 1500 clones per agar plate (230 mm × 230 mm) and were incubated 3–4 d at 28°C. Colonies obtained were picked into 384-well microtiter plates and filled with freezing medium by a robot. The colony-picking program was adapted to yeast cells. The robot-picking specifications are as follows: It has a resolution of 5 µm, a velocity of 3 m/sec, and it is able to pick about 3600 clones/h. The cells were grown in the microtiter wells for 3–4 d at 28°C, and afterward the clones were replicated by a 384-pin replicator. All copies were stored frozen at –80°C.

Sequence Analysis

For sequencing, the *S. cerevisiae* clones were transferred into 96-well plates filled with 65 µL freezing medium and were incubated 4 d at 28°C. The inserts of the cDNA clones were amplified by PCR directly from liquid cultures as described previously (Lueking et al. 2000). For PCR amplification, the primers pYEXbx5 (CATATAGAAGTCATCGA) and pYEXbx3 (TTTGCAG CTACCACATT) were used.

Generation of High-Density Filters

For generation of high-density filters the yeast clones were grown in the 384-well microtiter plates and were gridded onto PVDF membrane (Millipore) for protein analysis as described (Meier-Ewert et al. 1994; Büssow et al. 1998) with the following modifications. Before spotting, the filters were soaked in medium (0.67% YNB, 2% dextrose, 40 mg/L histidine, 0.5% calcium propionate). The clones were gridded in duplicate in a 3 × 3 spotting pattern at a density of 9216 clones per 222 mm × 222 mm PVDF membrane with ink guide dots. The prepared filters were placed onto square SD-leu agar plates containing 0.5% calcium propionate (antifungal agent). Following colony growth for 2 d at 28°C, protein filters were transferred to agar plates supplemented with 0.5 mM CuSO₄ to induce protein expression and were then incubated for an additional 2 d at 28°C. Then protein filters were placed on Whatman 3MM paper soaked in 4 M NaOH and left at room temperature for 1 h. The membranes were transferred onto Whatman 3MM paper soaked in denaturant solution 0.5 M NaOH, 1.5 M NaCl (10 min); neutralizing solution 1 M Tris-HCl (pH 7.5), 1.5 NaCl (10 min); 2× SSC (15 min). Yeast debris was wiped off with paper towels in TBST-T buffer (20 mM Tris-HCl at pH 7.5, 0.5 M NaCl, 0.05% Tween 20, 0.5% Triton X-100) and the membranes were washed two times briefly with TBS before using for antibody screening.

Antibody Screening on Protein Filters

The processed filters were blocked 1 h in blocking buffer (2% [w/v] BSA/TBST) and were incubated for 1–2 h in primary antibody solution diluted (1 : 2000) in blocking buffer according to the manufacturer's instructions. After washing 2×10 min in TBST, filters were incubated in horseradish peroxidase-conjugated secondary antibody diluted in 5% nonfat dry milk powder in TBST for 1 h. After washing four times for 10 min in TBST buffer, the filters were incubated in chemiluminescence substrate (NEN, Renaissance) for 1 min at room temperature. The excess chemiluminescence reagent was removed and the membranes either were exposed to autoradiography film, or a video system and a BioImager (428 nm) were used to generate pictures of filters, respectively.

Protein Expression

Selected clones of *S. cerevisiae* GRF18 were grown in 20–100 mL SD-leu medium at OD₆₀₀ of 1.0, and protein expression was induced by addition of 0.5 mM CuSO₄ for 3–4 h. The cells were harvested by centrifugation (10 min, 2500g, 4°C) and washed once with PBS (10 mM Na₂PO₄, 300 mM NaCl at pH 8.0). Washed pellets were resuspended in lysis buffer (10 mM Na₂PO₄, 300 mM NaCl at pH 8.0, 1% (v/v) Triton X-100) with protease inhibitor PMSF (1 mM). One volume sterile, acid-washed glass beads (0.5 mm) were added and cells were lysed by seven cycles of 1 min full-speed vortexing followed by 1 min cooling on ice. Beads were removed and the extract was cleared by centrifugation (15 min, 10,000g). Whole-cell proteins in the supernatant were separated directly by SDS-PAGE or purified by affinity chromatography (Ni-NTA agarose). Gels were blotted onto PVDF membranes, blocked with 2% BSA/TBST, and incubated for 1–2 h with the primary antibody diluted in blocking buffer according to the manufacturer's instructions. The secondary antibody was horseradish peroxidase-conjugated anti-mouse antibody for use with the NEN chemiluminescence protocol.

Protease Cleavage

The HIS3 fusion proteins were cleaved by using 2 units of PreScission Protease (Amersham Pharmacia Biotech) for each 100 µg of fusion protein in the eluate. The sample was adjusted to 1× cleavage buffer by addition of an appropriate volume of 10× cleavage buffer according to the manufacturer's instructions and was incubated for 4 h at 5°C.

ACKNOWLEDGMENTS

We thank Bernhard Korn (RZPD, Heidelberg) for the human fetal brain cDNA, Christine Lang (Technical University, Berlin) for the YdpH plasmid carrying the *HIS3* gene and the *S. cerevisiae* strain GRF18, Martin Horn for assistance of adaptation of the picking procedure to yeast cells, and Heike Göhler and Erich E. Wanker for valuable discussions. This work was funded by Bundesministerium für Bildung und Forschung (BMBF Grants 0311018, 0311323, and 0311870) and the Max-Planck-Gesellschaft.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

REFERENCES

- Büssow, K., Cahill, D., Nietfeld, W., Bancroft, D., Scherzinger, E., Lehrach, H., and Walter, G. 1998. A method for global protein expression and antibody screening on high-density filters of an arrayed cDNA library. *Nucleic Acids Res.* **26**: 5007–5008.
- Büssow, K., Nordhoff, E., Lübbert, C., Lehrach, H., and Walter, G. 2000. A human cDNA library for high-throughput protein expression screening. *Genomics* **65**: 1–8.
- Cahill, D.J. 2001. Protein and antibody arrays and their medical applications. *J. Immunol. Meth.* **250**: 81–91.
- Cahill, D.J., Nordhoff, E., O'Brien, J., Klose, J., Eickhoff, H., and Lehrach, H. 2000. Bridging genomics and proteomics. In *Proteomics* (eds. S. Pennington and M. Dunn), pp. 1–20. BIOS Scientific Publishers, Oxford.
- Clark, M.D., Panopoulou, G.D., Cahill, D.J., Buessow, K., and Lehrach, H. 1999. Construction and analysis of arrayed cDNA libraries. *Meth. Enzymol.* **303**: 205–33.
- Davis, C.A. and Benzer, S. 1997. Generation of cDNA expression libraries enriched for in-frame sequences. *Proc. Natl. Acad. Sci.* **94**: 2128–2132.
- Farabaugh, P.J. 1996. Programmed translational frameshifting. *Microbiol. Rev.* **60**: 103–134.
- Gan, W. and Rhoads, R.E. 1996. Internal initiation of translation directed by the 5'-untranslated region of the mRNA for eIF4G, a factor involved in the picornavirus-induced switch from cap-dependent to internal initiation. *J. Biol. Chem.* **271**: 623–626.
- Gietz, D., St. Jean, A., Woods, R.A., and Schiestl, R.H. 1992. Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* **20**: 1425.
- Grant, C.M. and Hinnebusch, A.G. 1994. Effect of sequence context at stop codons on efficiency of reinitiation in GCN4 translational control. *Mol. Cell. Biol.* **14**: 606–618.
- Gray, M.R., Mazzara, G.P., Reddy, P., and Rosbash, M. 1987. Searching for clones with open reading frames. *Meth. Enzymol.* **154**: 129–156.
- Harrison, S.M., Dunwoodie, S.L., Arkell, R.M., Lehrach, H., and Beddington, R.S.P. 1995. Isolation of novel tissue-specific genes from cDNA libraries representing the individual tissue constituents of the gastrulating mouse embryo. *Development* **121**: 2479–2489.
- Hinnebusch, A.G. 1997. Translational regulation of yeast GCN4. A window on factors that control initiator-trna binding to the ribosome. *J. Biol. Chem.* **272**: 21661–21664.
- Lueking, A., Holz, C., Gotthold, C., Lehrach, H., and Cahill, D.J. 2000. A system for dual protein expression in *Pichia pastoris* and *Escherichia coli*. *Protein Expr. Purif.* **20**: 372–378.
- Meier-Ewert, S., Rothe, J., Mott, R., and Lehrach, H. 1994. Establishing catalogues of expressed sequences by oligonucleotide fingerprinting of cDNA libraries. In *Identification of transcribed sequences* (eds. U. Hochgeschwender and K. Gardiner), pp. 253–260. Plenum, New York.
- Perret, E., Ferran, E.A., Marinx, O., Liauzun, P., Dumont, X., Fournier, J., Kaghad, M., Ferrara, P., and Caput, D. 1998. Improved differential screening approach to analyse transcriptional variations in organized cDNA libraries. *Gene* **208**: 103–115.
- Rhoads, R.E. and Lamphear, B.J. 1995. Cap-independent translation of heat shock messenger RNAs. *Curr. Top. Microbiol. Immunol.* **203**: 131–153.
- Romanos, M.A., Scorer, C.A., and Clare, J.J. 1992. Foreign gene expression in yeast: A review. *Yeast* **8**: 423–488.
- Schena, M., Shalon, D., Heller, R., Chai, A., Brown, P.O., and Davis, R.W. 1996. Parallel human genome analysis: Microarray-based expression monitoring of 1000 genes. *Proc. Natl. Acad. Sci.* **93**: 10614–10619.
- Schena, M., Heller, R.A., Thieriault, T.P., Konrad, K., Lachenmeier, E., and Davis, R.W. 1998. Microarrays: Biotechnology's discovery platform for functional genomics. *Trends Biotechnol.* **16**: 301–306.

Received January 23, 2001; accepted in revised form July 5, 2001.