



## A Novel $\gamma$ 2-Herpesvirus of the Rhadinovirus 2 Lineage in Chimpanzees

Vincent Lacoste, Philippe Maucière, Guy Dubreuil, et al.

*Genome Res.* 2001 11: 1511-1519

Access the most recent version at doi:[10.1101/gr.158601](https://doi.org/10.1101/gr.158601)

---

**References** This article cites 28 articles, 14 of which can be accessed free at:  
<http://genome.cshlp.org/content/11/9/1511.full.html#ref-list-1>

### License

**Email Alerting Service** Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

---

To subscribe to *Genome Research* go to:  
<https://genome.cshlp.org/subscriptions>

---

Cold Spring Harbor Laboratory Press

# A Novel $\gamma$ 2-Herpesvirus of the Rhadinovirus 2 Lineage in Chimpanzees

Vincent Lacoste,<sup>1</sup> Philippe Maucière,<sup>1,2</sup> Guy Dubreuil,<sup>3</sup> John Lewis,<sup>4</sup>  
Marie-Claude Georges-Courbot,<sup>3</sup> and Antoine Gessain<sup>1,5</sup>

<sup>1</sup>Unité d'Epidémiologie et Physiopathologie des Virus Oncogènes, Département du SIDA et des Rétrovirus, Institut Pasteur, 75724 Paris Cedex 15, France; <sup>2</sup>Centre Pasteur du Cameroun, BP 1274, Yaoundé, Cameroon; <sup>3</sup>Centre International de Recherches Médicales, Franceville, Gabon; <sup>4</sup>International Zoo Veterinary Group, Keighley, West Yorkshire BD21 1AG, UK

Old World monkeys and, recently, African great apes have been shown, by serology and polymerase chain reaction (PCR), to harbor different  $\gamma$ 2-herpesviruses closely related to Kaposi's sarcoma-associated Herpesvirus (KSHV). Although the presence of two distinct lineages of KSHV-like rhadinoviruses, RV1 and RV2, has been revealed in Old World primates (including African green monkeys, macaques, and, recently, mandrills), viruses belonging to the RV2 genogroup have not yet been identified from great apes. Indeed, the three yet known  $\gamma$ 2-herpesviruses in chimpanzees (PanRHV1a/PtRV1, PanRHV1b) and gorillas (GorRHV1) belong to the RV1 group. To investigate the putative existence of a new RV2 *Rhadinovirus* in chimpanzees and gorillas we have used the degenerate consensus primer PCR strategy for the Herpesviral DNA polymerase gene on 40 wild-caught animals. This study led to the discovery, in common chimpanzees, of a novel  $\gamma$ 2-herpesvirus belonging to the RV2 genogroup, termed Pan Rhadino-herpesvirus 2 (PanRHV2). Use of specific primers and internal oligonucleotide probes demonstrated the presence of this novel  $\gamma$ 2-herpesvirus in three wild-caught animals. Comparison of a 1092-bp fragment of the DNA polymerase obtained from these three animals of the *Pan troglodytes troglodytes* subspecies, one from Gabon and the two others from Cameroon, revealed <1% of nucleotide divergence. The geographic colocalization as well as the phylogenetic "relationship" of the human and simian  $\gamma$ 2-herpesviruses support the model according to which herpesviruses have diversified from a common ancestor in a manner mediating cospeciation of herpesviruses with their host species. By demonstrating the existence of two distinct *Rhadinovirus* lineages in common chimpanzees, our finding indicates the possible existence of a novel human  $\gamma$ 2-herpesvirus belonging to the RV2 genogroup.

[The Herpesviral DNA polymerase sequence data determined herein have been deposited at the GenBank database under accession nos. AF290601, AF346488, AF346489, and AF346490.]

The members of the family Herpesviridae have been grouped into three subfamilies, designated Alphaherpesvirinae, Betaherpesvirinae, and Gammaherpesvirinae (Roizmann et al. 1992). Herpesviruses are widespread in vertebrate species, sharing several moderately to well conserved genes, as determined from amino acid identity comparisons (e.g., DNA polymerase and glycoprotein B). Among Gammaherpesvirinae, Epstein-Barr virus (EBV) and Kaposi's sarcoma-associated Herpesvirus (KSHV), also named Human herpesvirus 8 (HHV8), are the human prototypes of the *Lymphocryptovirus* genus and the *Rhadinovirus* genus, respectively. Both of these viruses play a critical role in human multistep carcinogenesis, especially in immunodeficiency patients, leading to Burkitt's lymphoma (Magrath and Judde 1996) and Kaposi's sarcoma (KS) (Chang et al. 1994; Schulz 1998), respectively. Rhadinoviruses, or  $\gamma$ 2-herpesviruses, have also been found in several animal species including New World monkeys (*Herpesvirus ateles* and *Herpesvirus saïmiri*) (Albrecht and Fleckenstein 1990; Albrecht 2000) and Old World monkeys (macaques, African green monkeys, and recently mandrills) (Desrosiers et al. 1997; Rose et al. 1997; Auerbach et al. 2000; Greensill et al. 2000b; Lacoste et

al. 2000c; Strand et al. 2000). Comparison and phylogenetic analyses of available sequences support the existence of two distinct genogroups among the Old World monkey rhadinoviruses, called RV1 and RV2 for *Rhadinovirus* genogroups 1 and 2 (Bosch et al. 1998; Greensill et al. 2000b; Lacoste et al. 2000c; Schultz et al. 2000). KSHV belongs to the RV1 genogroup, whereas no human virus has yet been discovered in the RV2 group.

Considering that KSHV and Kaposi's sarcoma are highly endemic in Central Africa (Schulz 1998; Gessain et al. 1999) and no  $\gamma$ 2-herpesvirus sequence has been described in great apes (Sinkovics and Horvath 1999), the closest primate species to human in the animal kingdom, we decided to investigate the potential presence of KSHV-related viruses in chimpanzees and gorillas from Central Africa. Accordingly, we recently reported the detection and molecular characterization of the DNA polymerase gene fragment of three novel  $\gamma$ 2-herpesviruses in these great apes (Lacoste et al. 2000b). These three new and different rhadinoviruses, two present in *Pan troglodytes* (PanRHV1a and PanRHV1b) and the latest in *Gorilla gorilla* (GorRHV1), were more closely related to KSHV (70%–85% identity at the nucleotide level) than any other previously described virus of this genus. These three novel  $\gamma$ 2-herpesviruses belong to the RV1 genogroup as determined by phylogenetic analyses (Lacoste et al. 2000b). Moreover, an

## <sup>5</sup>Corresponding author.

E-MAIL [agessain@pasteur.fr](mailto:agessain@pasteur.fr); FAX 33 0 140-61-34-65.

Article and publication are at <http://www.genome.org/cgi/doi/10.1101/gr.158601>.

**Table 1. Epidemiological Data and Serological STLV-1/SIV and KSHV Results**

Name	Genus	KSHV serology titer	STLV-1/SIV serology	PCR and specific hybridization				Subspecies determination	Accession nos.
				PanRHV2	PanRHV1a	PanRHV1b	GorRHV1		
PanCamCar	<i>Pan troglodytes</i>	1/80	-/-	-	-	+	-	<i>vellerosus</i>	
PanCamChe	<i>Pan troglodytes</i>	1/320	-/-	-	-	-	-	<i>troglodytes</i>	AF346488
PanCamDja	<i>Pan troglodytes</i>	1/40	-/+	+	-	-	-	<i>troglodytes</i>	
PanCamEko	<i>Pan troglodytes</i>	1/80	-/-	-	-	+	-	<i>troglodytes</i>	
PanCamEpp	<i>Pan troglodytes</i>	-	-/-	-	-	-	-	<i>troglodytes</i>	AF346489
PanCamEtr	<i>Pan troglodytes</i>	-	-/-	+	-	-	-	<i>troglodytes</i>	
PanCamEtu	<i>Pan troglodytes</i>	-	-/-	-	-	-	-	<i>vellerosus</i>	
PanCamEwa	<i>Pan troglodytes</i>	1/80	-/-	-	+	-	-	<i>vellerosus</i>	
PanCamJac	<i>Pan troglodytes</i>	1/80	-/+	-	-	+	-	<i>vellerosus</i>	
PanCamJab	<i>Pan troglodytes</i>	1/320	-/-	-	-	+	-	<i>vellerosus</i>	
PanCamLou	<i>Pan troglodytes</i>	1/160	-/-	-	-	-	-	<i>troglodytes</i>	
PanCamMac	<i>Pan troglodytes</i>	1/160	-/-	-	+	-	-	<i>troglodytes</i>	
PanCamMay	<i>Pan troglodytes</i>	-	-/-	-	+	-	-	<i>troglodytes</i>	
PanCamMok	<i>Pan troglodytes</i>	1/160	-/-	-	-	-	-	<i>vellerosus</i>	
PanCamMuc	<i>Pan troglodytes</i>	1/160	-/-	-	+	-	-	<i>vellerosus</i>	
PanCamPern	<i>Pan troglodytes</i>	1/320	-/-	-	-	-	-	<i>vellerosus</i>	
PanCamPol	<i>Pan troglodytes</i>	1/80	-/-	-	-	-	-	<i>troglodytes</i>	
PanCamSam	<i>Pan troglodytes</i>	1/640	-/-	-	-	+	-	<i>troglodytes</i>	
PanCamSek	<i>Pan troglodytes</i>	1/80	-/-	-	+	-	-	<i>troglodytes</i>	
PanCamSus	<i>Pan troglodytes</i>	1/80	-/-	-	-	-	-	<i>troglodytes</i>	
PanCamTal	<i>Pan troglodytes</i>	-	-/-	-	-	-	-	<i>vellerosus</i>	
PanCamWan	<i>Pan troglodytes</i>	1/80	-/-	-	+	-	-	<i>vellerosus</i>	
PanGabNte	<i>Pan troglodytes</i>	1/160	ind./-	-	-	-	-	<i>troglodytes</i>	
PanGabNto	<i>Pan troglodytes</i>	1/80	ind./-	-	-	+	-	<i>troglodytes</i>	
PanGabBel	<i>Pan troglodytes</i>	1/320	ind./-	+	-	-	-	<i>troglodytes</i>	AF290601, AF346490
GorGabOmo	<i>Gorilla gorilla</i>	1/160	ind./-	-	-	-	+	<i>gorilla</i>	
GorGabCol	<i>Gorilla gorilla</i>	1/40	ind./-	-	-	-	-	<i>gorilla</i>	
GorCamNya	<i>Gorilla gorilla</i>	-	-/-	-	-	-	-	<i>gorilla</i>	
GorCamEla	<i>Gorilla gorilla</i>	-	-/-	-	-	-	-	<i>gorilla</i>	
GorCamEvi	<i>Gorilla gorilla</i>	-	-/-	-	-	-	-	<i>gorilla</i>	
France	<i>Gorilla gorilla</i>	1/80	-/-	-	-	-	-	<i>gorilla</i>	

Epidemiological data include name, genus, and geographical origin. STLV-1 serology was determined by Immunofluorescence Assay (IFA) and Western blot, SIV serology by specific Enzyme-Linked Immunosorbent Assay (ELISA) and Western blot confirmation, and KSHV serology by IFA assay at a dilution of 1/40. Distribution of the different novel herpesviruses, in the 31 chimpanzees and gorillas for which DNA was available, was determined by specific oligonucleotide probe hybridization on the YVGA-GD1D1B nPCR products (and on Pp2is-Ppas nPCR products for PanRHV2). Shaded boxes indicate viruses for which herpesviral DNA polymerase GenBank accession numbers are available.

**Table 2.** Sequences of Oligonucleotide Primers and Probes Used for Herpesviral DNA Polymerase Consensus and Specific PCRs, Southern Blot Hybridizations and mtDNA PCR Amplification

Oligonucleotide	Orientation	5'–3' sequence
DNA pol degenerate primers		
DFASA <sup>a</sup>	+	GTG TTC GAC TTY GCN AGY YTN TAY CC
VYGA <sup>a</sup>	+	ACG TGC AAC GCG GTG TAY GGN KTN ACN GG
GDTD1B <sup>a</sup>	–	CGG CAT GCG ACA AAC ACG GAG TCN GTR TCN CCR TA
P2s	+	GAG TTT CCK TCG GAR TAY GAC ATG
PanRHV2 specific primers		
Pan2as	–	TCA CTT AAT GCG GTT GGA TCT AG
P2eas	–	CAT TGC GTG TGA CCT TGA TG
P2ias	–	TCT GTT TTC GCT TGC TCA AC
Pp2es	+	GCA TCT TTA TCA TAC GCT AAC GG
Pp2is	+	CGT TTG TTC TCA GTG GAG GAA AG
Pp2as	–	ATG GTG CGT CCC TGG AGC
Specific probes		
PanRHV2-1	+	CTG CAT TAC CTG TTG TCC TAA CGC C
PanRHV2-2	+	ATA TGG TTT TAC TGG AGT AGC CAA CG
PanRHV1a	+	GTG GGT CTA CTG CGG AAG CCT ATA AAC GTC TC
PanRHV1b	+	TCT GTG CTT CTA CAT AGA CCA ATT GAG ACA CA
GorRHV1	+	GCG AGT CTC CTG CAA AAG CCC ATA GAC GTT CCC A
mtDNA primers		
MTD1S	+	CAC CAT TAG CAC CCA AAG CT
MTD1AS	–	CCT GAA GTA GGA ACC AGA TG

<sup>a</sup>Degenerate oligonucleotide primers described (Rose et al. 1997).

+, sense; –, antisense.

Positions of degeneracy are given. N = A, C, G and T; Y = C and T; R = A and G; K = G and T.

independent confirmation of the presence of PanRHV1a (named P<sub>TRV1</sub>) in a colony of captive common chimpanzees (*Pan troglodytes troglodytes*) was published recently (Greensill and Schulz 2000; Greensill et al. 2000a). The goals of the present study were therefore to search for other  $\gamma$ 2-herpesviruses belonging especially to the RV2 genogroup in African great apes, chimpanzees, and gorillas, and to study the prevalence and the species specificity of such identified novel herpesviruses in wild-caught great apes from Central Africa.


## RESULTS

To look for the presence of KSHV-related viruses in great apes, we first performed a serological analysis followed by a PCR-based study on the peripheral blood mononuclear cells (PBMCs) DNA. The plasma of 40 animals, 28 chimpanzees and 12 gorillas, mostly wild-caught and originating from the Western part of Central Africa, were tested by an immunofluorescence assay (IFA) that detects both latent and lytic KSHV antigens (Chatlynne et al. 1998). The results (Table 1) demonstrated a clear fluorescent reactivity to the KSHV antigens-producing cells (KS-1) in the plasma of 22/28 chimpanzees and 6/12 gorillas, with antibody titers ranging from 1/40 (initial dilution) to 1/640.

We then attempted to amplify a fragment of the very conserved Herpesvirus DNA polymerase gene from the PBMCs DNA of 31 out of the 40 great apes, by nested PCR with degenerate primers (Rose et al. 1997). Twenty DNA samples scored positive on the EtBr gel after the nested PCR. By using Southern blot analysis and specific probes for the recently described ape rhadinoviruses (PanRHV1a, PanRHV1b, and GorRHV1; Tables 1 and 2), we observed that 7 of the 31 studied animals were infected by a PanRHV1a, 6 by a PanRHV1b, and 1 by GorRHV1. Cloning and sequencing of two nested PCR products that did not hybridize with such specific viral probes revealed the presence of a novel Gamma-

herpesviral sequence. The 172-bp sequences (excluding primers) were identical to each other and exhibited 55%, 56%, 54%, and 59% nucleotide identity with the corresponding KSHV, PanRHV1a, PanRHV1b, and GorRHV1 fragments, respectively. Using the same nested PCR approach with, however, a specific reverse primer for the second PCR (Pan2as, designed from this novel sequence), we then amplified a second overlapping fragment of ~400 bp (DFASA-Pan2as) from 3 of the 31 animals and obtained finally a 476-bp fragment of the DNA polymerase gene for two chimpanzees and a 400-bp fragment for one other (Table 2; Fig. 1). Another heminested PCR, using P2s (a new degenerate forward primer based on conserved amino acid motifs within the DNA polymerase of  $\gamma$ 2-herpesviruses), P2eas, and P2ias (new virus-specific reverse primers designed from the DFASA-GDTD1B sequence; Table 2) yielded a further 870 bp of viral sequence. The resulting sequences were assembled to give a total of 1168 bp (excluding primers) of PanRHV2 for two chimpanzees and 1092 bp for one other.

Database searches using the BLAST Web server demonstrated that these novel sequences were most similar to the DNA polymerases of the  $\gamma$ 2-Herpesvirus subfamily. Comprehensive comparative analysis of these novel sequences with all the other available related Herpesviral sequences (Table 3; Figs. 2 and 3) revealed the existence of a novel and distinct chimpanzee KSHV-like viral strain that we propose to name PanRHV2 for Pan Rhadino-herpesvirus 2. This viral strain exhibited 63% identity at the nucleotide level and 73% identity at the amino acid level with KSHV in comparing a 454-bp DNA *pol* fragment (Table 3). Comparison of the 364 encoded amino acid sequences showed that the two Cameroones PanRHV2 strains were 99.7% identical to each other, and the Gabonese viral strain compared to the two Cameroones strains presents 99.2% and 99.5% amino acid identity, respectively.



**Figure 1** Relative position and orientation of primers and probes used for consensus and virus-specific DNA polymerase PCR and for Southern blot hybridization. Primers *above* the Herpesviral DNA polymerase gene (ORF 9) sequence represent the initial herpesvirus degenerate primers used in an nPCR assay to identify novel DNA polymerase sequences as well as the P2s degenerate primer. Primers *below* the sequence represent specific primers used in a degenerate (DFASA or P2s)–nondegenerate nPCR assay used to amplify the upstream DNA polymerase sequences as well as specific primers used in an nPCR assay (Pp2es–Pp2as followed by Pp2is–Pp2as) to study the PanRHHV2 prevalence. Relative positions of the specific oligonucleotide probes used for Southern blot hybridization on VYGA–GDTD1B nPCR products (PanRHHV2-1, PanRHHV1a, PanRHHV1b, and GorRHHV1) and also of PanRHHV2-2 for hybridization on Pp2is–Pp2as nPCR products are shown. The sequences of the oligonucleotides are given in Table 2.

Phylogenetic analyses using different methods (Neighbor Joining, DNA Maximum Parsimony) clearly placed this novel chimpanzee virus (PanRHHV2) within the *Rhadinivirus* genus (Fig. 2). Significantly, even though only partial fragments of the DNA polymerases were studied, the phylogenetic analysis presented in Figure 2 is in close agreement with the known clustering of Herpesviruses into  $\alpha$ ,  $\beta$ , and  $\gamma$  subfamilies. Moreover, phylogenetic analyses performed on the 1168-bp DNA polymerase gene fragment provide an identical tree topology (Table 3; data not shown). PanRHHV2 clusters with the macaque (RRV, MneRV2, MGVMn, MGVMm, and MGVMf), the African green monkey (ChRV2), and the recently reported mandrill (MndRHHV2) viral strains in the RV2 genogroup.

A second analysis restricted only to all the available primate *Rhadinivirus* polymerase genes (Fig. 3) demonstrated the existence of three major distinct separate lineages, supported by high bootstrap values among  $\gamma$ 2-herpesviruses. The first corresponds to the RV1 genogroup that comprises the human (KSHV), the chimpanzee (PanRHHV1a and PanRHHV1b), the gorilla (GorRHHV1), the mandrill (MndRHHV1) as well as the macaque (RFHVMm and RFHVMn), and the African green monkey (ChRV1) strains. This main lineage could be separated into three sublineages, each one being well supported (bootstrap values >75%). Among them, we can distinguish the lineage that contains only the Herpesviral strains from Old World monkeys (macaques, mandrills, and African green monkeys) from the two others constituted by Herpesviral strains of apes (gorilla, pan, and homo). The second main lineage, that is, the RV2 genogroup, comprises Old World

monkeys strains (ChRV2, RRV, MneRV2, MGVMn, MGVMm, MGVMf, and MndRHHV2) and the only new Herpesviral strain of apes, PanRHHV2, present in common chimpanzees. At least, this RV2 lineage is more closely connected to the third lineage of New World monkey (Spider and Squirrel monkeys) rhadinoviruses than is the RV1 genogroup.

To study the prevalence of PanRHHV2 infection, we hybridized the products of the heminested PCR (VYGA–GDTD1B) with a specific PanRHHV2 oligonucleotide-labeled probe (PanRHHV2-1). Three DNAs scored positive. We also developed a heminested PCR system with specific primers for the new PanRHHV2 polymerase gene (Table 2). Using these primers in a nested PCR assay followed by hybridization of the PCR products with another internal specific oligonucleotide probe (PanRHHV2-2), we detect the same three positive samples among the 31 DNAs.

Significantly, in our series only one case of multiple  $\gamma$ 2-herpesviral infection was observed in a SIVcpz-infected chimpanzee among the 31 great apes we tested. This common chimpanzee, described as a natural host of HIV-1-related viruses (Corbet et al. 2000), is therefore also a natural host for distinct  $\gamma$ 2-herpesviruses belonging to the two *Rhadinivirus* genogroups, PanRHHV1a and PanRHHV2.

To explore whether host-dependent evolution of chimpanzee rhadinoviruses exists, we determined the subspecies identity of the animals from which these novel viruses were derived. Four chimpanzee subspecies with nonoverlapping geographic ranges have been proposed on the basis of genetic differences in mitochondrial DNA sequences (Morin et al. 1994; Gonder et al. 1997; Gagneux et al. 1999). We amplified and sequenced a 498-bp fragment of mitochondrial DNA displacement loop (mtDNA D-loop) for the 15 chimpanzees infected by PanRHHV1a, PanRHHV1b, or PanRHHV2. Comparison of these newly derived mtDNA sequences to representative sequences from the four different chimpanzee subspecies revealed that 9 out of the 15 infected chimpanzees belonged to the *Pan troglodytes troglodytes* subspecies, and the 6 others belonged to *Pan troglodytes vellerosus* (Table 1). Classification of the chimpanzees was unambiguous as their mtDNA sequences fell within well-defined subspecies clusters and was further corroborated by the known geographic origins of these animals (Cameroon and Gabon).

## DISCUSSION

The novel data presented in this paper as well as recently published data (Greensill et al. 2000a; Lacoste et al. 2000b) indicate that chimpanzees and gorillas are natural hosts for at least four novel and distinct gammaherpesviruses belonging to the two known  $\gamma$ 2-herpesvirus lineages. The new  $\gamma$ 2-

**Table 3.** Percent of Nucleotide and Amino Acid Identities between the Novel  $\gamma$ 2-Herpesvirus PanRHV2 and the Other Primate Gammaherpesviruses

Virus	PanRHV2 <sup>a</sup>			
	% Nucleotide and amino acid sequence identity on 454bp		% Nucleotide and amino acid sequence identity on 1168bp	
	Nucleotide	Amino acid	Nucleotide	Amino acid
KSHV	63	73	66	74
PanRHV1a	62	72		
PtRV1 <sup>b</sup>			67	75
PanRHV1b	64	71		
GorRHV1	63	70		
RFHVMn	62	70	66	73
RFHVMm	63	68	67	73
MndRHV1	64	69		
ChRV1	60	68	66	73
ChRV2	66	76		
MndRHV2	67	74		
MneRV2	67	76	71	79
MGVMn	67	76		
MGVMf	67	75		
MGVMm	67	74		
RRV	67	74	71	78
HVS	60	68	64	71
HVA3	60	66	64	70
EBV	57	59	61	60

<sup>a</sup>PanRHV2 strain accession no. AF346490.<sup>b</sup>The comparison with the PtRV1 strain was performed only on the common 967-bp fragment.

herpesvirus PanRHV2 described in this report corresponds to the first strain of the RV2 genogroup identified in great apes. In contrast, the three other DNA sequences previously detected in chimpanzees and gorillas (PanRHV1a/PtRV1, PanRHV1b, and GorRHV1) belonging to the RV1 genogroup are the closest known homologs to KSHV.

In instances in which particular animal species (macaques, Mandrills, and African green monkeys) have been thoroughly analyzed for the presence of  $\gamma$ 2-herpesviruses, at least two distinct viruses, each one belonging to a particular genogroup RV1 or RV2, have been identified (Desrosiers et al. 1997; Rose et al. 1997; Auerbach et al. 2000; Greensill et al. 2000b; Lacoste et al. 2000c; Schultz et al. 2000). To date, therefore, five distinct macaque  $\gamma$ 2-herpesviruses have been characterized, two in *Macaca mulatta* (RFHVMm and RRV/MGVMm), two in *Macaca nemestrina* (RFHVMn and MneRV2/MGVMn), one in *Macaca fascicularis* (MGVMf), and two in African green monkeys (ChRV1 and ChRV2) as well as in mandrills (MndRHV1 and MndRHV2). Among these viruses, RFHVMm, RFHVMn, ChRV1, and MndRHV1 belong to the RV1 genogroup, and RRV, MGVMm, MneRV2, MGVMn, MGVMf, ChRV2, and MndRHV2 belong to RV2. Data obtained here from great apes extend the existence of the two phylogenetically distinct groups of  $\gamma$ 2-herpesviruses to the chimpanzees and demonstrate the existence of sublineages, within RV1 and RV2, of both Old World monkey and ape rhadinoviruses.

Regarding the comparison of the serological results for KSHV with the PCR detection of the new viruses, it is difficult to provide convincing epidemiological findings. This is because of the limited series of animal tested (40 by serology and 31 by nPCR), but also the fact that there is no serological assay specific for any of these new viruses. However, there is an overall good concordance, as seen in Table 1, between the serological results and the presence of PanRHV1a or PanRHV1b because 12 of 13 PCR positive samples were found in KSHV seropositive animals, but only one PCR positive DNA was detected among the 8 seronegative individuals.


We found that among the PanRHV1a-infected chimpanzees, there were 3 *P. t. vellerosus* and 4 *P. t. troglodytes*, and among the PanRHV1b-infected animals, there were 3 *P. t. vellerosus* and 3 *P. t. troglodytes* (Table 1). This indicates that for PanRHV1 viruses there is no specific association between a peculiar virus strain (1a or 1b) and a chimpanzee subspecies. Nevertheless, among the PanRHV2-infected chimpanzees, there were only 3 *Pan t. troglodytes* individuals. These data indicate a possible species specificity for this new virus, but such preliminary findings need to be confirmed on a larger series of animals.

It is worth noting that several of these animals, despite living in close contact in the same enclosure for several months or years, harbor different viruses. For example, PanCamDja and PanCamJac were housed in the same enclosure for 5 yr before being tested and they harbor different gammaherpesviruses. This suggests that viral infection took place before their arrival in the rescue center. Mother-to-offspring transmission is a possibility that has been suggested for KSHV in highly endemic areas of Central and East Africa (Plancoulaine et al. 2000).


Such data, taken as a whole, indicate that Central African great apes constitute an important reservoir of novel  $\gamma$ 2-herpesviruses. Although there are no available data supporting this hypothesis, the close identity of these viruses with their human pathogenic counterpart KSHV, their presence in peripheral blood mononuclear cells, and the high genetic relationship between apes and humans indicate that they are potentially transmissible to humans.

Regarding the diseases associated with  $\gamma$ 2-herpesviruses, RFHVMm and RFHVMn have been identified in retroperitoneal fibromatosis, a vascular fibroproliferative neoplasm with many morphological and histological similarities to Kaposi's sarcoma (Rose et al. 1997). RRV has also been isolated from simian immunodeficiency-virus-infected macaques with lymphoproliferative disorder reminiscent of multicentric Castleman's disease (Searles et al. 1999). Although the four  $\gamma$ 2-herpesviruses found in chimpanzees and gorillas are closely related to their human pathogenic counterpart KSHV, no clinical pathology has yet been identified in association with infection. The question therefore remains as to whether there is any disease associated with these novel herpesviruses in their natural hosts and especially in the case of multiple infection by SIVcpz, PanRHV1a, and PanRHV2 as observed in one wild-caught animal of our series. Followup of both experimentally HIV-infected chimpanzees (Greensill et al. 2000a) and of naturally SIV-infected animals may therefore provide some important clues regarding the physiopathology and natural history of infection by these novel  $\gamma$ 2-herpesviruses. Efforts are ongoing to establish a cell culture system for the propagation and extensive characterization of these viruses, which may allow the comparison with KSHV strains.

In the RV2 genogroup, only one viral sequence PanRHV2



**Figure 2** Phylogenetic tree resulting from analysis of selected 454-bp fragments (primers QAHNA and GDTD1B) (Rose et al. 1997) of the herpesvirus DNA polymerase gene, which is available for all viruses. The phylogeny was derived by the neighbor-joining method applied to pairwise sequence distances calculated using the Kimura two-parameter method (transition/transversion ratio set to 2). Horizontal branch lengths are drawn to scale with the bar indicating 0.1 nucleotide replacements per site. Numbers at each node indicate the percentage of bootstrap samples (out of 100) in which the cluster to the right is supported. Brackets on the right indicate previously defined subfamily and genus Herpesviral classification. Previously published sequences included and their accession numbers are as follows: HHV1/HSV1 (X04771), HHV2/HSV2 (M16321), HHV3/VZV (X04370), HHV4/Epstein-Barr Virus (V01555), HHV5/HCMV (M14709), HHV6A (X83413), HHV7 (U43400), KSHV/HHV8 (U75698, U93872, and AF005477), HVS (M31122), HVA3 (AF083424), PanRHHV1a (AF250879 and AF250880), PanRHHV1b (AF250881 and AF250882), GorRHHV1 (AF250886), MndRHHV1 (AF282943), MndRHHV2 (AF282937–AF282940), MndHVβ (AF282942), MndCMV (AF282941), ChRV1 (AJ251573), ChRV2 (AJ251574), RFHVMn (AF005478), RFHVMm (AF005479), RRV/17577 (AF083501), RRV/H26-95 (AF029302), MneRV2 (AF204167), Macaca  $\gamma$  virus strains *Macaca mulatta* (AF159033), *Macaca fascicularis* (AF159032), and *Macaca nemestrina* (AF159031) (named MGVMm, MGVMf, and MGVMn, respectively), PRV (L24487), BHV1 (Z78205), EHV1 (M86664), GHV2 (L40431), MCMV (U68299), RhHV5 (AF0033184), MHV68 (U97553), BHV4 (AF031811), EHV2 (U20824), BLHV (AF031808), AHV1 (AF005370), and OHV2 (AF031812).



**Figure 3** Neighbor-joining protein distance tree for the 151 amino acid residues encoded by the 454-bp fragment (primers QAHNA and GDTD1B) (Rose et al. 1997) of DNA polymerase. Sequences were aligned by using *ClustalW* and analyzed by using the *PROTDIST* and *NEIGHBOR* programs in *PHYLIP*. One hundred replica samplings were subjected to bootstrap analysis (*SEQBOOT*). The branch lengths are proportional to the evolutionary distance (scale bar) between the taxa. Previously published sequences included and their accession numbers are as follows: HHV6A (X83413), HVS (M31122), HVA3 (AF083424), HHV4/Epstein-Barr Virus (V01555), KSHV/HHV8 (U75698, U93872, and AF005477), PanRHV1a (AF250879 and AF250880), PanRHV1b (AF250881 and AF250882), GorRHV1 (AF250886), MndRHV1 (AF282943), MndRHV2 (AF282937–AF282940), ChRV1 (AJ251573), ChRV2 (AJ251574), RFHVMn (AF005478), RFHVMm (AF005479), RRV/17577 (AF083501), RRV/H26-95 (AF029302), MneRV2 (AF204167), *Macaca*  $\gamma$  virus strains *Macaca mulatta* (AF159033), *Macaca fascicularis* (AF159032), and *Macaca nemestrina* (AF159031) (named MGVMm, MGVMf, and MGVMn, respectively).

has yet been identified among great apes. This sequence branches off alone in the RV2 group, independently of the Old World monkey viral strains, indicating that this sequence may represent the prototype strain of a great ape lineage within this group. These comparative data, obtained for all the nonhuman primate species, raise the possibility of the existence of another  $\gamma$ 2-herpesvirus, belonging to the RV2 lineage, in humans, in which only KSHV, belonging to the RV1 genogroup, has been identified to date. The identification of novel RV2  $\gamma$ 2-herpesvirus sequences in other nonhuman primate species and the generation of new consensus degenerate primers targeted to the Herpesviral DNA polymerase may be helpful in the detection and identification of this putative human RV2 herpesvirus. Furthermore, the use of degenerate and consensus primers derived from all the primate  $\gamma$ -herpesvirus polymerase genes, including the four novel ones recently described, will allow us to understand the full extent of the natural infection by these viruses among great apes and the frequency of the eventual zoonotic transmission to humans. This virus hunt could be initially focused on captive or free living great apes and on persons at high risk through contact with such animals, including personnel of zoos and animal centers, as well as hunters and their relatives in Central Africa.

The dogma has always been that herpesviruses have diversified from a common ancestor, in a manner mediating cospeciation of herpesviruses with their host species through latent infection. Indeed, analyses of our phylogenetic results strongly support the notion of host-linked evolution of the

$\gamma$ 2-herpesviruses, at least for the RV1 genogroup, because chimpanzees and gorillas, the nonhuman primate species closest to humans, are infected by the  $\gamma$ 2-herpesvirus homologs closest to KSHV, the human  $\gamma$ 2-herpesvirus prototype.

## METHODS

### Animals

Blood specimens from 40 great apes including 28 chimpanzees and 12 gorillas were studied. The larger series comprises 27 wild-born animals (23 chimpanzees and 4 gorillas), originating from different parts of Cameroon, where they were originally kept as pets after their mothers had been killed by hunters. They were then gathered in a wildlife rescue center in the South West province of Cameroon, in which some of them were kept in the same enclosure, often in close contact (Corbet et al. 2000). The second group (three chimpanzees and two gorillas) originated from the large animal center of the Centre International de Recherches Médicales, Franceville (CIRMF) in Gabon (Georges-Courbot et al. 1996). The other animals came from two different zoos in France, five gorillas and two chimpanzees from La Palmyre Zoo (kindly provided by T. Petit) and one gorilla from Saint Martin la Plaine Zoo (kindly provided by P. Thivillon). For all the animals from France, except one (Gorph682), only serum was available. All these great apes were seronegative for simian immunodeficiency viruses/human immunodeficiency viruses (SIV/HIV) except two animals from Cameroon (Cam 3 and Cam 4, named in our study PanCamDja and PanCamJac, respectively) from which a SIVcpz has recently been isolated and characterized (Corbet et al. 2000). Great apes from the CIRMF exhibited an HTLV Western blot seroindeterminate profile (Georges-Courbot et al. 1996), but all the 35 other animals were negative for HTLV-1/STLV-1 infection. Data in Table 1 concern only animals for which we had both serum and DNA.

### KSHV Serological Analysis

All the plasma were tested, at a 1/40 dilution, for the KSHV-specific IgG, using an immunofluorescence assay (KSHV IFA, ABI). This assay, using the KS-1 cell line as the KSHV source of antigens, detects antibodies directed against both latent and lytic KSHV antigens, and is well adapted to conduct epidemiological works (Chatlynne et al. 1998; Plancoulaine et al. 2000). This assay does not detect any other human herpesviruses than KSHV.

### DNA Extraction and Herpesvirus DNA Polymerase Gene Amplification

DNA was extracted from buffy coats with the QIAamp DNA Blood mini kit (QIAGEN) following the manufacturer's instructions.

Herpesvirus DNA polymerase gene sequences were amplified by consensus heminested PCR based on a previously described method (Rose et al. 1997). We slightly modified the reported cycling conditions as: 10 min at 94°C, 5 cycles of 30 sec at 94°C, 1 min at 60°C, 1 min at 72°C; followed by 30 cycles of 30 sec at 94°C, 30 sec at 46°C, 30 sec at 72°C. An extension of 10 min at 72°C was realized on the last cycle (Perkin Elmer GeneAmp PCR system 9600 thermal cycler). The initial round of PCR contained 500 ng of genomic DNA, 30 pmoles of degenerate primers, 2 mM MgCl<sub>2</sub>, 0.2 mM each dNTP, 5  $\mu$ L of 10 $\times$  PCR buffer, and 0.5  $\mu$ L of Taq Gold DNA polymerase in a volume of 50  $\mu$ L; 1  $\mu$ L of this reaction was used in the heminested reaction. After two rounds of

heminested PCR (GDTD1B and DFASA primers in the initial round followed by GDTD1B and VYGA), 172-bp fragments (excluding primer sequences) were obtained. Specific reverse primer (Pan2as) was designed from these sequences (Table 2; Fig. 1) and used in heminested PCRs with the primer DFASA to finally obtain a 476-bp fragment of viral DNA polymerase sequence (excluding primers).

Finally, we designed an additional primer (P2s) derived from a conserved amino acid motif within the DNA *pol* gene of herpesviruses to allow amplification of a longer DNA polymerase fragment (Table 2; Fig. 1). PCR cycling conditions were 10 min at 94°C, 30 cycles of 30 sec at 94°C, 30 sec at 52°C, 1 min at 72°C; followed by a final extension of 10 min at 72°C. After two rounds of heminested PCR (P2s and P2eas in the initial round then P2s and P2ias) a 870-bp fragment of viral sequence was obtained. Primers P2eas and P2ias were derived from the DFASA-GDTD1B sequences previously determined. For all experiments, stringent precautions against PCR contamination were taken. The amplification mixes were made in a special room physically separated from the laboratory, and at least two negative controls (mix, water, or cellular DNA prepared from a KSHV-negative sample) were included.

### Specific PanRHV2 DNA Polymerase Gene Amplification

To determine the PanRHV2 prevalence, PanRHV2-specific PCR primers were designed from alignments of PanRHV2 sequences obtained by the degenerate DFASA-GDTD1B PCR procedure. PanRHV2-specific PCR outer primer pair Pp2es and Pp2as generates a 312-bp product, whereas the inner primer set Pp2is and Pp2as (Table 2; Fig. 1) amplifies a 274-bp product. Template volume and reagent concentrations were identical to the DNA *pol* consensus assay, and PCR conditions were 10 min at 94°C, 30 cycles of 30 sec at 94°C, 30 sec at 53°C, 30 sec at 72°C; and a final extension of 10 min at 72°C. Nested reactions used 2% of primary reaction product as template with the reaction component concentrations and PCR cycling conditions identical to the primary reaction.

### Mitochondrial DNA (mtDNA) Amplification

Single-round PCR amplification and sequence analysis, without interim cloning, of chimpanzee mitochondrial (mt) DNA was performed on a 498-bp segment of the mitochondrial D-loop control region (corresponding to position 15998–16497 of the human mitochondrial sequence; Anderson et al. 1981) from PBMC DNA as previously described (Gao et al. 1999; Corbet et al. 2000). The sequences of the primers used for chimpanzee mtDNA amplification are given in Table 2.

### Southern Blot Analysis

Nested PCR products (VYGA-GDTD1B or Pp2is-Pp2as) were size-fractionated by 1.5% agarose gel electrophoresis. Following electrophoresis, gels were incubated for 30 min in 0.5 M NaOH–1.5 M NaCl and then for 30 min in 3 M sodium acetate (pH 5.0), after which they were transferred overnight by capillarity onto Biotodyne A nylon membranes (Pall Corporation). DNA was cross-linked to the membranes by exposure to UV light in a UV Stratallinker (Stratagene Cloning Systems) and incubated for at least 6 h in hybridization buffer containing 6× Saline Sodium Phosphate EDTA (SSPE), 0.1% SDS, 5× Denhardt's solution, 50% deionized formamide, and 100 µg/mL of fragmented salmon sperm DNA at 42°C (prehybridization). Hybridizations were performed in the same buffer after the addition of the [<sup>32</sup>P]dATP end-labeled internal corresponding probes. The hybridized membranes were washed first for 1 h in 2× SSPE and 0.1% SDS and then for 15 min in 0.2× SSPE and 0.1% SDS at temperatures ranging from 45°C to 65°C, depending on the probe used. Washed membranes were exposed to phosphor screens and analyzed in a Phos-

phorimager (Molecular Dynamics, Amersham-France SA). The sequences of the probes are given in Table 2.

### Cloning, DNA Sequencing, and Phylogenetic Analyses

The TA cloning procedure, DNA sequencing, as well as the phylogenetic tree constructions using the *PHYLIP* package have been described previously (Lacoste et al. 2000a, 2000c).

Regarding the names of the new primate herpesvirus described in this paper, we have tentatively and provisionally named it PanRHV2 for Pan Rhadino-herpesvirus 2. However, among the specialists in the field, there is discussion and some debate about a new proposal for primate *Rhadinivirus* nomenclature. When new names are approved by their consensus, we will naturally modify the names of these new herpesviruses.

### ACKNOWLEDGMENTS

V.L. is a fellow of the CANAM. This work was partly supported by grants from the Agence Nationale de Recherches sur le SIDA (ANRS), SIDACTION, Association de Recherches sur le Cancer (ARC), and Action Concertée from the network of the Institut Pasteur. We acknowledge Peter Jenkins and Liza Gadsby (the Pandrillus Directors) for their great help in obtaining some of the blood samples studied and their continuous interest in this work. We also thank Thierry Petit from la Palmyre Zoo and Pierre Thivillon from the Saint Martin la Plaine Zoo for providing some of the studied samples.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

### REFERENCES

- Albrecht, J.C. 2000. Primary structure of the *Herpesvirus atele* genome. *J. Virol.* **74**: 1033–1037.
- Albrecht, J.C. and Fleckenstein, B. 1990. Structural organization of the conserved gene block of *Herpesvirus saimiri* coding for DNA polymerase, glycoprotein B, and major DNA binding protein. *Virology* **174**: 533–542.
- Anderson, S., Bankier, A.T., Barrell, B.G., de Bruijn, M.H., Coulson, A.R., Drouin, J., Eperon, I.C., Nierlich, D.P., Roe, B.A., Sanger, F., et al. 1981. Sequence and organization of the human mitochondrial genome. *Nature* **290**: 457–465.
- Auerbach, M.R., Czajak, S.C., Johnson, W.E., Desrosiers, R.C., and Alexander, L. 2000. Species specificity of macaque *Rhadinivirus* glycoprotein B sequences. *J. Virol.* **74**: 584–590.
- Bosch, M.L., Strand, K.B., and Rose, T.M. 1998. Gammaherpesvirus sequence comparisons. *J. Virol.* **72**: 8458–8459.
- Chang, Y., Cesarman, E., Pessin, M.S., Lee, F., Culpepper, J., Knowles, D.M., and Moore, P.S. 1994. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science* **266**: 1865–1869.
- Chatlynne, L.G., Lapps, W., Handy, M., Huang, Y.Q., Masood, R., Hamilton, A.S., Said, J.W., Koeffler, H.P., Kaplan, M.H., Friedman-Kien, A., et al. 1998. Detection and titration of human herpesvirus-8-specific antibodies in sera from blood donors, acquired immunodeficiency syndrome patients, and Kaposi's sarcoma patients using a whole virus enzyme-linked immunosorbent assay. *Blood* **92**: 53–58.
- Corbet, S., Muller-Trutwin, M.C., Versmisse, P., Delarue, S., Ayouba, A., Lewis, J., Brunak, S., Martin, P., Brun-Vezinet, F., Simon, F., et al. 2000. *env* sequences of simian immunodeficiency viruses from chimpanzees in Cameroon are strongly related to those of human immunodeficiency virus group N from the same geographic area. *J. Virol.* **74**: 529–534.
- Desrosiers, R.C., Sasseville, V.G., Czajak, S.C., Zhang, X., Mansfield, K.G., Kaur, A., Johnson, R.P., Lackner, A.A., and Jung, J.U. 1997. A herpesvirus of rhesus monkeys related to the human Kaposi's sarcoma-associated herpesvirus. *J. Virol.* **71**: 9764–9769.
- Gagneux, P., Wills, C., Gerloff, U., Tautz, D., Morin, P.A., Boesch, C., Fruth, B., Hohmann, G., Ryder, O.A., and Woodruff, D.S. 1999. Mitochondrial sequences show diverse evolutionary histories of

- African hominoids. *Proc. Natl. Acad. Sci. USA* **96**: 5077–5082.
- Gao, F., Bailes, E., Robertson, D.L., Chen, Y., Rodenburg, C.M., Michael, S.F., Cummins, L.B., Arthur, L.O., Peeters, M., Shaw, G.M., et al. 1999. Origin of HIV-1 in the chimpanzee *Pan troglodytes troglodytes*. *Nature* **397**: 436–441.
- Georges-Courbot, M.C., Moisson, P., Leroy, E., Pingard, A.M., Nerrienet, E., Dubreuil, G., Wickings, E.J., Debels, F., Bedjabaga, I., Poaty-Mavoungou, V., et al. 1996. Occurrence and frequency of transmission of naturally occurring simian retroviral infections (SIV, STLV, and SRV) at the CIRMF Primate Center, Gabon. *J. Med. Primatol.* **25**: 313–326.
- Gessain, A., Mauclere, P., van Beveren, M., Plancoulaine, S., Ayoub, A., Essame-Oyono, J.L., Martin, P.M., and de The, G. 1999. Human herpesvirus 8 primary infection occurs during childhood in Cameroon, Central Africa. *Int. J. Cancer* **81**: 189–192.
- Gonder, M.K., Oates, J.F., Disotell, T.R., Forstner, M.R., Morales, J.C., and Melnick, D.J. 1997. A new West African chimpanzee subspecies? *Nature* **388**: 337.
- Greensill, J. and Schulz, T.F. 2000. Rhadinoviruses (gamma2-herpesviruses) of Old World primates: Models for KSHV/HHV8-associated disease? *AIDS* **14**: S11–19.
- Greensill, J., Sheldon, J.A., Murthy, K.K., Bessonette, J.S., Beer, B.E., and Schulz, T.F. 2000a. A chimpanzee *Rhadinovirus* sequence related to Kaposi's sarcoma-associated herpesvirus/human herpesvirus 8: Increased detection after HIV-1 infection in the absence of disease. *AIDS* **14**: F129–135.
- Greensill, J., Sheldon, J.A., Renwick, N.M., Beer, B.E., Norley, S., Goudsmit, J., and Schulz, T.F. 2000b. Two distinct gamma-2 herpesviruses in African green monkeys: A second gamma-2 herpesvirus lineage among Old World primates? *J. Virol.* **74**: 1572–1577.
- Lacoste, V., Judde, J.G., Briere, J., Tulliez, M., Garin, B., Kassa-Kelembho, E., Morvan, J., Couppie, P., Clyti, E., Forteza Vila, J., et al. 2000a. Molecular epidemiology of human herpesvirus 8 in Africa: Both B and A5 K1 genotypes, as well as the M and P genotypes of K14.1/K15 loci, are frequent and widespread. *Virology* **278**: 60–74.
- Lacoste, V., Mauclere, P., Dubreuil, G., Lewis, J., Georges-Courbot, M.C., and Gessain, A. 2000b. KSHV-like herpesviruses in chimps and gorillas. *Nature* **407**: 151–152.
- Lacoste, V., Mauclere, P., Dubreuil, G., Lewis, J., Georges-Courbot, M.C., Rigoulet, J., Petit, T., and Gessain, A. 2000c. Simian homologues of human gamma-2 and betaherpesviruses in mandrill and drill monkeys. *J. Virol.* **74**: 1193–1199.
- Magrath, I. and Judde, J. 1996. Epstein-Barr virus and neoplasia. In *Molecular biology of cancer* (ed. M. Bertino). Academic Press, New York.
- Morin, P.A., Moore, J.J., Chakraborty, R., Jin, L., Goodall, J., and Woodruff, D.S. 1994. Kin selection, social structure, gene flow, and the evolution of chimpanzees. *Science* **265**: 1193–1201.
- Plancoulaine, S., Abel, L., van Beveren, M., Tregouet, D.A., Joubert, M., Tortevoe, P., de The, G., and Gessain, A. 2000. Human herpesvirus 8 transmission from mother to child and between siblings in an endemic population. *Lancet* **356**: 1062–1065.
- Roizmann, B., Desrosiers, R.C., Fleckenstein, B., Lopez, C., Minson, A.C., and Studdert, M.J. 1992. The family Herpesviridae: An update. The Herpesvirus Study Group of the International Committee on Taxonomy of Viruses. *Arch. Virol.* **123**: 425–449.
- Rose, T.M., Strand, K.B., Schultz, E.R., Schaefer, G., Rankin, G.W., Jr., Thouless, M.E., Tsai, C.C., and Bosch, M.L. 1997. Identification of two homologs of the Kaposi's sarcoma-associated herpesvirus (human herpesvirus 8) in retroperitoneal fibromatosis of different macaque species. *J. Virol.* **71**: 4138–4144.
- Schulz, T.F. 1998. Kaposi's sarcoma-associated herpesvirus (human herpesvirus-8). *J. Gen. Virol.* **79**: 1573–1591.
- Schultz, E.R., Rankin, G.W., Jr., Blanc, M.P., Raden, B.W., Tsai, C.C., and Rose, T.M. 2000. Characterization of two divergent lineages of macaque rhadinoviruses related to Kaposi's sarcoma-associated herpesvirus. *J. Virol.* **74**: 4919–4928.
- Searles, R.P., Bergquam, E.P., Axthelm, M.K., and Wong, S.W. 1999. Sequence and genomic analysis of a rhesus macaque *Rhadinovirus* with similarity to Kaposi's sarcoma-associated herpesvirus/human herpesvirus 8. *J. Virol.* **73**: 3040–3053.
- Sinkovics, J.G. and Horvath, J.C. 1999. Kaposi's sarcoma: Breeding ground of herpesviridae: A tour de force over viral evolution. *Int. J. Oncol.* **14**: 615–646.
- Strand, K., Harper, E., Thormahlen, S., Thouless, M.E., Tsai, C., Rose, T., and Bosch, M.L. 2000. Two distinct lineages of macaque gamma herpesviruses related to the Kaposi's sarcoma associated herpesvirus. *J. Clin. Virol.* **16**: 253–269.

Received February 14, 2001; accepted in revised form June 4, 2001.