



A Microarray-Based Antibiotic Screen Identifies a Regulatory Role for Supercoiling in the Osmotic Stress Response of *Escherichia coli*

Kevin J. Cheung, Vasudeo Badarinarayana, Douglas W. Selinger, et al.

Genome Res. 2003 13: 206-215

Access the most recent version at doi:[10.1101/gr.401003](https://doi.org/10.1101/gr.401003)

References This article cites 62 articles, 21 of which can be accessed free at:
<http://genome.cshlp.org/content/13/2/206.full.html#ref-list-1>

License

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Cold Spring Harbor Laboratory Press

A Microarray-Based Antibiotic Screen Identifies a Regulatory Role for Supercoiling in the Osmotic Stress Response of *Escherichia coli*

Kevin J. Cheung, Vasudeo Badarinarayana, Douglas W. Selinger, Daniel Janse, and George M. Church¹

Department of Genetics, Harvard Medical School, Boston, Massachusetts 02115, USA

Changes in DNA supercoiling are induced by a wide range of environmental stresses in *Escherichia coli*, but the physiological significance of these responses remains unclear. We now demonstrate that an increase in negative supercoiling is necessary for transcriptional activation of a large subset of osmotic stress-response genes. Using a microarray-based approach, we have characterized supercoiling-dependent gene transcription by expression profiling under conditions of high salt, in conjunction with the microbial antibiotics novobiocin, pefloxacin, and chloramphenicol. Algorithmic clustering and statistical measures for gauging cellular function show that this subset is enriched for genes critical in osmoprotectant transport/synthesis and *rpoS*-driven stationary phase adaptation. Transcription factor binding site analysis also supports regulation by the global stress σ factor *rpoS*. In addition, these studies implicate 60 uncharacterized genes in the osmotic stress regulon, and offer evidence for a broader role for supercoiling in the control of stress-induced transcription.

Changes in the level of DNA supercoiling coincide with a diverse spectrum of environmental events including nutritional upshift, entry into stationary phase, temperature stress, peroxide stress, and osmotic shock (Hengge-Aronis 1999; Lopez-Garcia and Forterre 2000; Weinstein-Fischer et al. 2000; Travers et al. 2001). These phenomena have been best characterized in the stress responses of *Escherichia coli* but have been described in a number of other bacterial species as well (Rohde et al. 1994; Jordi et al. 1995; Alice and Sanchez-Rivas 1997; Ali et al. 2002). Concurrently, expression and activity of global transcriptional regulators such as Fis (*fis*), cyclic AMP receptor protein (*crp*), and stress-induced σ factor σ^S (*rpoS*) have been shown to be dependent on the supercoiling state of the cell (Finkel and Johnson 1992; Schneider et al. 1999). Supercoiling modulates the transcriptional effects of these regulators directly by affecting the efficiency of protein binding to their DNA targets, or indirectly by altering the transcriptional expression of the regulators themselves. These observations have raised the possibility that DNA supercoiling may play a functional role in coupling stress signals to transcriptional activity (Dorman 1996).

In this paper, we report the application of whole-genome microarrays toward understanding the significance of supercoiling in the osmotic stress response. Osmotic shock is among the most common environmental challenges faced by bacterial organisms (Kempf and Bremer 1998; Wood 1999), and serves as an ideal model system because of the rapidity of its effects. High osmolarity causes a rapid increase in negative supercoiling, with subsequent relaxation to preinduction values as the cell recovers. These supercoiling effects are accom-

panied by a distinct sequence of cellular events: plasmolysis, followed by active potassium influx and glutamate synthesis to restore intracellular water, and finally, replacement of potassium glutamate with osmoprotectants more compatible with cell growth. Artificially induced positive supercoiling delays recovery and retards growth. Meanwhile, increased negative supercoiling is reported to stimulate transcription of several osmoregulated genes in vitro and in vivo, including the primary active transporter for the osmoprotectants proline and glycine- β -betaine (*proU*) (Higgins et al. 1988; Jordi and Higgins 2000) and the lipoprotein *osmE* (Conter et al. 1997), as well as to repress transcription of the outer-membrane β -barrel porin *ompF* (Graeme-Cook et al. 1989).

Here we demonstrate that increased negative supercoiling is necessary for proper elicitation of the *E. coli* osmotic shock response. Using a whole-genome approach (Lockhart et al. 1996; DeRisi et al. 1997; Richmond et al. 1999; Selinger et al. 2000), we have characterized the genetic subprograms activated in response to osmotic shock, clustered by expression profiles under salt stress and perturbed with three antibiotics: novobiocin, pefloxacin, and chloramphenicol. Statistical assignment of cellular function using the GenProtEC *E. coli* database (Riley and Serres 2000) identifies enrichment for several major functional categories and one subset of genes with transcriptional kinetics consistent with supercoiling-dependent regulation. This subset is composed of genes implicated in the adaptation of *E. coli* to osmotic stress, including genes in the major osmoprotectant synthesis/transport families, betaine (*betABIT*), trehalose (*otsAB*), and proline (*proVWX*). Statistical analyses of known transcription factor binding sites support the involvement of *rpoS*, and computational predictions implicate other global regulators including *crp* and *fis* that are sensitive to supercoiling. We propose a functional role for supercoiling in the osmotic stress response, and suggest that supercoiling may have broader significance in stress-induced transcription.

¹Corresponding author.

E-MAIL church@arep.med.harvard.edu; **FAX** (617) 432-7266.

Article and publication are at <http://www.genome.org/cgi/doi/10.1101/gr.401003>. Article published online before print in January 2003.

RESULTS

Microarray Studies of Genome-Wide *E. coli* Transcription in Response to Salt and a Panel of Microbial Antibiotics

We examined whole-genome mRNA expression in *E. coli* MG1655 (Blattner et al. 1997) following exposure to five different conditions: novobiocin, high salt, novobiocin plus salt, pefloxacin, and chloramphenicol (Table 1). Samples obtained at 0 and 10 min after treatments with novobiocin or salt, at 0, 7, and 10 min with pefloxacin, and at 0, 10, and 40 min with chloramphenicol were fluorescently labeled and hybridized to microarrays.

The microbial antibiotics novobiocin and pefloxacin were chosen for their effects on supercoiling in the cell (Maxwell 1997). *E. coli* topoisomerase II, also known as gyrase, is unusual among topoisomerases in that it can introduce negative supercoils into DNA by ATP hydrolysis (Luttinger 1995; Wang 1996; Champoux 2001). Novobiocin binding decreases the affinity of *gyrB* (the β subunit of topoisomerase II) for the ATP nucleotide, which is required for DNA breakage and strand passage (Gellert et al. 1976). Novobiocin therefore causes increased positive supercoiling. The quinolone antibiotic pefloxacin also targets gyrase, but possesses a different mechanism of inhibition. Pefloxacin, in contrast to novobiocin, stabilizes the transition state of gyrase after DNA breakage, leading to the formation of a cleavable ternary complex (Drlica and Zhao 1997). This complex can form a barrier, which on collision with replication forks, leads to double-stranded breaks. Joint treatment with novobiocin and salt produces an intermediate response (Conter et al. 1997). Chloramphenicol blocks the 23S subunit of the ribosome and thus protein synthesis. This antibiotic is not reported to have direct effects on supercoiling, but we include it to facilitate in the discovery of coregulated genes.

Total RNA samples were enriched for mRNA, biotinylated, hybridized to Affymetrix *E. coli* arrays, and scanned according to the Affymetrix protocol. Genes below detection threshold as reported by Affymetrix software were eliminated from consideration. For further stringency, a z-test was used to eliminate probes near threshold as defined by negative controls found on the microarrays. Using these criteria, 2146 of 4249 mRNAs and untranslated RNAs were deemed above detection with *P* values ≤ 0.01 .

Table 1. List of Experimental Conditions

Condition	Time point (min)	Growth media
Baseline	0	Minimal
Novobiocin	10	Minimal
Salt	10	Minimal
Novobiocin + salt	10	Minimal
Baseline	0	Rich
Pefloxacin	1	Rich
Pefloxacin	7	Rich
Baseline	0	Rich
Chloramphenicol	10	Rich
Chloramphenicol	40	Rich

All conditions were sampled in duplicate with the exception of pefloxacin at 1 min and 7 min, and chloramphenicol at 10 min. Pearson correlation coefficients between replicates over the set of all mRNAs and noncoding RNAs ranged from 0.90 to 0.99.

Table 2. Top Five Up- and Down-Regulated Genes in the Osmotic Stress Response

Gene	Product	log ₂ fold change
<i>proX</i>	High-affinity transport system for glycine betaine and proline	5.9
<i>proW</i>	High-affinity transport system for glycine betaine and proline	5.7
<i>proV</i>	ATP-binding component of transport system for glycine, betaine, and proline	4.5
<i>otsA</i>	Trehalose-6-phosphate synthase	4.5
<i>b1481</i>	Orf, hypothetical protein	4.0
<i>ykgM</i>	Putative ribosomal protein	-2.9
<i>upp</i>	Uracil phosphoribosyltransferase	-3.2
<i>treC</i>	Trehalase 6-P hydrolase	-3.5
<i>yifE</i>	Orf, hypothetical protein	-4.0
<i>lamB</i>	Phage lambda receptor protein; maltose high-affinity receptor	-4.5

The log₂ fold change was calculated as log₂ (gene expression with salt treatment)/(gene expression with no treatment).

gyrB and *topA* Transcription Match Predicted Effects of Supercoiling

E. coli topoisomerases are regulated by supercoiling in a negative feedback loop (Menzel and Gellert 1987). Positive supercoiling stimulates gyrase expression, whereas negative supercoiling stimulates topoisomerase I expression. Results shown in Figure 1A and B support this model. *gyrB* mRNA expression increases with novobiocin and pefloxacin treatments, but decreases with salt. In contrast, expression of the *topA* gene (encoding topoisomerase I) increases with salt, and is unchanged with novobiocin. These results are consistent with the known effects of these treatment conditions on DNA supercoiling (Fig. 1C).

Functional Characterization of the Osmotic Stress Regulon

Genes significantly induced or repressed by salt were identified by changes in expression greater than twofold, and in the same direction consistently between replicates. One hundred seventy-five genes satisfied these criteria. We list the top 5 up-regulated and top 5 down-regulated genes in Table 2. In order to functionally characterize the osmotic stress response, we used a two-stage approach. Genes were first clustered using self-organizing maps. Six-cluster partitioning was chosen for optimal balance between separation and number of clusters (Fig. 2). Second, the resulting clusters were assigned cellular functions using the GenProtEC: *E. coli* genome and protein database (Riley and Labeledan 1997; Riley and Serres 2000) (Table 3). Following the method of Tavazoie et al. (1999), we estimated the probability that an observed set of genes with a common functional role could have segregated into an individual cluster at random (Sokal and Rohlf 1995). *P* values below the Bonferroni-corrected significance threshold (*P* value < 0.01) are suggestive of functional enrichment (see Methods).

As shown in Table 3, several major functional classes of genes govern the osmotic stress response. These include adaptation to stress and osmoregulation (cluster 0), peptidoglycan biosynthesis (cluster 1), macromolecular biosynthesis (cluster 2), amino acid biosynthesis (cluster 4), and the PTS Mannose-Fructose-Sorbose family (cluster 5). A majority of the known osmoregulated genes as listed by GenProtEC are




Figure 1 *topA* and *gyrB* gene expression mirror predicted effects of supercoiling. *topA* encodes topoisomerase I, and *gyrB* encodes the β subunit of gyrase. (A) *Escherichia coli* MG1655 was cultured to optical density (OD) 0.4 and sampled (no treatment), or induced with novobiocin, salt, or novobiocin + salt. Treated cultures were sampled 10 min postinduction. (B) *E. coli* was cultured in a separate experiment to OD 0.6 and treated with pefloxacin. (C) Predicted changes in DNA supercoiling from salt or gyrase inhibitors based on the literature.

found in cluster 0 (P value $\leq 5.4 \times 10^{-4}$). These results provide evidence that the transcriptional response to osmotic shock greatly extends beyond what is currently known about osmoregulated genes, and provides a promising set of candidates for future experimental validation.

Potassium and Osmoprotectant Genes Segregate into Different Clusters

The immediate effect of hyperosmotic stress is plasmolysis, leading to reduced respiration and growth arrest (Ingraham and Marr 1996). Restoration of intracellular water by potassium influx and glutamate synthesis occurs within minutes, but is only an interim solution because high intracellular ion concentrations are also unfavorable for cell growth (Record Jr. et al. 1998). Osmoprotectants, in contrast, allow re-

sumption of growth, but must be synthesized or brought into the cell by active transport. Thus, two emergency systems, one fast and one slow, are triggered by osmotic stress (Wood 1999). In support of this model, our data demonstrate that Trk family and Kdp family potassium transport genes (clusters 1 and 3) cocluster separately from osmoprotectant genes (cluster 0). Major components for synthesis/transport of osmoprotectants betaine (*betABIT*), trehalose (*otsAB* and *treF*), and proline transport (*proVWX*) are all located in cluster 0. In contrast, *trkH* is located in cluster 1, whereas *kdpABCD*, although outside the twofold threshold, exhibits expression profiles similar to cluster 3 ($r = 0.89$ – 0.99). Likewise, the low affinity proline transporter (*proP*, cluster 3) and components of the murein oligopeptide transporter (*oppA*, cluster 3; *oppDEF*, cluster 1) segregate outside of cluster 0. *proP* is activated in medium hyperosmolarity and its synthesis coincides with potassium influx. Glutamate is a necessary building block in the synthesis of the peptidoglycan wall (Van Heijenoort 1996). Up-regulation of murein oligopeptide transporters (Goodell and Higgins 1987) may therefore reflect coupling of peptidoglycan degradation with potassium counter-ion synthesis.

Salt Stress Causes a Reduction in Metabolic Synthesis

Clusters 2, 4, and 5 also demonstrate the presence of a significant down-regulated component in the osmotic stress response. These include biosynthetic genes for amino acids (cluster 4), flagellar biosynthetic genes (cluster 2), and genes encoding components for galactitol and maltose transport (clusters 2 and 5, respectively). Maltose transport has been previously linked to trehalose synthesis, and is repressed by increased osmolarity in the absence of inducer (Boos and Shuman 1998). Galactitol, however, has not been described in the context of osmotic stress (Nobelmann and Lengeler 1996). Peptidoglycan-associated lipoprotein, *pal*, and peptidoglycan synthetase, *mrdA*, are also repressed by salt (cluster 2), as is trehalase 6-phosphate hydrolase (*treC*), which degrades intracellular trehalose. These findings are consistent with the mirror-image up-regulation seen for murein oligopeptide uptake and trehalose synthesis genes, respectively.

Cluster 0 is Enriched for Supercoiling-Dependent Transcription

If supercoiling regulates the expression of osmotic stress genes, then we would predict that nonphysiological supercoiling would cripple the osmotic stress response. Genes regulated by supercoiling should be down-regulated in novobio-




Figure 2 The osmotic stress response partitions into distinct clusters of drug sensitivity. Mean-variance normalized expression profiles for the 175 genes changed greater than twofold with salt treatment were partitioned into six clusters using self-organizing maps. The black line denotes the mean expression profile of the cluster, and the gray lines indicate one standard deviation above and below this mean (N) Novobiocin; (S) salt; (NS) novobiocin + salt; (P) pefloxacin; (C) chloramphenicol.

cin, and pefloxacin, and up-regulated in salt, or vice versa. Conversely, genes that fit this pattern are candidates for regulation by a supercoiling-dependent mechanism. Topoisomerase genes *gyrB* and *topA* (Fig. 1), for example, have expression profiles satisfying this criterion. Using this working model, we scored the 175 genes for their consistency with this expression pattern (Fig. 3A). Scores were based on 10 criteria or equivalently two replicates of five constraints. Higher scores correspond to greater consistency with our predicted model. For a given score, we calculated the number of genes with at least this score or better for each cluster, and tested the null hypothesis that this distribution could have occurred by chance. These calculations demonstrate that at all score cut-offs, only cluster 0 is statistically enriched for supercoiling-dependent transcription (Fig. 3B and Table 4). Given the functional significance of cluster 0 in osmoprotectant synthesis and stress adaptation (Table 3), we propose that negative supercoiling is a physiologically necessary activation signal for osmotically induced transcription.

The Global Transcription Factor *rpoS* Is Implicated in the Regulation of Cluster 0

Cluster 0 expression (Fig. 2) is significantly lower in the absence of salt stress, in support of a model in which supercoil-

ing does not act alone, but rather is a co-activator of gene expression. Several lines of evidence indicate that *rpoS*, encoding the stationary phase/stress-activated σ factor σ^S , may be a partner in this interaction (Hengge-Aronis 1996). First, *rpoS* is a member of cluster 0, and by our assay, shows supercoiling-dependent regulation (see Table 4). In addition, the *rpoS* transcript has a short half-life that is dramatically stabilized by high osmolarity (Muffler et al. 1996). Second, cluster 0 is enriched for stationary phase genes. *rpoS* was initially characterized as a critical regulator of the stationary phase response; *rpoS* mutants demonstrate both reduced transcription of a number of stationary phase-activated genes as well as decreased viability (Lange and Hengge-Aronis 1991). Cross-referencing stationary and log phase data by Selinger et al. (2000), we have found that, of 50 genes significantly increased in stationary phase relative to log phase, 29 are found in cluster 0, corresponding to a P value of enrichment of 1.2×10^{-4} . This result indicates overlap between stationary phase and osmotic stress responses, very likely through *rpoS*. Third, a number of genes in cluster 0 have been shown to be directly controlled by *rpoS*. Of 13 *rpoS*-regulated genes described by Hengge-Aronis (1996) that are changed greater than twofold in the salt condition, 11 are also located in this cluster (P value = 1×10^{-4}). Interestingly, the lipoprotein *nlpD*, found in cluster 0, has been shown to be part of an operon with *rpoS* (Lange and Hengge-Aronis 1994). More recently, glutaredoxin 2 (*grxB*), also in this cluster, has been shown to be *rpoS*-dependent (Potamitou et al. 2002). We therefore consider it likely that the other genes in this cluster are also regulated by a similar mechanism. Fourth, transcription by σ^S (*rpoS*) is enhanced by both high salt conditions (Ohnuma et al. 2000) and by more positively supercoiled templates in *in vitro* studies (Kusano et al. 1996). This indicates that *rpoS* may be sensitive to supercoiling topology *in vivo*. These findings are consistent with the hypothesis that the effects of supercoiling are mediated through *rpoS*.

We have also calculated enrichment for predicted transcription factor binding sites within upstream noncoding regions of genes in the osmotic stress response (Table 5). Predicted binding sites were generated computationally from a library of DNA-binding site matrices built from known transcription factor binding sites (Robinson et al. 1998; Roth et al. 1998). Eleven DNA-binding motifs were enriched at P values below 0.05, with three motifs corresponding to global regulators known to interact with supercoiling. These include σ^S (*rpoS*), Fis (*fis*), and cyclic AMP receptor protein (*crp*). We were unable to resolve motifs further to individual clusters by this method. Many of the global transcription factors linked to supercoiling recognize structural features in addition to and sometimes preferentially over explicit sequence motifs. Structural recognition presents a challenge to computational bind-

Table 3. The Osmotic Stress Response Partitions into Distinct Classes of Cellular Function

Cluster	<i>p</i> value	GenProt designation	Description	Observed (F _{NC})	Expected (F _{NC})
0	7.06E-05	5.5	Adaptation to stress	12	4.9
0	5.38E-04	5.5.1	Osmotic pressure	10	4.2
1	8.99E-05	1.7.34	Peptidoglycan (murein) turnover, recycling	3	0.1
1	1.93E-04	6	Cell structure	6	1.5
2	4.32E-03	1.6	Macromolecules (cellular constituent) biosynthesis	6	2.0
2	5.03E-03	1.1.1	Carbon compounds	11	5.3
2	9.11E-03	6.4	Flagellum	4	1.1
2	9.11E-03	1.6.12	Flagella	4	1.1
2	9.11E-03	5.3	Motility (incl. chemotaxis, energytaxis, aerotaxis, redoxaxis)	4	1.1
2	9.23E-03	1.1	Carbon utilization	12	6.5
3	9.29E-04	1.7.1	Unassigned reversible reactions	3	0.3
4	8.13E-06	1.5	Building block biosynthesis	13	4.2
4	4.41E-04	1.5.1	Amino acids	9	2.9
5	7.06E-04	4.5	Substrate	11	4.4
5	8.77E-04	4	Transport	12	5.2
5	2.31E-03	4.4.A.6	The PTS Mannose-Fructose-Sorbose (Man) family	3	0.4
5	6.45E-03	1.1	Carbon utilization	9	4.0
5	6.70E-03	1.1.1	Carbon compounds	8	3.3
5	7.34E-03	4.4	Group translocators	4	1.0
5	7.34E-03	4.4.A	Phosphotransferase systems (PEP-dependent PTS)	4	1.0

The six clusters were screened using a hypergeometric test measuring statistical enrichment for functional categories from the GenProtEC *Escherichia coli* database. Observed (F_{NC}) denotes the number of genes in the specified functional category that are found in the cluster. Expected (F_{NC}) denotes the number of genes expected by chance to lie within the cluster. *P* values below 0.01 were deemed significant using Bonferroni correction for multiple hypothesis testing phosphoenolpyruvate (PEP).

ing site prediction by sequence data alone. In these cases, genome-wide in vivo protein–DNA cross-linking experiments (genome-wide ChIP) will be expected to be particularly informative (Lieb et al. 2001; Simon et al. 2001).

Expression Profiles of Temperature Stress Genes Indicate Supercoiling-Dependent Regulation

We observed that the heat shock gene, *clpB*, was among the highest fold up-regulated with novobiocin treatment (2.7-fold), whereas the cold shock gene, *cspA*, was among the highest fold down-regulated (0.57-fold). It has been reported that transient increases in positive supercoiling follow heat shock; conversely, increased negative supercoiling is observed following cold shock (Ogata et al. 1996; Tse-Dinh et al. 1997; Phadtare et al. 1999; Yura and Nakahigashi 1999). To examine this further, we clustered known temperature stress genes into two categories (see Fig. 4). Strikingly, cold shock genes predominate in cluster 0, whereas heat shock genes predominate in cluster 1. Among heat shock genes found in cluster 0, *rpoE*, *rseA*, and *htrA* are all members of the σE regulon (Pallen and Wren 1997) involved in envelope protein stress, and *mseB* is a suppressor of *htrB*, found in cluster 1. Thus, at least four of six heat shock genes found in our putative cold shock cluster (cluster 0) would not be expected a priori to cluster with up-regulated heat shock genes. Similarly, the single cold shock gene found in cluster 1 is *gyrA*. *gyrA* synthesis is stimulated following cold stress. A short lag phase, however, precludes this increased transcription, and, in fact, *gyrA* synthesis actually decreases during this lag phase (Jones et al. 1992). These results offer the possibility that supercoiling-driven regulation is a general property of bacterial stress responses.

DISCUSSION

Here we demonstrate that increased negative supercoiling is necessary for induction of a functionally significant set of

genes activated in the osmotic stress response. Through a series of genome-wide expression profiling experiments under conditions of high salt and perturbation with drug antibiotics, we have identified several major categories of cellular function involved in the osmotic stress response (Fig. 2, Table 3) including adaptation to stress and known osmoregulation (cluster 0), peptidoglycan biosynthesis (cluster 1), macromolecular biosynthesis (cluster 2), amino acid biosynthesis (cluster 4), and the PTS Mannose-Fructose-Sorbose family (cluster 5). Our choice of experimental conditions permitted identification of supercoiling-dependent gene transcription. These supercoiling-dependent transcription profiles are enriched in cluster 0 (Fig. 3B). Significantly, cluster 0 constitutes 35% of the OSR (61/175), and is enriched for genes involved in stress protection such as osmoprotectant synthesis. We propose that increased negative supercoiling is critical for recovery because it mediates the activation of an important subcomponent of the osmotic stress response program.

Computational analysis of predicted DNA regulatory binding domains found upstream of genes in the osmotic stress response also indicates involvement of a number of global regulators known to interact with supercoiling, including *fis*, *crp*, and *rpoS* (Table 5). Comparison of known downstream targets with their distribution among our cluster supports regulation of cluster 0 by *rpoS* transcriptional activation. We consider it likely that supercoiling exerts its effects through these transcription factors. More complex modeling of the genetic network will be expedited by algorithms that incorporate the combinatorial interactions of these transcription factors, as well as binding site data derived from genome-wide ChIP analysis, and in vitro binding specificity assays (Bulyk et al. 2001; Pilpel et al. 2001; Simon et al. 2001).

It is an interesting question as to what causes negative supercoiling to increase following osmotic shock. ATP/ADP ratios are observed on treatment with salt (Hsieh et al. 1991). Because gyrase activity is coupled to the phosphorylation po-




Figure 3 Cluster 0 is enriched for supercoiling-dependent gene regulation. (A) Scoring function for supercoiling-dependent transcription. A total of five tests with two replicates were incorporated into our scoring function. Because genes could be up-regulated or down-regulated in response to supercoiling, we took the better score assuming either of these two cases. Thus, scores could range from 5–10. (B) The 175 genes in the putative osmotic stress response were assayed for their consistency with a supercoiling-dependent profile using the scoring function described in A. The minimum score denotes the lowest score or better that is included in the set of supercoiling-regulated genes. (C) For each minimum score, a hypergeometric test was used to gauge statistical enrichment over all clusters. Only cluster 0 shows statistically significant enrichment.

tential of the cell (van Workum et al. 1996), increased ATP concentration following salt stress is hypothesized to stimulate gyrase activity. In support of this mechanism, an increased ATP/ADP ratio is also observed following nutritional upshift, with a similar increase in negative supercoiling (Balke and Gralla 1987; Travers et al. 2001). Because long-term survival is contingent on adaptation to a new environment, transcriptional activity of the genes found in cluster 0 may be hardwired for optimal expression at high levels of negative supercoiling. We acknowledge that our osmotic shock experiments are performed in a rough *E. coli* K12 strain. Future experiments should include a comparison of our microarray data with that derived from an *E. coli* strain with full-length lipopolysaccharide (LPS) or alternatively, *Salmonella typhimurium*. Interestingly, the *his* operon in *S. typhimurium* has been shown to be induced by novobiocin and repressed by high osmolarity. An analogous role for supercoiling may therefore exist in this bacterium as well.

Our data provide a number of bioinformatics insights. These include a genome-wide characterization of the osmotic

shock response and tentative assignments for 60 uncharacterized genes in this transcriptional program. The 30 genes in cluster 0 with supercoiling-dependent transcription (Table 4) should expand the number of gene targets available for future study of this regulatory mechanism. Gmuender et al. (2001) have previously examined the genome-wide effects of novobiocin and ciprofloxacin (an antibiotic in the same class as pefloxacin) in *Haemophilus influenzae*. The microarray data described here allow comparative study of drug inhibition in both *H. influenzae* and *E. coli* (Table 6), and should facilitate the discovery of potential drug targets. For example, *gmhA* mutants show increased outer membrane permeability from aberrant lipopolysaccharide synthesis, leading to increased sensitivity to hydrophobic compounds such as novobiocin (Brooke and Valvano 1996). As shown in Table 6, however, both novobiocin and quinolone classes of antibiotics repress *gmhA* synthesis. An interesting possibility is that the efficacy of these drugs can be attributed in part to their ability to increase membrane permeability. Finally, we present here a set of computational tools to gauge statistical enrichment for cellular function drawn from GenProtEC as well for transcription factor binding sites. These technologies should help accelerate the functional annotation of the genome. Both programs and datasets are available on our Web site, <http://arep.med.harvard.edu/supercoiling/supplement.htm>.

We propose that transiently induced changes in supercoiling may be relevant in environmental challenges beyond osmotic stress. Clustering of temperature stress genes yields expression profiles consistent with reported changes in supercoiling under heat shock and cold shock (Fig. 4). We have also found that *oxyR* transcription increases with positive supercoiling. Hydrogen peroxide stress induces a transient increase in positive supercoiling; *oxyR* mutants show delayed resupercoiling and peroxide response (Weinstein-Fischer et al. 2000). An intriguing question is whether bacteria will fare worse if challenged with multiple stresses that require conflicting supercoiling responses. Identification of pharmacological effects on supercoiling may therefore aid in the design of drug combinations with synergistic potency.

METHODS

Array Design

Using an array of oligos capable of specifically hybridizing to their target sequence (RNA or DNA), an entire mRNA popu-

Table 4. Cluster 0 is Enriched for Supercoiling-Dependent Gene Regulation

Gene	Product
<i>acnA</i>	Aconitate hydratase 1
<i>b1664</i>	Possible enzyme
<i>b1724</i>	Orf, hypothetical protein
<i>b2809</i>	Orf, hypothetical protein
<i>bax</i>	Putative ATP-binding protein
<i>btuE</i>	Vitamin B ₁₂ transport
<i>dps</i>	Global regulator, starvation conditions
<i>gcd</i>	Glucose dehydrogenase
<i>grxB</i>	Glutaredoxin 2
<i>nlpD</i>	Lipoprotein
<i>osmE</i>	Activator of <i>ntrL</i> gene
<i>otsA</i>	Trehalose-6-phosphate synthase
<i>otsB</i>	Trehalose-6-phosphate phosphatase, biosynthetic
<i>poxB</i>	Pyruvate oxidase
<i>proV</i>	ATP-binding component of transport system for glycine, betaine, and proline
<i>proW</i>	High-affinity transport system for glycine, betaine, and proline
<i>proX</i>	High-affinity transport system for glycine, betaine, and proline
<i>rpoS</i>	RNA polymerase, sigma S (sigma38) factor; synthesis of many growth phase-related proteins
<i>sugE</i>	Suppresses groEL, may be chaperone
<i>tktB</i>	Transketolase 2 isozyme
<i>wrbA</i>	Trp repressor binding protein; affects association of trp repressor and operator
<i>yacK</i>	Orf, hypothetical protein
<i>ybaL</i>	Putative transport protein
<i>ybaY</i>	Glycoprotein/polysaccharide metabolism
<i>ybeL</i>	Putative alpha helical protein
<i>ygaM</i>	Orf, hypothetical protein
<i>yggB</i>	Putative transport protein
<i>ykfE</i>	Orf, hypothetical protein
<i>ynhG</i>	Orf, hypothetical protein
<i>yrbL</i>	Orf, hypothetical protein

Thirty of 61 genes in cluster 0 have scores of 8 or better using the metric described in Fig. 3A.

lation can be probed in parallel (Lockhart et al. 1996). The oligonucleotide array used here is a 544 × 544 grid divided into 24 × 24- μ m regions (Affymetrix). Each region contains ~10⁷ copies of a 25-mer oligonucleotide probe. Photolithography and combinatorial chemical methods are used to synthesize the oligonucleotides directly on a derivatized glass plate. Probe oligonucleotides are grouped into pairs consisting of a perfect match (PM) probe and a mismatch (MM) probe. The PM probe is complementary to the target sequence, whereas the MM probe contains a single base pair mismatch. The MM oligo serves as a control used in identifying cross-hybridization. Probe pairs are, in turn, grouped into probe sets that correspond to mRNA transcripts.

Experimental Design

Ten conditions were assayed in our experiments (see Table 1) using *E. coli* MG1655 (provided by Fred Blattner, University of Wisconsin, Madison, WI). For novobiocin and salt treatments, *E. coli* was grown to optical density (OD) 0.4 in M9 minimal media supplemented with 0.4% glucose at 37°C, and aliquoted into two separate flasks. In the first flask, prewarmed novobiocin/dH₂O was added for a final concentration of 300 μ g/mL of novobiocin. In the second flask, prewarmed NaCl/dH₂O was added for a final osmolarity of 0.8. All samples were spun down, flash-frozen in dry ice-ethanol, and stored at -70°C in accordance to the procedures outlined

by DeRisi et al. (1997). For pefloxacin and chloramphenicol treatments, *E. coli* MG1655 was grown in rich media to OD 0.6 at 37°C. Pefloxacin was added to a final concentration of 1 μ g/mL. The chloramphenicol concentration was 0.32 μ g/uL. Samples were taken directly to phenol. All conditions are averages of two biological replicates (i.e., replicates of separate independent experiments) except for pefloxacin 1-min, pefloxacin 7-min, and chloramphenicol 10-min time points, which are generated from single experiments only.

Sample Preparation/Labeling/Measurement

RNA was isolated by phenol-chloroform extraction and prepared following the Affymetrix protocol for *E. coli* arrays. This protocol includes an mRNA enrichment step using RNase H enzyme to reduce cross-hybridization from ribosomal RNA. Chips were read using an HP-Affymetrix scanner, and quantified using Affymetrix GeneChip 3.2 software. The expression of each probe set was quantified based on the intensities of its PM and MM probes by a composite statistic, the "Average Difference" intensity. The Average Difference (AvgDiff) is calculated as the mean of all PM-MM pairs after removal of outliers. The AvgDiff is a more representative measure of target sequence concentration than PM intensities alone, because it accounts for cross-hybridization. The resultant data were then background subtracted and total intensity was normalized to 5000 using Affymetrix GeneChip version 3.2 software.

Normalization Specifications

We eliminated noisy gene expression data by a significance test developed previously (Selinger et al. 2000). We assumed that (1) intensities observed among 80 negative control probe sets on the chip are representative of cross-hybridization and other forms of noise only, and (2) noise is approximately normally distributed. For each gene, we calculated its mean expression across all conditions (MEC). We then estimated a mean MEC and standard deviation based on 80 negative control probes on the array. For each gene on the chip, we performed the following procedure. If the MEC was 2.33 standard deviations above the mean MEC of our estimated noise distribution, we rejected the null hypothesis that the observed gene expression was noise with a *P* value \leq 0.01. Genes that were below the 2.33 standard deviation cutoff were called "absent" by Affymetrix GeneChip software in greater than

Table 5. Upstream Regions of Genes in the Osmotic Stress Response Are Enriched for Regulatory Binding Sites

DNA-binding protein	<i>p</i> value	# of putative motifs found in OSR	# expected in OSR
<i>crp</i>	9.13E-05	86	62.4
<i>soxS</i>	1.09E-04	94	70.3
<i>fls</i>	2.03E-04	120	97.5
<i>glpR</i>	2.85E-04	125	103.4
<i>ompR</i>	4.00E-04	93	71.4
<i>rpoS</i>	1.11E-03	124	104.8
<i>lrp</i>	8.88E-03	139	125.3
<i>cytR</i>	1.28E-02	34	23.5
<i>tyrR</i>	1.97E-02	85	71.6
<i>malT</i>	4.15E-02	8	4.0
<i>cpxR</i>	5.00E-02	22	15.4

Regulatory binding sites predicted by 55 DNA-binding site matrices found on our website (www.arep.med.harvard.edu) were tallied among genes in the putative osmotic stress response. Statistical enrichment was gauged as described previously (osmotic stress response [OSR]).

Supercoiling Is a Regulator of the Osmotic Stress Response




Figure 4 Heat shock and cold shock genes segregate into supercoiling-responsive profiles. A total of 32 temperature stress genes were selected from the 2146 genes above noise as defined by Blattner notation (Blattner et al. 1997), GenProtEC (Karp et al. 2002), and EcoGene (Rudd 2000). Genes were partitioned into two clusters using the conditions denoted in Fig. 2. (A) Expression profiles for novobiocin (N) and salt (S) conditions are plotted. (B) The distribution of cold shock and heat shock genes are shown for each cluster.

eight conditions, or that contained negative values in any condition were eliminated from further analysis.

Functional Enrichment Analysis

Functional categories from the genProtEC database (Riley et al. 1997; Riley and Serres 2000) were downloaded from <http://>

genprotec.mbl.edu. The actual scoring algorithm is described as follows. Given a file of clustered genes, the perl script catscore.pl (CATEgory Score) counts over all functional categories and clusters the number of genes in a given cluster that are members of a particular functional category. Tabulated data are subsequently analyzed using the statistical test developed by Tavazoie et al (1999):

$$P(k) = 1 - \sum_{a=0}^{k-1} p(a) = 1 - \sum_{a=0}^{k-1} \frac{\binom{f}{a} \binom{n-f}{n-a}}{\binom{n}{c}}$$

where $P(k)$ = the cumulative probability of observing at least k genes in a functional category within a given cluster, f = the number of genes within the functional category, n = the total number of genes, a = the number of genes that match the functional category within the cluster, and c = the number of genes inside the cluster.

We note that this test is mathematically equivalent to a one-tailed Fisher's exact test for 2×2 contingency tables (Sokal and Rohlf 1995). With multiple clusters, we use the Bonferroni correction for reducing experimental error rate. The procedure is simply:

$$\alpha' = 1 - (1 - \alpha)^{1/(m-1)}$$

where α is the desired error rate, α' is the necessary cutoff to achieve this error rate, and m is the number of clusters. For example, if $\alpha = 0.05$, and there are six clusters, $\alpha' = 1 - (1 - 0.05)^{1/5} = 0.01$.

Computational Analysis of Motif Enrichment

Motifs were generated with the AlignACE program (Robinson et al. 1998; Roth et al. 1998), using known footprinting sites from the DPInteract database. The highest scoring motif for each protein was then scanned against the *E. coli* genome using the program ScanACE. High-scoring sequences were defined to be two standard deviations below the mean of the input binding sites or greater, and constituting greater than

Table 6. Common Effects of Novobiocin and Quinolone Antibiotic Treatment for Both *Escherichia coli* and *Haemophilus influenzae*

In <i>Escherichia coli</i> MG1655		In <i>Haemophilus influenzae</i> KW20		%ident	Drug effect
gene	description	gene	description		
<i>b2255</i>	Putative transformylase	<i>hi0623</i>	Methionyl-tRNA formyltransferase	28	↑
<i>fmt</i>	10-Formyltetrahydrofolate:L-methionyl-tRNA(fMet) N-formyltransferase	<i>hi0623</i>	Methionyl-tRNA formyltransferase	65	↑
<i>b2392</i>	Putative transport system permease	<i>hi1728</i>	Conserved hypothetical protein	25	↓
<i>gyrA</i>	DNA gyrase, subunit A, type II topoisomerase	<i>hi1264</i>	DNA gyrase, subunit A	71	↑
<i>gyrB</i>	DNA gyrase subunit B, type II topoisomerase, ATPase activity	<i>hi0567</i>	DNA gyrase, subunit B	75	↑
<i>nusB</i>	Transcription termination; L factor	<i>hi1304</i>	N utilization substance protein B	55	↑
<i>topA</i>	DNA topoisomerase type I, omega protein	<i>hi1365</i>	DNA topoisomerase I	72	↓
<i>ygjU</i>	Putative transport protein	<i>hi1545</i>	Sodium dicarboxylate symporter protein	62	↓
<i>dctA</i>	Uptake of C4-dicarboxylic acids	<i>hi1154</i>	Proton glutamate symporter protein, putative	25	↓
<i>gmhA</i>	Phosphoheptose isomerase	<i>hi1181</i>	Phosphoheptose isomerase	74	↓
<i>yraO</i>	Orf, hypothetical protein	<i>hi1181</i>	Phosphoheptose isomerase	45	↓

H. influenzae genes that were changed greater than two-fold in both novobiocin and ciprofloxacin treatments (Gmuender et al. 2001) were compared against *E. coli* genes with similar changes in both novobiocin and pefloxacin. Sequence homology scores (%ident) were obtained from the Comprehensive Microbial Resource at TIGR (Peterson et al. 2001). *P* values for all gene pairs were better than 10^{-8} .

50% noncoding sequence. The enrichment for high-scoring sequences was then assayed using catscore.pl.

ACKNOWLEDGMENTS

We thank F. Ausubel and S. Lory for helpful comments in a preliminary draft of this work, as well as members of the Church lab for their generous comments and support. This research was supported by grants from the DOE, ONR, and NSF to G.M.C.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

REFERENCES

- Ali, N., Herron, P.R., Evans, M.C., and Dyson, P.J. 2002. Osmotic regulation of the *Streptomyces lividans* thioestrepton-inducible promoter, *ptipA*. *Microbiology* **148**: 381–390.
- Alice, A.F. and Sanchez-Rivas, C. 1997. DNA supercoiling and osmoresistance in *Bacillus subtilis* 168. *Curr. Microbiol.* **35**: 309–315.
- Balke, V.L. and Gralla, J.D. 1987. Changes in the linking number of supercoiled DNA accompany growth transitions in *Escherichia coli*. *J. Bacteriol.* **169**: 4499–4506.
- Blattner, F., Plunkett III, G., Bloch, C.A., Perna, N.T., Burland, V., Riley, M., Collado-Vides, J., Glasner, J.D., Rode, C.K., Mayhew, G.F., et al. 1997. The complete genome sequence of *Escherichia coli* K-12. *Science* **277**: 1453–1474.
- Boos, W. and Shuman, H. 1998. Maltose/maltodextrin system of *Escherichia coli*: Transport, metabolism, and regulation. *Microbiol. Mol. Biol. Rev.* **62**: 204–229.
- Brooke, J.S. and Valvano, M.A. 1996. Molecular cloning of the *Haemophilus influenzae* gmhA (*lpcA*) gene encoding a phosphoheptose isomerase required for lipooligosaccharide biosynthesis. *J. Bacteriol.* **178**: 3339–3341.
- Bulyk, M., Choo, Y., Huang, X., and Church, G.M. 2001. Exploring DNA binding specificities of zinc fingers with DNA microarrays. *Proc. Natl. Acad. Sci.* **98**: 7158–7163.
- Champoux, J.J. 2001. DNA topoisomerases: Structure, function, and mechanism. *Annu. Rev. Biochem.* **70**: 369–413.
- Conter, A., Menchon, C., and Gutierrez, C. 1997. Role of DNA supercoiling and *rpoS* σ factor in the osmotic and growth phase-dependent induction of the gene *osmE* of *Escherichia coli* K12. *J. Mol. Biol.* **273**: 75–83.
- DeRisi, J.L., Iyer, V.R., and Brown, P.O. 1997. Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* **278**: 680–686.
- Dorman, C.J. 1996. Flexible response: DNA supercoiling, transcription and bacterial adaptation to environmental stress. *Trends Microbiol.* **4**: 214–216.
- Drlica, K. and Zhao, X. 1997. DNA gyrase, topoisomerase IV, and the 4-quinolones. *Microbiol. Mol. Biol. Rev.* **61**: 377–392.
- Finkel, S.E. and Johnson, R.C. 1992. The Fis protein: It's not just for DNA inversion anymore. *Mol. Microbiol.* **6**: 3257–3265.
- Gellert, M., O'Dea, M.H., Itoh, T., and Tomizawa, J. 1976. Novobiocin and coumermycin inhibit DNA supercoiling catalyzed by DNA gyrase. *Proc. Natl. Acad. Sci.* **73**: 4474–4478.
- Gmuender, H., Kuratli, K., Di Padova, K., Gray, C.P., Keck, W., and Evers, S. 2001. Gene expression changes triggered by exposure of *Haemophilus influenzae* to novobiocin or ciprofloxacin: Combined transcription and translation analysis. *Genome Res.* **11**: 28–42.
- Goodell, E.W. and Higgins, C.F. 1987. Uptake of cell wall peptides by *Salmonella typhimurium* and *Escherichia coli*. *J. Bacteriol.* **169**: 3861–3865.
- Graeme-Cook, K.A., May, G., Bremers, E., and Higgins, C.F. 1989. Osmotic regulation of porin expression: A role for DNA supercoiling. *Mol. Microbiol.* **3**: 1287–1294.
- Hengge-Aronis, R. 1996. Regulation of gene expression during entry into stationary phase. In *Escherichia coli and Salmonella: cellular and molecular biology* (eds. F.C. Neidhardt et al.), pp. 1497–1512. American Society for Microbiology, Washington D.C.
- . 1999. Interplay of global regulators and cell physiology in the general stress response of *Escherichia coli*. *Curr. Opin. Microbiol.* **2**: 148–152.
- Higgins, C.F., Dorman, C.J., Stirling, L., Waddell, L., Booth, I.R., May, G., and Bremer, E. 1988. A physiological role for DNA supercoiling in the osmotic regulation of gene expression in *S. typhimurium* and *E. coli*. *Cell* **52**: 569–584.
- Hsieh, L., Rouviere-Yaniv, J., and Drlica, K. 1991. Bacterial DNA supercoiling and [ATP]/[ADP] ratio: Changes associated with salt shock. *J. Bacteriol.* **173**: 3914–3917.
- Ingraham, J.L. and Marr, A.G. 1996. Effect of Temperature, Pressure, pH, and Osmotic Stress on Growth. In *Escherichia coli and Salmonella: cellular and molecular biology*. (eds. F.C. Neidhardt et al.), pp. 1570–1577. American Society for Microbiology, Washington D.C.
- Jones, P., Krah, R., Tafuri, S.R., and Wolffe, A.P. 1992. DNA gyrase, CS7.4, and the cold shock response in *Escherichia coli*. *J. Bacteriol.* **174**: 5798–5802.
- Jordi, B. and Higgins, C.F. 2000. The downstream regulatory element of the proU operon of *Salmonella typhimurium* inhibits open complex formation by RNA polymerase at a distance. *J. Biol. Chem.* **275**: 12123–12128.
- Jordi, B.J., Owen-Hughes, T.A., Hulton, C.S., and Higgins, C.F. 1995. DNA twist, flexibility and transcription of the osmoregulated proU promoter of *Salmonella typhimurium*. *EMBO J.* **14**: 5690–5700.
- Karp, P.D., Riley, M., Saier, M., Paulsen, I.T., Collado-Vides, J., Paley, S.M., Pellegrini-Toole, A., Bonavides, C., and Gama-Castro, S. 2002. The EcoCyc database. *Nucleic Acids Res.* **30**: 56–58.
- Kempf, B. and Bremer, E. 1998. Uptake and synthesis of compatible solutes as microbial stress responses to high-osmolarity environments. *Arch. Microbiol.* **170**: 319–330.
- Kusano, S., Ding, Q., Fujita, N., and Ishihama, A. 1996. Promoter selectivity of *Escherichia coli* RNA polymerase E σ 70 and E σ 38 holoenzymes. Effect of DNA supercoiling. *J. Biol. Chem.* **271**: 1998–2004.
- Lange, R. and Hengge-Aronis, R. 1991. Identification of a central regulator of stationary-phase gene expression in *Escherichia coli*. *Mol. Microbiol.* **5**: 49–59.
- . 1994. The *nlpD* gene is located in an operon with *rpoS* on the *Escherichia coli* chromosome and encodes a novel lipoprotein with a potential function in cell wall formation. *Mol. Microbiol.* **13**: 733–743.
- Lieb, J.D., Liu, X., Botstein, D., and Brown, P.O. 2001. Promoter-specific binding of Rap1 revealed by genome-wide maps of protein-DNA association. *Nat. Genet.* **28**: 327–334.
- Lockhart, D.J., Dong, H., Byrne, M.C., Follettie M.T., Gallo, M.V., Chee, M.S., Mittmann, M., Wang, C., Kobayashi, M., Horton, H., et al. 1996. Expression monitoring by hybridization to high-density oligonucleotide arrays. *Nat. Biotechnol.* **14**: 1675–1680.
- Lopez-Garcia, P. and Forterre, P. 2000. DNA topology and the thermal stress response, a tale from mesophiles and hyperthermophiles. *Bioessays* **22**: 736–746.
- Luttinger, A. 1995. The twisted 'life' of DNA in the cell: Bacterial topoisomerases. *Mol. Microbiol.* **15**: 601–606.
- Maxwell, A. 1997. DNA gyrase as a drug target. *Trends Microbiol.* **5**: 102–109.
- Menzel, R. and Gellert, M. 1987. Modulation of transcription by DNA supercoiling: A deletion analysis of the *Escherichia coli* *gyrA* and *gyrB* promoters. *Proc. Natl. Acad. Sci.* **84**: 4185–4189.
- Muffler, A., Fischer, D., and Hengge-Aronis, R. 1996. Posttranscriptional osmotic regulation of the *s* subunit of RNA polymerase in *Escherichia coli*. *J. Bacteriol.* **178**: 1607–1613.
- Nobelmann, B. and Lengeler, J.W. 1996. Molecular analysis of the *gat* genes from *Escherichia coli* and of their roles in galactitol transport and metabolism. *J. Bacteriol.* **178**: 6790–6795.
- Ogata, Y., Mizushima, T., Kataoka, K., Kita, K., Miki, T., and Sekimizu, K. 1996. DnaK heat shock protein of *Escherichia coli* maintains the negative supercoiling of DNA against thermal stress. *J. Biol. Chem.* **271**: 29407–29414.
- Ohnuma, M., Fujita, N., Ishihama, A., Tanaka, K., and Takahashi, H. 2000. A carboxy-terminal 16-amino-acid region of σ (38) of *Escherichia coli* is important for transcription under high-salt conditions and σ activities in vivo. *J. Bacteriol.* **182**: 4628–4631.
- Pallen, M. and Wren, B.W. 1997. The HtrA family of serine proteases. *Mol. Microbiol.* **26**: 209–221.
- Peterson, J.D., Umayam, L.A., Dickinson, T.M., Hickey, E.K., and White, O. 2001. The comprehensive microbial resource. *Nucleic Acids Res.* **29**: 123–125.
- Phadtare, S., Alsina, J., and Inouye, M. 1999. Cold-shock response and cold-shock proteins. *Curr. Opin. Microbiol.* **2**: 175–180.
- Pilpel, Y., Sudarsanam, P., and Church, G.M. 2001. Identifying regulatory networks by combinatorial analysis of promoter elements. *Nat. Genet.* **29**: 153–159.

- Potamitou, A., Neubauer, P., Holmgren, A., and Vlamis-Gardikas, A. 2002. Expression of *Escherichia coli* glutaredoxin 2 is mainly regulated by ppGpp and σ S. *J. Biol. Chem.* **277**: 17775–17780.
- Record Jr., T.M., Courtenay, E.S., Cayley, D.S., and Guttman, H.J. 1998. Responses of *E. coli* to osmotic stress: Large changes in amounts of cytoplasmic solutes and water. *Trends Biochem. Sci.* **23**: 143–148.
- Richmond, C., Glasner, J.D., Mau, R., Jin, H., and Blattner, F.R. 1999. Genome-wide expression profiling in *Escherichia coli* K-12. *Nucleic Acids Res.* **27**: 3821–3835.
- Riley, M. and Labeledan, B. 1997. Protein evolution viewed through *Escherichia coli* protein sequences: Introducing the notion of a structural segment of homology, the module. *J. Mol. Biol.* **268**: 857–868.
- Riley, M. and Serres, M.H. 2000. Interim report on genomics of *Escherichia coli*. *Annu. Rev. Microbiol.* **54**: 341–411.
- Robinson, K., McGuire, A.M., and Church, G.M. 1998. A comprehensive library of DNA-binding site matrices for 55 proteins applied to the complete *Escherichia coli* K-12 genome. *J. Mol. Biol.* **284**: 241–254.
- Rohde, J.R., Fox, J.M., and Minnich, S.A. 1994. Thermoregulation in *Yersinia enterocolitica* is coincident with changes in DNA supercoiling. *Mol. Microbiol.* **12**: 187–199.
- Roth, F.P., Hughes, J.D., Estep, P.W., and Church, G.M. 1998. Finding DNA regulatory motifs within unaligned noncoding sequences clustered by whole-genome mRNA quantification. *Nat. Biotechnol.* **16**: 939–945.
- Rudd, K. 2000. EcoGene: A genome sequence database for *Escherichia coli* K-12. *Nucleic Acids Res.* **28**: 60–64.
- Schneider, R., Travers, A., Kutateladze, T., and Muskhelishvili, G. 1999. A DNA architectural protein couples cellular physiology and DNA topology in *Escherichia coli*. *Mol. Microbiol.* **34**: 953–964.
- Selinger, D.W., Cheung, K.J., Mei, R., Johansson, E.M., Richmond, C.S., Blattner, F.R., Lockhart, D.J., and Church, G.M. 2000. RNA expression analysis using a 30 base pair resolution *Escherichia coli* genome array. *Nat. Biotechnol.* **18**: 1262–1268.
- Simon, I., Barnett, J., Hannett, N., Harbison, C.T., Rinaldi, N.J., Volkert, T.L., Wyrick, J.J., Zeitlinger, J., Gifford, D.K., Jaakkola, T.S., et al. 2001. Serial regulation of transcriptional regulators in the yeast cell cycle. *Cell* **106**: 697–708.
- Sokal, R.R. and Rohlf, F.J. 1995. *Biometry: The principles and practice of statistics in biological research*. W.H. Freeman, New York.
- Tavazoie, S., Hughes, J.D., Campbell M.J., Cho, R.J., and Church, G.M. 1999. Systematic determination of genetic network architecture. *Nat. Genet.* **22**: 281–285.
- Travers, A., Schneider, R., Muskhelishvili, G. 2001. DNA supercoiling and transcription in *Escherichia coli*: The FIS connection. *Biochimie* **83**: 213–217.
- Tse-Dinh, Y., Qi, H., and Menzel, R. 1997. DNA supercoiling and bacterial adaptation: Thermotolerance and thermoresistance. *Trends Microbiol.* **5**: 323–326.
- Van Heijenoort, J. 1996. Murein biosynthesis. In *Escherichia coli and Salmonella: cellular and molecular biology* (eds. F.C. Neidhardt et al.), pp. 1025–1034. American Society for Microbiology, Washington, D.C.
- van Workum, M., van Dooren, S.J., Oldenburg, N., Molenaar, D., Jensen, P.R., Snoep, J.L., and Westerhoff, H.V. 1996. DNA supercoiling depends on the phosphorylation potential in *Escherichia coli*. *Mol. Microbiol.* **20**: 351–360.
- Wang, J.C. 1996. DNA topoisomerases. *Annu. Rev. Biochem.* **65**: 635–692.
- Weinstein-Fischer, D., Elgrably-Weiss, M., and Altuvia, S. 2000. *Escherichia coli* response to hydrogen peroxide: A role for DNA supercoiling, topoisomerase I and Fis. *Mol. Microbiol.* **35**: 1413–1420.
- Wood, J.M. 1999. Osmosensing by bacteria: Signals and membrane-based sensors. *Microbiol. Mol. Biol. Rev.* **63**: 230–262.
- Yura, T. and Nakahigashi, K. 1999. Regulation of the heat-shock response. *Curr. Opin. Microbiol.* **2**: 153–158.

WEB SITE REFERENCES

- <http://arep.med.harvard.edu/supercoiling/supplement.htm>; complete datasets and software as described in manuscript.
- <http://genprotec.mbl.edu>; functional category assignments for the *E. coli* MG1655 genome.
- www.arep.med.harvard.edu; resources and supplementary material for publications from the Church laboratory.

Received May 7, 2002; accepted in revised form October 22, 2002.