



***Drosophila* telomeric retrotransposons derived from an ancestral element that was recruited to replace telomerase**

Alfredo Villasante, José P. Abad, Rosario Planelló, et al.


Genome Res. 2007 17: 1909-1918 originally published online November 7, 2007
Access the most recent version at doi:[10.1101/gr.6365107](https://doi.org/10.1101/gr.6365107)

References This article cites 47 articles, 16 of which can be accessed free at:
<http://genome.cshlp.org/content/17/12/1909.full.html#ref-list-1>

Open Access Freely available online through the *Genome Research* Open Access option.

License Freely available online through the Genome Research Open Access option.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Copyright © 2007, Cold Spring Harbor Laboratory Press

Drosophila telomeric retrotransposons derived from an ancestral element that was recruited to replace telomerase

Alfredo Villasante,^{1,4} José P. Abad,¹ Rosario Planelló,^{1,3} María Méndez-Lago,¹ Susan E. Celniker,² and Beatriz de Pablos¹

¹Centro de Biología Molecular "Severo Ochoa" (CSIC-UAM), Cantoblanco, 28049 Madrid, Spain; ²Berkeley *Drosophila* Genome Project, Department of Genome Sciences, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

Drosophila telomeres do not have arrays of simple telomerase-generated G-rich repeats. Instead, *Drosophila* maintains its telomeres by occasional transposition of specific non-long terminal repeat (non-LTR) retrotransposons to chromosome ends. The genus *Drosophila* provides a superb model system for comparative telomere analysis. Here we present an evolutionary study of *Drosophila* telomeric elements to ascertain the significance of telomeric retrotransposons (TRs) in the maintenance of *Drosophila* telomeres. PCR and in silico surveys in the sibling species of *Drosophila melanogaster* and in more distantly related species show that multiple TRs maintain telomeres in *Drosophila*. In addition to TRs with two open reading frames (ORFs) capable of autonomous transposition, there are deleted telomeric retrotransposons that have lost their ORF2, which we refer to as half telomeric-retrotransposons (HTRs). The phylogenetic relationship among these telomeric elements is congruent with the phylogeny of the species, suggesting that they have been vertically inherited from a common ancestor. Our results suggest that an existing non-LTR retrotransposon was recruited to perform the cellular function of telomere maintenance.

[Supplemental material is available online at www.genome.org. The sequence data from this study have been submitted to GenBank under accession nos. AM161543, AM040250-6, AM040240-1, AM040246-9, and AM040237-9.]

The molecular study of *Drosophila* telomeres began initially by analyzing the plasmid cDm356 (Rubin 1978) and afterward the phage λ T-A (Young et al. 1983). In both cases, the cloned DNA hybridized with all telomeres of *Drosophila melanogaster*. It took a long time to realize that these clones derived from heterochromatin: cDm356 from the centromeric region of chromosome Y (Agudo et al. 1999) and λ T-A from the pericentromeric region of chromosome 3 (Losada et al. 1999). The unexpected findings of telomeric elements at the centromeric region of the Y could be understood if this region derived from an ancestral telomere, as has been proposed by Agudo et al. (1999).

It was not until 10 years after Gerald Rubin had observed hybridization at the telomeres with cDm356 that a telomere-specific retrotransposon, named *HeT-A*, was described as an element capable of healing broken or receding chromosome ends (Traverse and Pardue 1988; Biessmann et al. 1990, 1992a,b). The second element, *TART*, was discovered in the analysis of a native *Drosophila* telomere (Levis et al. 1993). The *TART* element has two open reading frames (ORF1 and ORF2), typical of non-LTR retrotransposons; ORF1 encodes a putative RNA-binding protein and ORF2 encodes a protein with domains related to both reverse transcriptase (RT) and endonuclease (Sheen and Levis 1994). However, *HeT-A* is an atypical element because it lacks ORF2, so the RT for its transposition is produced in *trans* from a yet-unknown source. This type of nonautonomous element has been

previously described in several species. For example, half-elements derived from LINE elements, by deletion of its ORF2, have arisen in both the rat and mouse lineages (Smit 1999; Rat Genome Sequencing Project Consortium 2004). And, it has been shown that LINE-encoded proteins can act in *trans* to promote retrotransposition of half-LINE elements (Wei et al. 2001). In *Drosophila* telomeric arrays, *HeT-A* and *TART* are intermingled and frequently truncated at the 5' end (Abad et al. 2004b). This could partially explain why it has been so difficult to clone and characterize complete copies of these elements. The *HeT-A* and *TART* elements never appear in euchromatin, but they are found in centromeric and pericentromeric heterochromatin (Traverse and Pardue 1988; Danilevskaya et al. 1993; Levis et al. 1993; Losada et al. 1997, 1999; Agudo et al. 1999; Abad et al. 2004a). Recently, the third and less abundant telomere-specific retrotransposon, *TAHRE*, was discovered during the sequence analysis of the telomeres of the *y; cn bw sp* strain of *Drosophila melanogaster* (Abad et al. 2004b,c). The 5'UTR, ORF1, and 3'UTR of *TAHRE* are very similar to the corresponding sequences of *HeT-A*, and its ORF2 encodes a RT that has some similarity to the RT of *TART* (Abad et al. 2004c). The structural and phylogenetic analyses of *HeT-A*, *TAHRE*, and *TART* imply that *HeT-A* derived from ancestral *TAHRE* elements (Abad et al. 2004c). However, Pardue and collaborators favor the hypothesis that *HeT-A* derived from telomerase-encoding sequences (Pardue et al. 1996, 2005; Pardue and DeBaryshe 1999, 2002, 2003).

The identification of *TART* and *HeT-A* elements in *Drosophila yakuba* and *Drosophila virilis* has shown that these elements have been performing the cellular function of telomere maintenance during more than 45 million yr (Danilevskaya et al. 1998b; Casacuberta and Pardue 2002, 2003a,b).

³Present address: Facultad de Ciencias, UNED Senda del Rey 9, 28040 Madrid, Spain.

⁴Corresponding author.

E-mail avillasante@cbm.uam.es; fax 34-91-4974799.

Article published online before print. Article and publication date are at <http://www.genome.org/cgi/doi/10.1101/gr.6365107>. Freely available online through the *Genome Research* Open Access option.

The molecular characterization of *Drosophila* telomeres has been so difficult, in part, because of the intrinsic difficulty in isolating large DNA clones containing chromosome ends (Adams et al. 2000; Celniker et al. 2002; Hoskins et al. 2002). But, recently, the use of sheared DNA libraries has allowed the cloning of these chromosomal regions from *D. melanogaster* (Abad et al. 2004b) as well as from other species (Osoegawa et al. 2006).

In this report we have used a phylogenetic analysis of *Drosophila* telomeric DNA to demonstrate that multiple autonomous and nonautonomous telomere-specific retrotransposons have diverged from the common ancestor that was recruited to perform the cellular function of telomere maintenance. The novel vision of the telomeres in *Drosophila*, presented in this work, could have never been achieved without the sequencing and the comparative analysis of the genomes of the 12 *Drosophila* species.

Results

TAHRE elements in the *D. melanogaster* species subgroup

To investigate the rate and nature of the evolution of *Drosophila* telomeres, it is essential to analyze telomere sequences from species that are at various levels of divergence. Thus, we initiated PCR (see Methods) and in silico surveys in the sibling species of *D. melanogaster* and in more distant species. To identify TAHRE homologs in silico, we performed TBLASTN searches using the

ORF2 of *Dmel*\TAHRE as query for the genomic databases. Subsequent to the in silico survey, we have principally used scaffolds in which the telomeric retrotransposons (TRs) appear in the typical telomeric head-to-tail arrays with no interspersed nontelomeric elements. Using this approach we have identified scaffolds in which a subtelomeric minisatellite (named telomere-associated sequences or TAS) often appears between the most proximal telomeric retrotransposon in the array and the most distal nonrepetitive sequences. In these cases we identified the putative telomere from which they derived, validating our approach. As the success of our analyses depends on the coverage and assembly quality, the number and nature of telomeric contigs we have obtained vary among the sequenced species (Figs. 1B, 2).

In the *Drosophila sechellia* genome we found a scaffold that contains a typical tandem array of two *Dsec*\TART, two *Dsec*\TAHRE, one *Dsec*\HeT-A, and one *Dsec*\TART (Fig. 1B). However, in *D. yakuba* and *Drosophila erecta* there were no single scaffolds containing long TAHRE sequences, so we have constructed representative TAHREs by connecting scaffolds that had overlapping regions that were >97% identical (see Methods). Dot-matrix comparisons of each TAHRE with its corresponding HeT-A clearly show that the HeT-A elements are TAHREs without their ORF2 (Fig. 1A). To determine the chromosomal location and ascertain whether these elements are found at telomeres, we performed fluorescence in situ hybridization using TAHRE-ORF2




Figure 1. TAHRE elements in the *D. melanogaster* species subgroup. (A) Dot-matrix comparisons of the nucleotide sequence of each TAHRE (*Dsec*\TAHRE scaffold 182, 15703–7717; *Dyak*\TAHRE and *Dere*\TAHRE are listed in Supplemental Fig. S12) with its corresponding HeT-A (*Dsec*\HeT-A scaffold 182, 7716–2747; *Dyak*\HeT-A AF043258; *Dere*\HeT-A scaffold 4836, 38316–32346). The three-frame ORF maps of TAHRE elements are indicated (the tall tick marks correspond to stop codons; the short tick marks correspond to methionines; the ORFs are the long stretches without tall tick marks). (B) Diagram of scaffold 182 from *D. sechellia*. The color code for the TRs is indicated in the box below. (C) FISH of TAHRE-ORF2 probes to polytene chromosomes of *D. simulans* and *D. yakuba*. Hybridization signals are shown in green on the chromosomes counterstained with DAPI (blue).




Figure 2. Diagrams of scaffolds containing head-to-tail arrays of telomeric retrotransposons from *D. ananassae*, *D. persimilis*, *D. mojavensis*, *D. virilis*, and *D. grimshawi*. The putative origin of each scaffold appears between brackets. The color code for TRs, HTRs, TAS, and euchromatin is indicated in boxes. In the euchromatin regions a distal gene is indicated. Ns indicates the presence of unsequenced regions.

probes to polytene chromosomes of *Drosophila simulans* and *D. yakuba*. The results showed the presence of TAHRE sequences at two telomeres of *D. simulans* and at three of *D. yakuba* (Fig. 1C). These elements are not found in all telomeres, similar to the situation in *D. melanogaster* (Abad et al. 2004c), but they seem to be present across the *melanogaster* subgroup. As the PCR analysis has obvious limitations (see Methods), and the sequences of the 12 species are at their first assembly stage (Comparative Analysis Freeze 1), it is not surprising that a full-length TAHRE has only been found in *D. melanogaster*, in which identified telomeric clones have been sequenced (Abad et al. 2004c).

Recurrent evolution of HTRs from TRs across the *Drosophila* genus

During our analyses in more distant species, we have found up to 37 phylogenetically distinct telomeric elements, of which 19 were half TRs (HTRs) because they had lost their ORF2. The names of the new elements were given after performing phylogenetic analysis. In order to help the reading and comprehension, a color code based on the phylogenetic relationships was added to the graphical representation of the elements. Thus, elements from different species that share the same color in Figure 2 are closely related to each other since they appear in the same cluster in the trees (see Figs. 5, 6, below). In addition to the 37 previously mentioned elements, we have also detected several subfamilies when the differences between the elements were low. The elements belonging to subfamilies were designated by adding a lower case letter to the name of the family and appear in the figures with the same color but different intensity. Given the high number of new telomeric retrotransposons, only a brief overview of the elements is given for each species.

Drosophila ananassae, a species in the *melanogaster* group, has the second largest genome of the 12 sequenced species, around 231 Mb. After the analysis of its genome, we have found that three scaffolds end in arrays of TRs and HTRs (Fig. 2). In these telomeric arrays, the element *Dana*\TR2A and its corresponding deleted element *Dana*\HTR2A have a characteristic long 3'UTR made of complex repeats (see dot-matrix comparison between them in Fig. 3).

The N-terminal part of the ORF1 of retroelements evolved quickly due to base substitutions and small deletions. However, in *D. ananassae* there are three HTRs (*Dana*\HTRX, *Dana*\HTRY, and *Dana*\HTRZ; Supplemental Fig. S1a) with an additional large deletion that removed the conserved motif called major homology region (MHR) (Supplemental Fig. S1b). Moreover, *Dana*\HTRX and *Dana*\HTRY have long 3'UTRs with several repeats (Supplemental Fig. S1a). Apart from those telomeric scaffolds, we found other scaffolds containing decayed TRs interspersed with other transposable elements. Interestingly, Jennifer Kirkland, John Belote, and Robert Levis had previously isolated a *D. ananassae* clone having those TRs (DQ114943) (Kirkland 2005). They hypothesized that it was probably derived from centromeric heterochromatin after observing that the elements were decayed and that the clone did not give any telomeric signal when hybridized to polytene chromosomes (R. Levis, pers. comm.).




Figure 3. Dot-matrix analysis of TRs with HTRs. Dot-matrix comparisons of the nucleotide sequence of *Dana\TR2A*: *Dana\HTR2A*, *Dper\TR1A*: *Dper\HTR1A*, *Dper\TR3A*: *Dper\HTR3A*, *Dper\TR3B*: *Dper\HTR3B*, *Dmoj\TR1Ab*: *Dmoj\HTR1Ab*, *Dmoj\TR1D*: *Dmoj\HTR1D*, *Dmoj\TR1C*: *Dmoj\HTR1C*, *Dmoj\TR1E*: *Dmoj\HTR1E*. The three-frame ORF maps are indicated.

The genome sequence of *Drosophila persimilis*, a sister species of *Drosophila pseudoobscura*, provides an opportunity to study whether a recent speciation, in the *obscura* group, has affected the structure of telomeric elements. Although, the *Drosophila persimilis* genome has only been sequenced at the $3\times$ level, we have been able to recognize telomeric scaffolds. Thus, we found three scaffolds containing arrays of *TRs* and *HTRs* that correspond to chromosome ends (Fig. 2). We have also found four additional scaffolds that are not linked to telomeres. In these scaffolds we identified five *TRs* (*Dper*\TR1A, *Dper*\TR2A, *Dper*\TR3A, *Dper*\TR3B, and *Dper*\TR4A) and three *HTRs* (*Dper*\HTR1A, *Dper*\HTR3A, and *Dper*\HTR3B). The dot-matrix comparisons between these *TRs* and their corresponding *HTRs* are shown in Figure 3. *Dper*\TR1A, *Dper*\TR2A, and *Dper*\TR3A have short 3'UTRs, *Dper*\TR3B has a long 3'UTR, and *Dper*\TR4A has a large repeated region in its 3'UTR (Fig. 4; Supplemental Figs. S2–S4). The ORF1 of the *Dper*\TR3A element contains dispersed short runs of polyglutamines (poly Qs) after the three zinc knuckle domains, whereas ORF2 has a nonhomogenous long run of poly Qs after the RT domain (Fig. 4).

As expected for two species whose separation has occurred recently, the telomeric elements found in *D. persimilis* are very similar (~98% of nucleotide identity) to those in *D. pseudoobscura* (data not shown).




Figure 4. Diagrams of the structure of *TRs* and *HTRs*. Diagrams are approximately to scale. The 5'UTR regions appear in yellow, the ORF1 in green, the ORF2 in orange, and the 3'UTR in magenta. The three zinc knuckles domain in ORF1 and the RT domain in ORF2 are indicated. The polyglutamines appear as a single "Q" and the polyasparagines as a single "N." The presence of repeats in the 3'UTR regions is also indicated.

Drosophila willistoni belongs to the *willistoni* group and has the largest genome of the 12 sequenced species (~237 Mb). Unfortunately, we could not identify any scaffolds containing *TR* sequences that, by our criteria, derive from a telomere. Nevertheless, we found small scaffolds that encoded partial *TRs* phylogenetically related to *TR2* elements.

Drosophila mojavensis, a cactophilic species in the *repleta* group, is a good model of incipient speciation. Among the new sequenced species, the *D. mojavensis* genome assembly appears to be the most complete (Gilbert 2006). As a consequence, we found five scaffolds linked to telomeres and one additional very long scaffold not linked (Fig. 2). We have identified up to six *TRs* (*Dmoj*\TR1A, *Dmoj*\TR1B, *Dmoj*\TR1C, *Dmoj*\TR1D, *Dmoj*\TR1E, and *Dmoj*\TR3A) and, among the *HTRs* found, four (*Dmoj*\HTR1Ab, *Dmoj*\HTR1B, *Dmoj*\HTR1C, and *Dmoj*\HTR1D) have their corresponding *TR* (see dot-matrix comparison between them in Fig. 3). We have also found three additional *HTRs* whose *TRs* counterparts were not detected (*Dmoj*\HTR2A, *Dmoj*\HTR3B, and *Dmoj*\HTR3C).

The *TRs* and *HTRs* in *D. mojavensis* display a number of different features. Five of the six *TRs* found have a short 3'UTR (*Dmoj*\TR1A, *Dmoj*\TR1B, *Dmoj*\TR1C, *Dmoj*\TR1D, and *Dmoj*\TR3A) and only *Dmoj*\TR1E has a long 3'UTR, with three repeated regions along it (Fig. 4; Supplemental Figs. S4–S7). The ORF1 of *Dmoj*\TR1C has two long runs of poly Qs following the zinc knuckles domain (Fig. 4). *Dmoj*\HTR2Aa has a long 3'UTR with two duplications along it (Fig. 4; Supplemental Fig. S4). *Dmoj*\HTR3Ba has a long 3'UTR with a large repeated region covering the majority of the 3'UTR, and its ORF has several tracts of poly Qs before and after the zinc knuckles domain (Fig. 4; Supplemental Fig. S7). *Dmoj*\HTR3Ca has two large repeated regions covering almost the entire length of its 3'UTR, and its ORF has two tracts of poly Qs after the zinc knuckles domain (Fig. 4; Supplemental Fig. S7). With this data on hand, it would be interesting to analyze whether diverging populations of *D. mojavensis* show clear differences in the structure of the telomeres.

The genome size of *D. virilis* is about twice as large as that of *D. melanogaster*, but this seems to be due to an increase in the amount of simple and complex satellite DNAs. In *D. virilis*, we have found two scaffolds linked to telomeres (Fig. 2). In both cases, between the proximal telomeric element and the euchromatic sequences there are subtelomeric repeats of 370 bp, corresponding to the TAS of *D. virilis*. Interestingly, these are the repeats that Biessmann et al. (2000) found at the telomeres and at internal euchromatic regions. These telomeres have five autonomous elements: *Dvir*\TR1A, *Dvir*\TR1B, *Dvir*\TR2A (previously called *Dvir*\TART by Casacuberta and Pardue 2003a), *Dvir*\TR2B, and *Dvir*\TR2C. They also have three half telomeric-retrotransposons: *Dvir*\HTR0A (previously called *Dvir*\HeT-A by Casacuberta and Pardue 2003b), *Dvir*\HTR0B, and *Dvir*\HTR2B. The *TRs* and *HTRs* of *D. virilis* have short and long 3'UTRs, respectively (Fig. 4; Supplemental Fig. S8). The ORF1s of *Dvir*\TR1B, *Dvir*\TR2C, and *Dvir*\HTR0A have short tracts of poly Qs following the three zinc knuckle domains (Fig. 4; Casacuberta and Pardue 2003b). However, the ORF1 of *Dvir*\TR2A has long runs of poly Q after the zinc knuckles domain (Casacuberta and Pardue 2003a) and the ORF of *Dvir*\HTR0B has long runs of poly Qs and a long tract of polyasparagines (poly Ns) following the zinc knuckles domain (Fig. 4). In *D. virilis*, only the ORF2 of *Dvir*\TR2A has long runs of poly Qs after the RT domain (Fig. 4; Casacuberta and Pardue 2003a).

The Hawaiian *Drosophila* lineage contains >1000 species. *Drosophila grimshawi*, the sequenced picture wing species, is representative of a lineage that began to rapidly diversify 26 million yr ago. In *D. grimshawi*, we identified a single telomeric scaffold (Fig. 2). This scaffold contains arrays made of one *TR* (*Dgri*\TR2A) and two *HTRs* (*Dgri*\HTR2B and *Dgri*\HTR2C). The *Dgri*\TR2A elements have tracts of poly Qs after the three zinc knuckles of the ORF1 and a short 3'UTR (Fig. 4; Supplemental Fig. S9). The *Dgri*\HTR2B has long tracts of poly Qs after the zinc knuckles domain and a large duplication in its 3'UTR (Fig. 4; Supplemental Fig. S9). The *Dgri*\HTR2C does not contain extended glutamine repeats and has a short 3'UTR (Fig. 4; Supplemental Fig. 9).

Since we have analyzed whole-genome shotgun draft sequence that contains many gaps within telomeric regions, we cannot exclude the possibility that corresponding *TRs* may exist in the genomes where only orphan *HTRs* have been detected. Nor can we exclude the possibility that additional elements and even additional families may exist in all of the sequenced species. Future analyses, when further sequencing has been done, are likely to find more telomere-specific elements than the ones described in this article.

Phylogenetic relationships among *Drosophila* telomeric retrotransposons

Normally, phylogenetic analyses of non-LTR retrotransposons are restricted to the RT domain, the only domain common to all elements. Therefore, we have initially constructed the phylogenetic relationships between the *TRs* based on their entire RT domains (for alignments see Supplemental Fig. S10). The limits of the RT domains were defined according to Malik et al. (1999). The phylogenetic tree identifies multiple families of *TRs* that belong, as expected, to the *jockey* clade (Fig. 5).

In *Drosophila*, the C-terminal half of ORF1 of *TRs* and the ORF of *HTRs* contains three conserved domains: the MHR, the zinc knuckles (also called CCHC), and the leucine zipper-like region (Rashkova et al. 2003). To verify the *TRs* phylogeny based on the RT domain, we have also analyzed the phylogenetic relationships between *TRs* and *HTRs* using these ORF1 domains (for alignments see Supplemental Fig. S11). In this analysis, the ORF1s from the *D. ananassae* *HTRs* were omitted, as they lack MHR domains. Importantly, although the ORF1 tree has less resolution than the RT tree, because the ORF1 region is smaller and less conserved than the RT domain, the phylogeny of the *TR* ORF1s is in

agreement with the RT phylogeny (Fig. 6). Moreover, it is in this type of analysis that the evolutionary relationships between *TRs* and their corresponding *HTRs* appear more clearly. Our finding of multiple pairs (Fig. 6) of *TRs* and *HTRs* is strong evidence that *HTRs*, like *HeT-A*, have derived from ancestral *TRs*, rather than from telomerase.

The phylogenetic relationships among the telomere-specific elements are congruent with the phylogeny of the species, which suggests vertical transmission of these elements from a common ancestor recruited to perform the cellular function of telomere maintenance.




Figure 5. Phylogenetic relationships of *TRs* based on their RT domains. The tree was inferred using the neighbor-joining method. Bootstrap values are given as percentage numbers. The color code for *TRs* is indicated. *TART* and *TAHRE* have not been renamed to keep their historical names.




Figure 6. Phylogenetic relationships of TRs and HTRs based on their ORF1 domains. The tree was inferred using the neighbor-joining method. Bootstrap values are given as percentage numbers. The color code for TRs and HTRs is indicated. TART, TAHRE, and HeT-A have not been renamed to keep their historical names.

Discussion

Drosophila telomeric sequences do not behave as regular heterochromatin sequences. While in centromeric heterochromatin there are multiple amplifications, deletions and insertions of transposable elements that produce the decay of the elements present in it, in terminal retrotransposon arrays amplifications and internal deletions still occur, but never interrupt the reading frames of the telomeric elements. In noncoding regions, the amplification events principally occur in the 3'UTRs, and it is possible that multiple amplifications events occur in the same 3'UTR (Fig. 4; Supplemental figures). In coding regions, amplifications in ORF1 and/or ORF2 can take place, where repeats of the triplets

CAG and CAA encode for tracts of polyglutamines (or AAC for polyasparagines). These polyglutamines or polyasparagines are not part of known functional domains and they have been suggested to provide a molecular basis for fast adaptation to environmental changes (Trifonov 2004). As the glutamine-rich and asparagine-rich regions form polar zippers, Perutz et al. (2002) proposed that they mediate interaction between different proteins. It is tempting to speculate that the initial expansion of triplets coding for poly Qs/Ns could have been selected because of the improvement in the ability to interact with other telomeric proteins. However, since extremely long expansions cause the formation of protein aggregates, the elimination of these tracts by unequal crossing-over is expected to happen over time. This could explain why some elements have poly Qs and most of them do not.

In telomere-specific elements it is normal to find deletions of varied sizes in different upstream regions with respect to the zinc knuckles domains of the ORF1. However, the most outstanding event that occurs in telomeric heterochromatin is a recurrent loss of most of the TRs' ORF2, giving rise to HTR elements. This finding is consistent with the suggested origin of HeT-A by deletion of the ORF2 of an ancestral TAHRE (Abad et al. 2004c).

The recent discoveries of the telomerase reverse transcriptase (*TERT*) of *Bombyx mori* (Lepidoptera), *Tribolium castaneum* (Coleoptera), and *Apis mellifera* (Hymenoptera) has confirmed the absence of telomerase orthologs in the available *Drosophila* and *Anopheles* genomes (Osanai et al. 2006; Robertson and Gordon 2006). This is consistent with the absence of TTAGG telomeric repeats in flies and mosquitoes. The phylogenetic relationships using the core RT domain of the three insect *TERT*s showed, as expected, that the insect telomerases cluster together, and that *A. mellifera* appears basal to *T. castaneum* (Robertson and Gordon 2006) in the same way that the Hymenoptera is basal to the Coleoptera in the Holometabola (Honey Bee Genome Sequencing Consortium 2006). These insect *TERT*s lack the N-terminal domain implicated in processivity in others telomerases, and this feature could explain why telomerase activity is low in the bee and virtually undetectable in the silkworm.

The unexpected complexity of the *Drosophila* telomeric sequences seems to challenge the requirement for the conventional telomerase-synthesized repeats with G-tracts. But, the 3' noncoding DNA of *Dmel\HeT-A* elements have the same strand

bias: the strand running 5'–3' toward the end of the chromosome is G+T rich (Danilevskaya et al. 1998a). Moreover, the 3' UTR of *Dmel\HeT-A* elements contains sequences with the propensity to form G-quadruplex DNA structures (Abad and Villasante 1999). The RNA templates of *TRs* and *HTRs* found in the genus *Drosophila* have a C+A bias not restricted to the noncoding regions. Interestingly, the RNA template of other members of the *jockey* clade such as, *Dmel\Doc*, *Dmel\F*, *Dmel\Juan*, and *Dmel\X* also has a C+A bias. Therefore, in *Drosophila*, the universal telomere strand bias seems to be maintained by repeated reverse transcription of telomere-specific C+A biased RNA templates primed by a 3'OH on the end of the chromosomal DNA.

In *D. melanogaster*, as in any eukaryote, recombination-based mechanisms maintain chromosome termini normally (Kahn et al. 2000). Therefore, it is plausible that a progressive telomerase inactivation (as seems to happen now in *B. mori*) would have led to its loss if another telomerase-independent mechanism would have been able to replace telomere function progressively. Interestingly, to compensate for the extremely low activity of the *T. castaneum* and *B. mori* telomerase's, these species seem to have co-opted non-LTR retrotransposons, which belong to the *R1* clade, for the maintenance of telomeres. (Fujiwara et al. 2005; Osanai et al. 2006). A scenario for the origin of *Drosophila* telomeres has been recently proposed (Abad et al. 2004c): once the telomerase was lost, and the recombination-based mechanism to maintain telomeres was acting normally, telomere erosion could activate the mobilization of C+A biased non-LTR retroelements via a DNA-damage signaling pathway (Rudin and Thompson 2001) and their addition to the chromosome ends would restore telomere function (Yamamoto et al. 2003). In agreement with this model, Morrish et al. (2007) have detected endonuclease-independent LINE-1 retrotransposition at dysfunctional telomeres. Finally, the extensive and rapid evolution of *Drosophila* telomere-specific elements could give a wide range of situations for evolutionary innovations.

Methods

Drosophila strains and DNA preparation

Four *Drosophila* species were used for experimental studies: *D. simulans* (14021–0251.176 from the Tucson Stock Center [TSC]), *D. mauritiana* (14021–0241.05, TSC), *D. sechellia* (14021–0248.08, TSC), and *D. yakuba* (14021–0261.00, TSC). Genomic DNA was obtained from adult flies as previously described (Pirrota et al. 1983).

PCR amplification and DNA cloning

The sequences of *Dmel\TAHRE* (AJ542581), *Dmel\TART* (AJ566116), and *Dmel\HeT-A* (U06920) were used to design primers for the PCR amplification of specific regions of telomeric retrotransposons. The sequence of these primers appears in Supplemental Table S1. Because of the great similarity of *HeT-A* and *TAHRE*, with *HeT-A* being much more abundant, we have not managed to PCR-amplify whole copies of *TAHRE* from genomic DNA. In all of the attempts we made, *HeT-A* was the only amplified product. To overcome these difficulties, we designed two sets of primers (Supplemental Table S1) to amplify two overlapping regions of *TAHRE*: one containing the ORF1–ORF2 junction and the other encompassing the ORF2–3'UTR junction, thus covering the entire ORF2 of *TAHRE*. Using this approach, overlapping fragments from *D. simulans*, *D. mauritiana*, and *D. sechellia* were isolated. The high degree of identity of the nucleotide sequence

showed that the reconstructed elements encoded *TAHRE* homologs. However, while *Dsim\TAHRE* (AM040252) and *Dsec\TAHRE* (AM040246) sequences show nondecayed ORFs, suggesting a telomere origin, all cloned *Dmau\TAHRE* (AM040237, AM040238, and AM040239) and some of the *Dsec\TAHRE* (AM040247, AM040248, and AM040249) and *Dsim\TAHRE* (AM040253) sequences have the typical deletions of the decayed telomeric elements found in centromeric and pericentromeric heterochromatin (Losada et al. 1997, 1999; Agudo et al. 1999; Abad et al. 2004a). When we tried the same approach in more distant species, we were unable to get PCR amplification using any set of primers. The amplified DNA was cloned in vector pGEM-T (Promega). DNA fragments containing the *TAHRE* ORF1–ORF2 junction were PCR amplified using genomic DNA from *D. mauritiana*, *D. sechellia*, and *D. simulans* and the pair of primers TH1 from the ORF1 and Tsc1.4 from ORF2. The amplified DNAs were cloned, sequenced, and named pThmau3.1 (AM040237), pThsch3.6 (AM040246), and pTsim3.6 (AM040252), respectively. In *D. simulans* a clone containing *HeT-A* sequences was also obtained (pHeTsim1.9; AM040254). The PCR amplifications of fragments containing the *TAHRE* ORF2–3'UTR junction were performed using the pair of primers TH5 from the ORF2 and TH9 from the 3'UTR. The cloned DNAs were sequenced and named pThmau2.1 (AM040238) and pThmau1.7 (AM040239) in *D. mauritiana*; pThsch2.1.0 (AM040247), pThsch2.1.1 (AM040248), and pThsch1.8 (AM040249) in *D. sechellia*, and pTsim2.1 (AM040253) in *D. simulans*. In *D. yakuba*, we have reconstructed the *TAHRE* homolog using two sequences encompassing the ORF1–ORF2 junction (AAEU02006918 and AAEU02009565), one with a portion of the ORF2 (AAEU02013304), and another one that contains a big region of the ORF2 and the whole 3'UTR (AAEU02000405). To confirm that the reconstructed ORF1–ORF2 junction truly exists in the genome, we PCR amplified a DNA fragment encompassing the junction using the pair of primers Thy1 and Thy2 (see Supplemental Table S1), and the PCR product was cloned (pThyak1.4) and sequenced (AM161543). Similarly, in *D. erecta*, we have found that scaffolds 3102 and 1433 contain an ORF1–ORF2 junction and an ORF2–3'UTR junction of a *TAHRE* homolog, respectively. The reconstructed sequences of *Dyak\TAHRE* and *Dere\TAHRE* are listed in Supplemental Figure S12. The unavailable *TART* ORF2 sequences from *D. mauritiana*, *D. sechellia*, and *D. simulans* were amplified with the primers T14 (AJ566116; 11235–11254) and T28 (AJ566116; 6997–7016), cloned, and sequenced: pTmau4.3 (AM040241), pTsch4.3 (AM040251), and pTsim4.3 (AM040256). The unavailable *HeT-A* sequences from different species were PCR amplified using the following pairs of primers: TH1 and Hs6 (AM040254, 1226–1246) for *D. mauritiana*, TH1 and Hs4 (AM040254, 1199–1216) for *D. sechellia*, and TH9 and Hs4 for *D. simulans*. The cloned DNA fragments were sequenced and named pHeTmau1.2 (AM040240), pHeTsch1.2 (AM040250), and pHeTsim2.7 (AM040255), respectively.

To perform standard PCR amplification, Taq DNA polymerase and reaction buffer from Promega were used in a Perkin-Elmer Thermal Cycler. The 50- μ L reaction mix contained 2 mM MgCl₂, 5 U of Taq polymerase, 100 μ M of dNTPs, 500 ng of genomic DNA, and 30 nmol of each primer. The amplification program was 30 cycles of 1 min, 20 sec at 94°C, 1 min at 50°C, and 2 min, 30 sec at 72°C, increasing the extension step of the last cycle to 7 min. To perform long PCR, the Expand Long Template PCR System kit from Roche Diagnostics was used. The 50- μ L reaction mix contained 1.75 mM MgCl₂, 3.75 U of Taq DNA polymerase, 350 μ M of dNTPs, 500 ng of genomic DNA, and 300 nmol of each primer. The amplification program was 30 cycles of

1 min, 20 sec at 94°C, 4 min at 52°C, and 4 min 30 sec at 68°C, increasing the extension step of the last cycle to 8 min.

DNA sequencing, sequence analyses, and annotation

The sequencing was done using the primers listed in Supplemental Table 1. All sequencing was performed using big dye-termination reagents and ABI 3700 automated sequencers. Sequence analyses were performed online against the DroSpeGe comparative genome database, the FlyBase database, and the database at the National Center for Biotechnology Information. The TBLASTN program (Altschul et al. 1990) was used for these searches. The major problem during the annotation of *TRs* and *HTRs* inside the positive scaffolds was to identify the boundaries of the elements. This task was performed manually using principally dot-matrix comparisons. The name of the *TR* or *HTR*, its length, and the coordinates within its corresponding scaffold appear in Supplemental Table S2. Normally, the translated sequences revealed intact ORFs, but in some cases, ORFs had to be manually reconstructed due to frameshifts in the original sequence. Also, in some cases the ORFs had to be extended upstream of the first encountered methionine. Multiple-sequence alignments were performed with ClustalX (Thompson et al. 1997), followed by manual adjustments of gaps. Phylogenetic trees were generated with the MEGA 2.1 program (Kumar et al. 2001) using a neighbor-joining method with bootstrap statistics (1000) after removal of the gaps. Dot-matrix analyses were performed with the COMPARE and DOTPLOT programs from the CGC package (University of Wisconsin, Madison).

Fluorescence in situ hybridization

The *Dsim\TAHRE* probe used for FISH was pThsim2.1 (AM040253). The *Dyak\TAHRE* probe used for FISH was an ORF2 fragment obtained by PCR using pThyak1.4 as the template and the pair of primers: Thy2 and Ths11: 5'-ACAATATAATCCG CAGCC-3'. Probes were labeled by nick translation using Biotin-16-dUTP (Roche Diagnostics). FISH experiments on polytene chromosomes were carried out under high-stringency conditions, as described in Pimpinelli et al. (2000). Post-hybridization washing was in $0.1 \times$ SSC at 60°C. For probe detection, a 3.3- μ g/mL FITC-conjugated avidin (DCS grade, Vector Laboratories) diluted in SBT (1% BSA, 0.1% Tween-20, $4 \times$ SSC) solution was used. Chromosomes were counterstained with 4',6-diamino-2-phenylindole (DAPI). Digital images were obtained using a Zeiss Axiovert 200 microscope equipped with a cooled CCD camera. The fluorescent signals from the FITC and the DAPI were recorded separately as grayscale digital images and then pseudocolored and merged using the Adobe Photoshop software.

Acknowledgments

We thank R. Levis, R. de Frutos, M.J. Martínez-Sebastián, J. Dopazo, H. Dopazo, and E. Torroja for critical comments and suggestions. We also thank the DroSpeGe comparative genome database and the FlyBase database for access to the 12 *Drosophila* draft genomes assemblies before publication. The work was supported by a grant from Ministerio de Educación y Ciencia (BFU2005-07690-C02-01) to A.V., an institutional grant from Fundación Ramón Areces to the CBMSO, and by a subcontract to Lawrence Berkeley National Laboratory (PI, S.C.) from UC Berkeley (HG000750, PI, G. Rubin) carried out under Department of Energy Contract DE-AC0376SF00098, University of California, Berkeley.

References

- Abad, J.P. and Villasante, A. 1999. The 3' non-coding region of the *Drosophila melanogaster* HeT-A telomeric retrotransposon contains sequences with propensity to form G-quadruplex DNA. *FEBS Lett.* **453**: 59–62.
- Abad, J.P., de Pablos, B., Agudo, M., Molina, I., Giovinnazo, G., Martín-Gallardo, A., and Villasante, A. 2004a. Genomic and cytological analysis of the Y chromosome of *Drosophila melanogaster*: Telomere-derived sequences at internal regions. *Chromosoma* **113**: 295–304.
- Abad, J.P., De Pablos, B., Osoegawa, K., De Jong, P.J., Martín-Gallardo, A., and Villasante, A. 2004b. Genomic analysis of *Drosophila melanogaster* telomeres: Full-length copies of HeT-A and TART elements at telomeres. *Mol. Biol. Evol.* **21**: 1613–1619.
- Abad, J.P., De Pablos, B., Osoegawa, K., De Jong, P.J., Martín-Gallardo, A., and Villasante, A. 2004c. TAHRE, a novel telomeric retrotransposon from *Drosophila melanogaster*, reveals the origin of *Drosophila* telomeres. *Mol. Biol. Evol.* **21**: 1620–1624.
- Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D., Amanatides, P.G., Scherer, S.E., Li, P.W., Hoskins, R.A., Galle, R.F., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* **287**: 2185–2195.
- Agudo, M., Losada, A., Abad, J.P., Pimpinelli, S., Ripoll, P., and Villasante, A. 1999. Centromeres from telomeres? The centromeric region of the Y chromosome of *Drosophila melanogaster* contains a tandem array of telomeric HeT-A- and TART-related sequences. *Nucleic Acids Res.* **27**: 3318–3324.
- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. 1990. Basic local alignment search tool. *J. Mol. Biol.* **215**: 403–410.
- Biessmann, H., Mason, J.M., Ferry, K., d'Hulst, M., Valgeirsdottir, K., Traverse, K.L., and Pardue, M.-L. 1990. Addition of telomere-associated HeT DNA sequences “heals” broken chromosome ends in *Drosophila*. *Cell* **61**: 663–673.
- Biessmann, H., Champion, L.E., O'Hair, M., Ikenaga, K., Kasravi, B., and Mason, J.M. 1992a. Frequent transpositions of *Drosophila melanogaster* HeT-A transposable elements to receding chromosome ends. *EMBO J.* **11**: 4459–4469.
- Biessmann, H., Valgeirsdottir, K., Lofsky, A., Chin, C., Ginther, B., Levis, R.W., and Pardue, M.-L. 1992b. HeT-A, a transposable element specifically involved in “healing” broken chromosome ends in *Drosophila melanogaster*. *Mol. Cell. Biol.* **12**: 3910–3918.
- Biessmann, H., Zurovcova, M., Yao, J.G., Lozovskaya, E., and Walter, M.F. 2000. A telomeric satellite in *Drosophila virilis* and its sibling species. *Chromosoma* **109**: 372–380.
- Casacuberta, E. and Pardue, M.-L. 2002. Coevolution of the telomeric retrotransposons across *Drosophila* species. *Genetics* **161**: 1113–1124.
- Casacuberta, E. and Pardue, M.-L. 2003a. Transposon telomeres are widely distributed in the *Drosophila* genus: TART elements in the *virilis* group. *Proc. Natl. Acad. Sci.* **100**: 3363–3368.
- Casacuberta, E. and Pardue, M.-L. 2003b. HeT-A elements in *D. virilis*: Retrotransposon telomeres are conserved across the *Drosophila* genus. *Proc. Natl. Acad. Sci.* **100**: 14091–14096.
- Celniker, S.E., Wheeler, D.A., Kronmiller, B., Carlson, J.W., Halpern, A., Patel, S., Adams, M., Champe, M., Dugan, S.P., Frise, E., et al. 2002. Finishing a whole-genome shotgun: Release 3 of the *Drosophila melanogaster* euchromatic genome sequence. *Genome Biol.* **3**: doi: 10.1186/gb-2002-3-12-research0079.
- Danilevskaya, O., Lofsky, A., Kurenova, E.V., and Pardue, M.-L. 1993. The Y chromosome of *Drosophila melanogaster* contains a distinctive subclass of HeT-A-related repeats. *Genetics* **134**: 531–543.
- Danilevskaya, O.N., Lowenhaupt, K., and Pardue, M.-L. 1998a. Conserved subfamilies of the *Drosophila* HeT-A telomere-specific retrotransposon. *Genetics* **148**: 233–242.
- Danilevskaya, O.N., Tan, C., Wong, J., Alibhai, M., and Pardue, M.-L. 1998b. Unusual features of the *Drosophila melanogaster* telomere transposable element HeT-A are conserved in *Drosophila yakuba* telomere elements. *Proc. Natl. Acad. Sci.* **95**: 3770–3775.
- Fujiwara, H., Osanai, M., Matsumoto, T., and Kojima, K.K. 2005. Telomere specific non-LTR retrotransposons and telomere maintenance in the silkworm, *Bombyx mori*. *Chromosome Res.* **13**: 455–467.
- Gilbert, D.G. 2006. DroSpeGe: Rapid access database for new *Drosophila* species genomes. *Nucleic Acids Res.* **35**: D480–D485. doi: 10.1093/nar/gk1997.
- Honey Bee Genome Sequencing Consortium. 2006. Insights into social insects from the genome of the honeybee *Apis mellifera*. *Nature* **443**: 931–949.
- Hoskins, R.A., Smith, C.D., Carlson, J., Carvalho, B.A., Halpern, A., Kaminker, J.S., Kennedy, C., Mungall, C.J., Sullivan, B.A., Sutton,

- G.G., et al. 2002. Heterochromatic sequences in a *Drosophila* whole genome shotgun assembly. *Genome Biol.* **3**: doi: 10.1186/gb-2002-3-12-research0085.
- Kahn, T., Savitsky, M., and Georgiev, P. 2000. Attachment of HeT-A sequences to chromosomal termini in *Drosophila melanogaster* may occur by different mechanisms. *Mol. Cell. Biol.* **20**: 7634–7642.
- Kirkland, J. 2005. "Molecular cloning and initial characterization of a TART-like sequence in *Drosophila ananassae*." M.S. thesis, University of Syracuse, NY.
- Kumar, S., Tamura, K., Jakobsen, I.B., and Nei, M. 2001. MEGA2: Molecular evolutionary genetics analysis software. *Bioinformatics* **17**: 1244–1245.
- Levis, R.W., Ganesan, R., Houtchens, K., Tolar, L.A., and Sheen, F.-M. 1993. Transposons in place of telomeric repeats at a *Drosophila* telomere. *Cell* **75**: 1083–1093.
- Losada, A., Abad, J.P., and Villasante, A. 1997. Organization of DNA sequences near the centromere of the *Drosophila melanogaster* Y chromosome. *Chromosoma* **106**: 503–512.
- Losada, A., Agudo, M., Abad, J.P., and Villasante, A. 1999. HeT-A telomere-specific retrotransposons in the centric heterochromatin of *Drosophila melanogaster* chromosome 3. *Mol. Gen. Genet.* **262**: 618–622.
- Malik, H.S., Burke, W.D., and Eickbush, T.H. 1999. The age and evolution of non-LTR retrotransposable elements. *Mol. Biol. Evol.* **16**: 793–805.
- Morrish, T.A., Garcia-Perez, J.L., Stamato, T.D., Taccioli, G.E., Sekiguchi, J., and Moran, J.V. 2007. Endonuclease-independent LINE-1 retrotransposition at mammalian telomeres. *Nature* **446**: 208–212.
- Osanai, M., Kojima, K.K., Futahashi, R., Yaguchi, S., and Fujiwara, H. 2006. Identification and characterization of the telomerase reverse transcriptase of *Bombyx mori* (silkworm) and *Tribolium castaneum* (flour beetle). *Gene* **376**: 281–289.
- Osoegawa, K., Vessere, G.M., Li Shu, C., Hoskins, R.A., Abad, J.P., de Pablos, B., Villasante, A., and de Jong, P.J. 2006. BAC clones generated from sheared DNA. *Genomics* **89**: 291–299.
- Pardue, M.-L. and DeBaryshe, P.G. 1999. Telomeres telomerase: More than the end of the line. *Chromosoma* **108**: 73–82.
- Pardue, M.-L. and DeBaryshe, P.G. 2002. Telomeres transposable elements. In *Mobile DNA II* (ed. N.L. Craig), pp. 870–887. ASM Press, Washington, D.C.
- Pardue, M.-L. and DeBaryshe, P.G. 2003. Retrotransposons provide an evolutionary robust non-telomerase mechanism to maintain telomeres. *Annu. Rev. Genet.* **37**: 485–511.
- Pardue, M.-L., Danilevskaya, O.N., Lowenhaupt, K., Slot, F., and Traverse, K.L. 1996. *Drosophila* telomeres: New views on chromosome evolution. *Trends Genet.* **12**: 48–52.
- Pardue, M.-L., Rashkova, S., Casacuberta, E., DeBaryshe, P.G., George, J.A., and Traverse, K.L. 2005. Two retrotransposons maintain telomeres in *Drosophila*. *Chromosome Res.* **13**: 443–453.
- Perutz, M.F., Pope, B.J., Owen, D., Wanker, E.E., and Scherzinger, E. 2002. Aggregation of proteins with expanded glutamine and alanine repeats of the glutamine-rich and asparagine-rich domains of Sup35 and of the amyloid beta-peptide of amyloid plaques. *Proc. Natl. Acad. Sci.* **99**: 5596–5600.
- Pimpinelli, S., Bonaccorsi, S., Fanti, L., and Gatti, M. 2000. Preparation and analysis of *Drosophila* mitotic chromosomes. In *Drosophila: A Laboratory Manual* (eds. W. Sullivan et al.), pp. 1–24. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Pirrotta, V., Hadfield, C., and Pretorius, G.H.J. 1983. Microdissection and cloning of the white locus and the 3B1–3C2 region of the *Drosophila* X chromosome. *EMBO J.* **2**: 927–934.
- Rashkova, S., Athanasiadis, A., and Pardue, M.-L. 2003. Intracellular targeting of Gag proteins of the *Drosophila* telomeric retrotransposons. *J. Virol.* **77**: 6376–6384.
- Rat Genome Sequencing Project Consortium. 2004. Genome sequence of the Brown Norway rat yields insights into mammalian evolution. *Nature* **428**: 493–521.
- Robertson, H.M. and Gordon, K.H. 2006. Canonical TTAGG-repeat telomeres and telomerase in the honey bee, *Apis mellifera*. *Genome Res.* **16**: 1345–1351.
- Rubin, G.M. 1978. Isolation of a telomeric DNA sequence from *Drosophila melanogaster*. *Cold Spring Harb. Symp. Quant. Biol.* **42**: 1041–1046.
- Rudin, C.M. and Thompson, C.B. 2001. Transcriptional activation of short interspersed elements by DNA-damaging agents. *Genes Chromosomes Cancer* **30**: 64–71.
- Sheen, F.-M. and Levis, R.W. 1994. Transposition of the LINE-like retrotransposon TART to *Drosophila* chromosome termini. *Proc. Natl. Acad. Sci.* **91**: 12510–12514.
- Smit, A.F. 1999. Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Curr. Opin. Genet. Dev.* **9**: 657–663.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., and Higgins, D.G. 1997. The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**: 4876–4882.
- Traverse, K.L. and Pardue, M.-L. 1988. A spontaneously opened ring chromosome of *Drosophila melanogaster* has acquired HeT DNA sequences at both new telomeres. *Proc. Natl. Acad. Sci.* **85**: 8116–8120.
- Trifonov, E.N. 2004. The tuning function of tandemly repeating sequences: A molecular device for fast adaptation. In *Evolutionary theory and processes: Modern horizons* (ed. S.P. Wasser), pp. 115–138. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Wei, W., Gilbert, N., Ooi, S.L., Lawler, J.F., Ostertag, E.M., Kazazian, H.H., Boeke, J.D., and Moran, J.V. 2001. Human L1 retrotransposition: Cis preference versus trans complementation. *Mol. Cell. Biol.* **21**: 1429–1439.
- Yamamoto, Y., Fujimoto, Y., Arai, R., Fujie, M., Usami, S., and Yamada, T. 2003. Retrotransposon-mediated restoration of *Chlorella* telomeres: Accumulation of Zepp retrotransposons at termini of newly formed minichromosomes. *Nucleic Acids Res.* **31**: 4646–4653.
- Young, B.S., Pession, A., Traverse, K.L., French, C., and Pardue, M.L. 1983. Telomere regions in *Drosophila* share complex DNA sequences with pericentric heterochromatin. *Cell* **34**: 85–94.

Received February 6, 2007; accepted in revised form May 16, 2007.