



Identification of Three Novel Ca²⁺ Channel γ Subunit Genes Reveals Molecular Diversification by Tandem and Chromosome Duplication

Daniel L. Burgess, Caleb F. Davis, Lisa A. Gefrides, et al.

Genome Res. 1999 9: 1204-1213

Access the most recent version at doi:[10.1101/gr.9.12.1204](https://doi.org/10.1101/gr.9.12.1204)

References

This article cites 48 articles, 11 of which can be accessed free at: <http://genome.cshlp.org/content/9/12/1204.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

An advertisement banner with a teal background. On the left, the text reads "CRISPR and RNAi Genetic Screening. Your new superpower." In the center, there is a white box with the words "LEARN MORE". On the right, there is a photograph of a woman wearing a red mask and a red cape, and the Cellecta logo, which consists of a green molecular structure and the word "CELLECTA" in white capital letters.

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Cold Spring Harbor Laboratory Press

Identification of Three Novel Ca²⁺ Channel γ Subunit Genes Reveals Molecular Diversification by Tandem and Chromosome Duplication

Daniel L. Burgess,^{1,2} Caleb F. Davis,¹ Lisa A. Gefrides,¹ and Jeffrey L. Noebels¹

¹Department of Neurology, Baylor College of Medicine, Houston, Texas 77030 USA

Gene duplication is believed to be an important evolutionary mechanism for generating functional diversity within genomes. The accumulated products of ancient duplication events can be readily observed among the genes encoding voltage-dependent Ca²⁺ ion channels. Ten paralogous genes have been identified that encode isoforms of the α_1 subunit, four that encode β subunits, and three that encode $\alpha_2\delta$ subunits. Until recently, only a single gene encoding a muscle-specific isoform of the Ca²⁺ channel γ subunit (*CACNG1*) was known. Expression of a distantly related gene in the brain was subsequently demonstrated upon isolation of the *Cacng2* gene, which is mutated in the mouse neurological mutant stargazer (*stg*). In this study, we sought to identify additional genes that encoded γ subunits. Because gene duplication often generates paralogs that remain in close syntenic proximity (tandem duplication) or are copied onto related daughter chromosomes (chromosome or whole-genome duplication), we hypothesized that the known positions of *CACNG1* and *CACNG2* could be used to predict the likely locations of additional γ subunit genes. Low-stringency genomic sequence analysis of targeted regions led to the identification of three novel Ca²⁺ channel γ subunit genes, *CACNG3*, *CACNG4*, and *CACNG5*, on chromosomes 16 and 17. These results demonstrate the value of genome evolution models for the identification of distantly related members of gene families.

[The sequence data described in this paper have been submitted to the GenBank data library under accession numbers AFI42618–AFI42625 and AFI48220.]

Voltage-dependent Ca²⁺ channels couple membrane depolarization in a wide variety of cellular processes, including action potential generation, neurotransmitter and hormone release, muscle contraction, neurite outgrowth, synaptogenesis, Ca²⁺-dependent gene expression, synaptic plasticity, and cell death. This broad range of biological activity is regulated by distinct channel subtypes whose biophysical properties are determined predominantly by subunit isoform composition. Ca²⁺ channels are believed to be heteromultimers of α_1 , β , $\alpha_2\delta$, and γ subunits that associate in a 1:1:1:1 stoichiometry (De Waard et al. 1996). To date, 10 genes have been identified and localized that encode isoforms of the pore-forming α_1 subunit (α_{1A} – α_{1I} , α_{1S} ; Chin et al. 1991; Powers et al. 1991; Drouet et al. 1993; Gregg et al. 1993a; Iles et al. 1993; Diriong et al. 1995; Fisher et al. 1997; Cribbs et al. 1998; Perez-Reyes et al. 1998; Lee et al. 1999), 4 that encode β subunits (β_{1-4} ; Gregg et al. 1993b; Collin et al. 1994; Taviaux et al. 1997), 3 that encode $\alpha_2\delta$ subunits ($\alpha_2\delta_{1-3}$; Powers et al. 1994; Klugbauer et al. 1999), and 2 that encode γ subunits (γ_1 , γ_2 ; Powers et al. 1993; Letts et al. 1998). All but the two skeletal muscle isoforms, α_{1S} and γ_1 , are expressed in the central nervous system (CNS).

Until recently, only a single gene encoding a muscle-specific Ca²⁺ channel γ subunit was known

(*CACNG1*). Subsequent isolation of the molecular defect in the mouse neurological mutant stargazer (*stg*) identified a second γ subunit gene, *Cacng2*, expressed exclusively in the CNS (Letts et al. 1998). Expression of a single isoform in neurons distinguishes the γ subunit from other Ca²⁺ channel subunits, which utilize genetic heterogeneity as an important mechanism for generating functional diversity in these cells. In addition, the expression of significant levels of either *CACNG1* or *CACNG2* mRNA has not been reported in tissues such as heart, kidney, and testis, which express high levels of other Ca²⁺ channel subunits and produce measurable Ca²⁺ currents (Jay et al. 1990; Letts et al. 1998). We therefore hypothesized the existence of additional γ subunit genes. The low level of amino acid identity between the γ_1 and γ_2 proteins suggested that novel γ subunit paralogs might be difficult to identify using gene isolation methods dependent only on nucleic acid hybridization, such as low-stringency cross-hybridization to known γ subunit cDNA fragments or PCR amplification between conserved domains using degenerate oligonucleotides. An alternative approach based on the use of similarity search algorithms to screen genome-wide sequence databases for homologous genes can be utilized, but this sometimes produces large numbers of ambiguous identifications when low levels of homology are predicted and low-stringency search criteria are employed. We hy-

²Corresponding author.

E-MAIL dburgess@bcm.tmc.edu; FAX (713) 798-7528.

pothesized that a modification of genome-wide database searching might prove useful under these conditions. To test this, we applied a search paradigm based on sequence similarity analyses but restricted to small genomic regions predicted by gene duplication models as likely locations of unidentified γ subunit genes.

The expansion of gene families through evolution is thought to rely on two principal mechanisms of gene duplication (Ohno 1970; Nadeau and Sankoff 1997). Tandem duplication generates paralogs that often remain in close proximity on the same chromosome. Chromosome or whole-genome duplication results in the simultaneous duplication of many genes, which retain their initial order on paralogous daughter chromosomes. Both models were used to predict the most likely locations of additional γ subunit genes. We then performed an extensive low-stringency comparative analysis of *CACNG1* and *CACNG2* cDNA and amino acid sequences to all available genomic sequences localized to the predicted regions. We report the identification of three novel Ca^{2+} channel γ subunit genes, *CACNG3*, *CACNG4*, and *CACNG5*, on chromosomes 16 and 17. Phylogenetic analysis supports a complex model of γ subunit gene family evolution requiring a minimum of two ancient tandem duplications that preceded at least two chromosome duplication events. The identification of expressed sequences in the brain corresponding to *CACNG3* and *CACNG4* suggests that the γ subunit, like the α_1 , β , and $\alpha_2\delta$ subunits, regulates Ca^{2+} currents in the CNS from multiple genetic loci.

RESULTS

Three Novel Members of the Ca^{2+} Channel γ Subunit Gene Family

The Ca^{2+} channel γ subunit genes *CACNG1* and *CACNG2* are located on chromosome bands 17q24 and 22q12–q13, respectively (Iles et al. 1993; Powers et al. 1993; Letts et al. 1998). We reasoned that any unidentified paralogous genes generated by tandem duplication would most likely have remained close to *CACNG1* and *CACNG2* in these regions throughout evolution. Chromosome band 16p11–p13 was the only additional genomic region we identified that contained several genes with paralogs on 17q11–q25 and 22q11–q24 (Giles et al. 1998) and was therefore a good candidate location for γ subunit genes created by ancient chromosome or whole-genome duplications. A target sequence database was constructed that contained only those human genomic sequences from the GenBank database that could be localized unambiguously to the paralogous chromosome bands 17q11–q25, 22q11–q24, and 16p11–p13 (Methods). The estimated number of nonredundant sequence residues contained in this target region database ($\sim 1.5 \times 10^7$) comprised <0.6% of the total number of sequence resi-

dues contained in the concurrent release of GenBank (release 110.0; 2.57×10^9).

Low-stringency similarity searches of the target database with human *CACNG1* and mouse *Cacng2* sequences identified exons of the genes *CACNG1* (17q24) and *CACNG2* (22q12–q13), as expected. Several additional related sequences were identified that were clearly distinct from *CACNG1* and *CACNG2* but were organized into similar gene structures. One of these putative genes was identical to a gene product located on chromosome band 16p12–p13.1 that had been predicted previously by automated gene identification programs associated with large-scale genome sequencing efforts (GenBank accession no. AAC15246). Because this gene shared significant sequence similarity with the Ca^{2+} channel γ_1 and γ_2 subunits, was organized into an intron–exon configuration identical to *CACNG1* and *CACNG2*, and was located within a paralogous chromosome region, we tentatively designated it *CACNG3* as a novel member of this gene family. The remaining sequences with similarity to *CACNG1* and *CACNG2* were all located on chromosome 17 and were derived from two partially overlapping bacterial artificial chromosome (BAC) clones that also contained the *CACNG1* gene. These sequences were organized into two putative genes that were nearly identical in structure to *CACNG1*, *CACNG2*, and *CACNG3* and were designated *CACNG4* and *CACNG5* (Fig. 1).

The existing annotation of *CACNG3* as an unknown gene product within its larger genomic sequence database entry distinguished it from *CACNG4* and *CACNG5* sequences, which were not annotated as potential genes. This difference reflects the fact that a subset of sequencing centers do not routinely perform or report gene identification analysis of large genomic sequences using methods such as XGRAIL, Genefinder, and Genscan. To determine if these analyses would have predicted *CACNG4* and *CACNG5*, we analyzed the relevant genomic sequences (GenBank accession nos. AC005544 and AC005988) with the Genscan program (Burge and Karlin 1997, 1998). Using default parameters, Genscan predicted the structure of both *CACNG4* and *CACNG5*. The accuracy was generally high, although variable among different exons. The borders of exons 1, 2, and 3 were predicted exactly as shown in Figure 1, while the Genscan-predicted end of exon 4 was 6 bp short for *CACNG4* and 321 bp short for *CACNG5*. *P* values were >0.99 for each of the predicted exons except exon 4 of *CACNG4* (*P* = 0.128) and exon 1 of *CACNG5* (*P* = 0.425). Genscan also predicted the promoter region of both genes upstream of the first exon.

Identification of Expressed Sequences from *CACNG3*, *CACNG4*, and *CACNG5*

Although comparisons based on sequence, gene




Figure 1 (A) Amino acid alignment of the voltage-dependent Ca^{2+} channel γ_{1-5} subunits. The relative positions of the three introns are indicated by dots, and the adjacent exons are numbered 1–4. The location of putative transmembrane domains predicted by the program TMpred (Hofmann and Stoffel 1993) are underlined (see Fig. 2). Dashes indicate gaps introduced to maintain optimal alignment. HG1, 2, 3, 4, and 5 were translated from sequences of the human genes *CACNG1*–*CACNG5*, respectively, with conceptual splicing at positionally conserved splice sites. Consensus *N*-glycosylation sites are double-underlined. Potential phosphorylation sites, indicated by carets (^), are consensus targets for one or more of the following: cAMP/cGMP-dependent protein kinase (Prosite PDOC00004), protein kinase C (PDOC00005), casein kinase II (PDOC00006), and tyrosine kinase (PDOC00007). Potential protein kinase C phosphorylation sites at amino acids 50 and 51 of HG2 are not marked. A single nontransmembrane region of well-conserved amino acid sequence is indicated by diamonds (◆). (B) Comparison of the intron–exon splice junctions and intron sizes of the human *CACNG1*–5 genes (HG1–5). Exon sequences are in uppercase letters; intron sequence in lowercase. Consensus splice acceptor and donor motifs (Mount 1982) are indicated at the bottom. (C) Alignment of *CACNG1*–5 translation initiation sites. The consensus translation initiation motif (Kozak 1984) is shown at the bottom. The putative first methionine codon is underlined. The sequences of the *CACNG2* and *CACNG3* genes are identical for 9 nucleotides preceding the start codon; however, this sequence is a poor match to the consensus (thymine is the most uncommon residue at positions –1 and –2 of vertebrate translation start sites, 9% and 11%, respectively). (M) A or C; (R) purine; (Y) pyrimidine; (N) any base.

structure, open reading frame, and chromosome location supported the inclusion of *CACNG3*, *CACNG4*, and *CACNG5* in the Ca^{2+} channel γ subunit gene

family, we sought additional evidence that these loci encoded functional genes rather than pseudogenes. The identification of several expressed sequence tags (ESTs) representing *CACNG3* and *CACNG4* was consistent with transcription and splicing of these genes as predicted by the genomic sequence motifs. ESTs corresponding to the 5' UTR (GenBank accession nos. H38324 and T07086) and 3' UTR (H38292, T23680, H04803, and H11477) of *CACNG3* were identified by sequence similarity searches of GenBank. Three of the four 3' ESTs terminated in poly(A) sequences 31 bp downstream of a consensus polyadenylation motif, ATTTAA. Three additional ESTs spanned the coding region of *CACNG3* (W29095, H11833, and H04905) and confirmed the splicing of all three predicted introns. The mRNA source of the ESTs was fetal adult brain tissue (except for a single cDNA derived from adult retina) and suggested that *CACNG3* was expressed in neurons or glia. *CACNG4* was also represented in GenBank by multiple human ESTs, corresponding to the 3' UTR (M78316 and AI207906) and the protein coding region (AA970202, AI423159, and AI146595). The sequence of one EST spanned exons 3 and 4 of *CACNG4* and confirmed that mRNA from this gene was also spliced as predicted. The *CACNG4* ESTs were derived from fetal brain, glioblastoma, and oligodendroglioma cDNA libraries, suggesting that this gene, like *CACNG3*, was also expressed in neurons or glia. We did not identify any EST or cDNA sequences in GenBank or other sequence databases corresponding to *CACNG5*. To determine if this gene was expressed, we generated oligonucleotide primers corresponding to exon 3 and exon 4 of the *CACNG5* genomic sequence and screened cDNA libraries by PCR. A single product of the expected size was amplified from a human fetal kidney cDNA library. Sequencing of this product demonstrated that it was identical to the predicted spliced cDNA of *CACNG5*, and confirmed that this gene was transcribed and the mRNA was processed (GenBank accession no. AF148220).

We further sought to determine if sequence from the promoter regions of *CACNG3*, *CACNG4*, and *CACNG5* contained consensus transcription factor binding motifs that might be useful for predicting tissue-specific expression patterns. However, although web-based promoter analysis programs (Methods) were successful in identifying numerous potential binding sites for various proteins, we did not identify any patterns that predicted the preferred transcription of these genes in specific tissues or in response to specific stimuli (data not shown).

Phylogenetic Relationships

Although the skeletal muscle γ_1 and neuronal γ_2 sub-

units exhibited low amino acid identity (~25%), the predicted transmembrane topologies were nearly indistinguishable (Letts et al. 1998), suggesting strong selective constraints on this aspect of secondary structure. We extended this analysis to include the γ_3 , γ_4 , and γ_5 subunits. Examination of all five isoforms predicted very similar transmembrane topologies for γ_1 , γ_2 , γ_3 , γ_4 , and γ_5 (Fig. 2). The presence of greater amino acid identity within the putative transmembrane domains, as compared to other regions of the protein, was consistent with selective conservation of sequence identity in these regions (Fig. 1). Comparison of the hydrophobicity plots indicated that γ_2 , γ_3 , and γ_4 were more similar to each other in secondary structure than to γ_1 or γ_5 , and that γ_5 was intermediate in structure between γ_1 and the others. This was somewhat unexpected, because *CACNG2*, *CACNG3*, and *CACNG4* are each located on different chromosomes (chromosomes 22, 16, and 17, respectively), and *CACNG4* is located between *CACNG1* and *CACNG5* on chromosome 17.

To clarify the evolutionary relationship among these genes, a phylogenetic analysis of γ subunit




Figure 2 Comparison of transmembrane topologies of Ca^{2+} channel γ_{1-5} subunits predicted by the TMpred program (Hofmann and Stoffel 1993). Positive TMpred values (y-axis, $\times 1000$) indicate likely membrane spanning segments. Amino acid position is shown on the x-axis. All five γ subunit isoforms are predicted to contain four transmembrane domains with amino and carboxyl termini located intracellularly.




Figure 3 (A) Molecular phylogeny of the Ca^{2+} channel γ subunit family. The alignment used to infer the tree was done independently for full-length proteins and for conserved regions alone (defined as translation start to the final residue of the fourth predicted transmembrane domain), and resulted in identical topologies. The number of trees with a particular node among 10,000 bootstrap replicates is indicated at the node (values for conserved domain trees shown above the horizontal line and for full length proteins below). A member of the related Claudin protein family, Claudin 4, was included in the comparison and defined as the out-group. Branch lengths are arbitrary and do not correspond to genetic distances. This tree is unrooted. (B) Pairwise amino acid identity among the Ca^{2+} channel γ_{1-5} subunits and mouse Claudin 4. Percent identity determined after alignment by the BLAST2 program is shown above the horizontal and by the ALIGN program below the horizontal. Homology between Claudin 4 and the γ subunits was below the threshold of detection by BLAST2. (MC4) Mouse Claudin 4; HG1, 3, 4, and 5 refer to human Ca^{2+} channel γ subunits; MG2 refers to the mouse γ_2 subunit; (n.a.), not applicable.

amino acid sequences was conducted under the assumption of maximum parsimony (Fig. 3A). The mouse protein Claudin 4, which we determined to be distantly related to the Ca^{2+} channel γ subunits by comparative sequence analysis, was defined as the out-group. The recently identified Claudin proteins comprise a family of four-transmembrane-domain proteins believed to be important in the formation of tight junctions (Morita et al. 1999). The topology of the inferred tree was consistent with the hydrophobicity analysis and strongly supported a close relationship between γ_2 and γ_3 . The remaining branchpoints were also inferred with high confidence levels and indicated the branching of γ_4 , γ_5 , and γ_1 in reverse chronological order from the $\gamma_2 - \gamma_3$ node. These relationships were

concordant with the levels of pairwise amino acid identity among the proteins (Fig. 3B).

Our data suggested a model of Ca^{2+} channel γ subunit gene family evolution in which at least two ancient tandem gene duplications preceded the chromosome duplication events that led to the modern chromosome regions 17q11–q25, 22q11–q24, and 16p11–p13 (Fig. 4). The phylogenetic clustering of γ_2 (chromosome 22) with γ_3 (chromosome 16), and their more distant relationship to γ_4 (chromosome 17), could be interpreted as evidence of more recent divergence between chromosomes 22 and 16. A logical extrapolation would be that other paralogous genes on chromosomes 16 and 22 would also be more closely related to each other, on average, than to any paralogs on chromosome 17. To investigate this hypothesis further, we examined sequences immediately surrounding the γ subunit genes on these three chromosomes. Several additional genes were identified, but comparisons among these failed to support a more recent divergence between chromosomes 16 and 22. In fact, the presence of paralogous protein kinase C genes (*PRKCB1* and *PRKCA*) immediately telomeric of *CACNG3* and *CACNG5*, respectively, and the absence of a *PRKC* paralog telomeric of *CACNG2* on chromosome 22, sup-

ported a greater similarity between chromosomes 16 and 17 in these regions (Fig. 5). To resolve this ambiguity, we carried out a comprehensive comparison of paralogous genes located on chromosome bands 17q11–q25, 22q11–q24, and 16p11–p13, including some novel loci that were identified through analysis of our regionally restricted target sequence database (Table 1). However, although the data demonstrated a clear relationship among all three chromosome regions, their ancestral relationships remained equivocal and additional studies will be needed to clarify this issue.

DISCUSSION

We applied two models of gene family expansion, tandem duplication and chromosome duplication, to facilitate the identification of three novel members of the Ca^{2+} channel γ subunit gene family: *CACNG3*, *CACNG4*, and *CACNG5*. The aim of this approach was to maximize the likelihood of correct gene identifications by low-stringency similarity searches of localized DNA sequences and reduce the large number of biologically irrelevant matches usually generated by genome-wide database analysis. The amino acid identity between the γ_1 subunit and the γ_3 , γ_4 , and γ_5 subunits was 22%–26%. This low degree of similarity may explain why the *CACNG3*, *CACNG4*, and *CACNG5* genes were previously undetected by standard whole-genome database searches, which often employ higher stringency alignment parameters as the default criteria for defining similarity. In contrast, the amino acid identity between the γ_2 subunit and the γ_3 , γ_4 , and γ_5 subunits was variable, measuring 84%, 64%, and 32%, respectively. Recently, Black and Lennon (1999) also identified the *CACNG3* gene by computer similarity searches of genomic sequence databases using the human *CACNG2* sequence as the query. The high degree of similarity between γ_2 and γ_3 (84%), and the pre-existing GenBank annotation of γ_3 as an unknown gene product, may have facilitated identification of *CACNG3* using standard search parameters. The fact that the *CACNG4* and *CACNG5* genes were not detected by that approach, although they are both located within 100 kb of *CACNG1*, demonstrates the value of gene duplication models for predicting gene location and improving gene identification efficiency. The additional observation that both *CACNG4* and *CACNG5* would have been predicted by gene identification software such as Genscan, underscores the importance of improved genomic sequence annotation.

Although our approach was successful in identifying *CACNG3*, *CACNG4*, and *CACNG5*, there are important limitations to its efficacy. Foremost, many paralogous members of gene families are not located in tandem or in duplicated chromosomal regions that exhibit conserved gene order with other family mem-




Figure 4 A model of Ca^{2+} channel γ subunit gene family evolution. Integration of chromosome locations and molecular phylogeny data suggest a minimum of two tandem duplications of an ancestral γ subunit gene, followed by two duplications of the precursor of modern chromosome bands 16p11–p13, 17q11–q25, and 22q11–q24. Four potential genes suggested by the model (C' , C'' , A' , and A'') but not identified in this study may have been lost during evolution sometime after the duplication indicated by arrow 3. Arrows 1 and 2 indicate tandem gene duplication events. Arrows 3 and 5 indicate regional (chromosome or whole genome) duplications. Letters (A, B, C) and symbols (' , '') designate inferred ancestors of modern γ subunit genes.




Figure 5 (A–C) Physical maps of the region surrounding the human Ca^{2+} channel γ subunit genes. Genomic sequences are indicated by thin lines below the name of the associated clone, with selected markers shown below. (●) Sequence overlap; (○) end sequences associated only with a single parent clone in GenBank. Characterized genes are represented by thick lines, with transcriptional orientation indicated by arrowheads. *PRKCB1* is interrupted by a gap in the genomic sequence of unknown length. Only the 3' ends of *PRKCA* and *ORFA05* are currently represented in GenBank. (*CBP-P22*) calcineurin-related gene; (*PRKCB1*) protein kinase C β 1; (*CACNG3*) calcium channel γ 3; *CSF2RB* colony-stimulating factor 2 receptor β ; (*NCF4*) neutrophil cytosolic factor 4; (*PVALB*) parvalbumin; (*HSRASR*) similar to *H. sapiens RAY1* gene; (*CACNG2*) calcium channel γ 2; (*HPSP1*) Hermansky–Pudlak syndrome pseudogene; (*EIF3-P66*) eukaryotic translation initiation factor 3, subunit 7 (ζ , 66/67 kD); (*TRX2*) thioredoxin 2; (*PRKCA*) protein kinase C α ; (*G6PDP1*) formerly *G6PDL*, glucose-6-phosphate dehydrogenase pseudogene 1; (*CACNG5*) calcium channel γ 5; (*CACNG4*) calcium channel γ 4; (*CACNG1*), calcium channel γ 1; (*ORFA05*) hypothetical myeloid cell line protein 5. StsG25737 represents an EST of *CACNG3*. Complete clone names and database accession numbers are given in Methods. (D–F) Comparison of the intron–exon structure of *CACNG1–5*. The scale bar in D applies to D–F.

bers. Some of these paralogs may have been generated by gene duplication mechanisms not considered in the model we employed. For example, genes duplicated by retrotransposition via an mRNA intermediate could theoretically insert anywhere in the genome with respect to the parental gene. In most cases, however, it seems probable that complex genomic rearrangements occurring over large time scales have obliterated the initial positional relationships among distantly related genes. Predictions of paralogous gene locations based on common duplication models will, by design, exclude such genes. This approach should therefore be considered primarily as a complement to other sequence-based gene identification techniques.

Role of γ Subunits in Ca^{2+} Channel Function

Two of the three genes identified in this study, *CACNG3* and *CACNG4*, were represented by several ESTs derived from brain mRNA. We found expressed sequences corresponding to *CACNG5* by PCR amplifi-

cation of a human fetal kidney cDNA library but did not exclude the possibility of *CACNG5* expression in the brain or other tissues. It is worth noting that several tissues that express multiple α_1 , β , and $\alpha_2\delta$ subunit isoforms, including testes, ovary, lung, pancreas, spleen, liver, and kidney, do not express γ_1 or γ_2 (Biel et al. 1990; Jay et al. 1990; Castellano and Perez-Reyes 1994; Yu 1995; Williams et al. 1999). The γ_3 , γ_4 , and γ_5 isoforms are possible components of Ca^{2+} channels in these tissues. The coexpression of multiple isoforms of α_1 , β , and $\alpha_2\delta$ subunits in individual neurons is a valuable mechanism for generating functional variability among Ca^{2+} channels in brain, and our data suggest that the γ subunit could contribute to channel diversity in a similar manner. For example, the mouse *Cacng2* gene is widely expressed in the brain (Letts et al. 1998) and may be coexpressed in some regions with the mouse homologs of *CACNG3* and *CACNG4*. If this is confirmed, then it will be important to examine the relative contributions of each γ subunit isoform to the structure and function of distinct Ca^{2+} channel types in vivo. A comprehensive comparative analysis of γ subunit gene expression in the brain and other tissues will provide insight into the physiological role of these isoforms. Confirmation will ultimately require the demonstration that the predicted γ_3 , γ_4 , and γ_5 proteins can directly modulate channel biophysical properties, or influence the stability or subcellular localization of the channel complex.

Less is understood about the function of the Ca^{2+} channel γ subunit than about α_1 , β , and $\alpha_2\delta$. Coexpression of the cardiac α_1 subunit (α_{1C}) with the skeletal muscle γ_1 isoform was used to demonstrate that it shifted the inactivation curve of the channel to negative potentials and accelerated current inactivation without significantly affecting other voltage-dependent properties (Singer et al. 1991; Eberst et al. 1997). Another study indicated that the γ_1 subunit did not have a significant effect on α_{1C} -mediated channel currents unless coexpressed with a β subunit (Wei et al. 1991). Coexpression analysis of the γ_2 subunit, which is disrupted in the mouse neurological mutant *stg*, showed that it increased the steady-state inactivation of α_{1A} -containing Ca^{2+} channels (Letts et al. 1998). Sequence similarity to γ_1 and γ_2 supports a prediction that the γ_3 , γ_4 , and γ_5 proteins may also regulate the inactivation properties of Ca^{2+} channels. In general, the effects of γ subunit regulation on Ca^{2+} currents that have been described are small in magnitude (Walker and De Waard 1998). However, if modulation of channel properties is dependent on the coex-

Table 1. Paralogous Genes Located Within Duplicated Regions on Chromosomes 16p11–p13, 17q11–q25, and 22q11–q24

gene	Chromosome 16			Chromosome 17			Chromosome 22		
	RH map	cytogenetic	gene	RH map	cytogenetic	gene	RH map	cytogenetic	
ARHGDI _G	19.85	16p13.3	ARHGDI _A	537.27	17q25.3	CSNK1E	131.36	22q12–q13	
SSTR5	25.14	16p13.3	CSNK1D	501.76–539.58	17q25	SSTR3	111.72–124.01	22q13.1	
CACNA1H	44.93–65.94	16p13.3	SSTR2	460.32	17q24	CACNA1I	137.86–138.61	22q12.3–q13.2	
SYNGR3	65.01–65.33	16p13.3	CACNA1G	347.18–351.66	17q22	SYNGR1	138.19	22q13	
RPL3L	65.01	16p13.3	SYNGR2	N.D.	17qtel	RPL3	138.19	22q13	
CREBBP	62.52–68.90	16p13.3	SOX9	448.88–466.09	17q24.3–q25.1	EP300	N.D.	22q13.2	
GRIN2A	95.4–100.92	16p13.2	GRIN2C	N.D.	17q24–q25	SOX10	115.15	22q13	
PMM2	N.D.	16p13.3–p13.2				PMM1	126.67–138.78	22q13.1	
TOP3L**	191.15	16p11.2				TOP3B	N.D.	22q11	
CRYM	191.25–192.97	16p13.11–p12.3	CRYBA1	N.D.	17q11.1–q12	CRYBA4	72.13–76.42	22q12.1	
						CRYBB1	72.13–76.42	22q11.2–q12.1	
						CRYBB2	72.13–76.42	22q11.2–q12.1	
						CRYBB3	72.13–76.42	22q11.2–q12.1	
PRKCB1	197–200.56	16p11.2	PRKCA	421.67	17q22–q23.2	CACNG2	112	22q12–q13	
CACNG3**	197	16p12–13.1	CACNG5**	421.16	17q24				
			CACNG4**	421.16	17q24				
			CACNG1	421.16	17q24				
TBX6	N.D.	16p11.2	TBX2	371.45	17q23	TBX1	N.D.	22q11.2	
			TBX4*	385.31	17q23				
			TBRL**	338.95	17q23				
			CLTC	376.31–377.45	17q11–qter	CLTCL	N.D.	22q11.2	

RH map values refer to 3000 cR on the Genebridge4 (GB4) radiation hybrid mapping panel (Gyapay et al. 1996; Deloukas et al. 1998; GeneMap '98). Cytogenetic locations are from OMIM. (*) Previously identified but not mapped. (**) Novel genes identified in this study. (ARHGDI) Rho gdp-dissociation inhibitor; (CSNK) casein kinase; (SOX) SRY (Sex determining region Y)-box; (SSTR) somatostatin receptor; (SYNGR) synaptogyrin; (RPL3L) ribosomal protein, L3-like; (RPL3) ribosomal protein L3; (CREBBP) CREB-binding protein; (EP300) E1A binding protein p300; (CACNA) voltage-gated calcium channel; (GRIN) ionotropic glutamate receptor; (PMM) phosphomannomutase; (TOP) topoisomerase; (CRY) crystallin; (PRKC) protein kinase C; (TBX) T-box transcription factor; (TBR) T-box brain protein; (CLT) clathrin; (N.D.) not determined. New gene symbols were approved by the HUGO/GDB Nomenclature Committee.

pression of specific α_1 and γ isoforms, the experimental results described above may not accurately represent γ subunit function in vivo. For example, the skeletal muscle γ_1 isoform is not expressed at high levels in heart with the cardiac α_{1C} isoform, and it is not known whether the γ_2 isoform actually associates with α_{1A} in the brain, although both are widely coexpressed (Tanaka et al. 1995; Letts et al. 1998). Instead, it is possible that other γ subunit isoforms associate preferentially with α_{1C} and α_{1A} in vivo. Functional coexpression of γ_3 , γ_4 , and γ_5 in combination with various α_1 , β , and $\alpha_2\delta$ isoforms in vitro may illustrate distinct regulatory functions for the γ subunit, whereas coimmunoprecipitation analysis of different tissues or brain regions will be helpful in determining which isoforms are associated preferentially in vivo. It is also worth noting that Ca^{2+} channel γ subunits exhibit a low level of amino acid identity and similar hydrophobicity profiles to several Claudin proteins, to the lens intrinsic membrane protein MP20, and to peripheral myelin protein PMP22 (data not shown). However, it is not known whether any functional similarities exist among the members of this extended family of four-transmembrane domain proteins.

The chromosome locations of the γ subunit genes, *CACNG3*, *CACNG4*, and *CACNG5*, suggests they could be candidates for involvement in hereditary disease. *CACNG3* is located on chromosome band 16p12–p13.1 in the vicinity of the *ICCA* locus [infantile convulsions and paroxysmal choreoathetosis; Online Mendelian Inheritance in Man (OMIM) no. 602066]. *CACNG3* is expressed in the brain and it is worth noting that mutation of the closely related *Cacng2* gene in the *stg* mouse results in epilepsy and ataxia (Noebels et al. 1990; Letts et al. 1998). The mouse ortholog of *CACNG3* is predicted to map to chromosome 7 near *Szv2*, a quantitative trait locus (QTL) influencing seizure response to kainic acid (Ferraro et al. 1997). *CACNG4* and *CACNG5* are located on chromosome band 17q24 in tight physical linkage to *CACNG1*. A locus for neuralgic amyotrophy with brachial predilection (*NAPB*; OMIM 162100) has been mapped to 17q24–q25 and is characterized by severe pain, weakness, wasting, depression of reflexes, and sensory loss (Jacob et al. 1961). However, *NAPB* was localized close to marker *D17S939* (Pellegrino et al. 1997), whereas *CACNG1* (and *CACNG4* and *CACNG5* by association) was significantly more centromeric, near *D17S807*, and is therefore an unlikely candidate for this disorder. Comparison of conserved linkage groups suggests that mouse *Cacng1*, *Cacng4*, and *Cacng5* are probably located near *Pkca*, which is on the consensus map of chromosome 11 at 68 cM [Mouse Genome Informatics (MGI) database]. Interestingly, this position is near a second locus for seizure susceptibility, *Szs3*, at 66 cM (Ferraro et al. 1997). The close association of epilepsy and ataxia with mutations

in other neuronal voltage-dependent Ca^{2+} channels (Burgess and Noebels 1999) suggests these are potential candidate phenotypes for defects in the *CACNG3*, *CACNG4*, or *CACNG5* genes.

METHODS

Target Region Sequence Database Construction

A database of human genomic sequences derived from chromosome bands 17q11–q25, 22q11–q24, and 16p11–p13 (target regions) was constructed using Microsoft Excel '97. Sequences were compiled from two sources: the Human Genome Sequencing Index (HGSI) database contains sequences from large clones (cosmids, BACs, or PACs), and clone contigs that have been localized unambiguously to specific genomic regions. Additional sequences were obtained by screening GenBank for genomic sequences identical to genes or cDNA that were localized previously to the target regions, using the programs BLASTN or TBLASTN (release 2.0; Altschul et al. 1997). Approximately 200–400 bp of nonrepetitive sequence from the ends of each genomic sequence obtained in this way was used for additional rounds of database screening and sequence contig extension, terminating when no additional sequences were identified.

Electronic Database Information

Data presented are consistent with the following databases as of September 1999: GenBank (GB), <http://www.ncbi.nlm.nih.gov/Web/Genbank/>; GeneMap'98, <http://www.ncbi.nlm.nih.gov/genemap/>; Genestream, <http://vega.crbm.cnrs-mop.fr/home.html>; Genscan, <http://gnomic.stanford.edu/~chris/GENSCANW.html>; HUGO/GDB Human Gene Nomenclature, <http://www.gene.ucl.ac.uk/nomenclature/>; LocusLink, <http://www.ncbi.nlm.nih.gov/LocusLink/>; Human Genome Sequencing Index (HGSI), <http://www.ncbi.nlm.nih.gov/HUGO/>; Mouse Genome Informatics (MGI), <http://www.informatics.jax.org/>; National Center for Biotechnology Information (NCBI), <http://www.ncbi.nlm.nih.gov/>; Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/Omim/>; Prosite, <http://www.expasy.ch/prosite/>.

Accession Numbers

Genes referred to in the text and figures are listed in alphabetical order, followed by genomic sequences listed by chromosome. Genes identified within larger genomic sequences are identified by base-pair (bp) position in the genomic sequence. All accession numbers refer to GenBank unless otherwise indicated: *CACNG1* (AC005544), *CACNG2* (Z83733, AL022313, AL031845), *CACNG3* (AC004125), *CACNG4* (AC005544, AC005988, AF142622, AF142623, AF142624, AF142625), *CACNG5* (AC005988, AF142618, AF142619, AF142620, AF142621, AF148220), *CBP-P22* (2695572), *CSF2RB* (OMIM 138981), *EIF3-P66* (U54558), *G6PDP1* (M12996; AC005988, bp 8473–9248), *HSRASR* (AL022729), *HPSP1* (U65676; AL022313, bp 90054–88792), *NCF4* (AH004909), *ORFA05* (D29677), *PRKCA* (X52479), *PRKCB1* (M13975), *PVALB* (OMIM 168890), *TRX2* (U78678). Chromosome 16 genomic sequences: CIT987–SKA–345G4 (AC002302), CIT987–SKA–113A6 (AC002299), CIT987SK–625P11 (AC004125). Chromosome 17 genomic sequences: hCIT.187_K_10 (AC006263), hRPK.115_C_3 (AC006947), hRPK.74_H_8 (AC005918), hRPK.299_G_24 (AC005988),

hRPK.349_A.8 (AC005544). Chromosome 22 genomic sequences: E132D12 (Z80897), 833B7 (AL008637), 24E5 (Z82185), 566H6 (AL031845), 1119A7 (AL022313), 126G10 (Z82184), 293L6 (Z82197; Z83733; Z83732), 4G12 (Z70289).

Low Stringency Similarity Searches

Low-stringency similarity searches for novel Ca²⁺ channel γ subunit genes were limited to sequences contained in the target region database. The BLASTN program (NCBI) was utilized for pairwise comparisons between large genomic sequences (10–150 kb) and cDNA sequences. Default filters used to mask sequences of low compositional complexity were turned off. The BLOSUM62 alignment scoring matrix was used. Alignment parameters were adjusted to reduce stringency (default value, value used): expectation value, $-e$ (10.0, 100.0); gap-opening penalty, $-G$ (5, 3); gap-extension penalty, $-E$ (2, 1); mismatch penalty in the blast portion of the run, $-q$ (–2, –1); word size (11, 7).

Sequence Analysis of Promoter Regions

Analysis of promoter region sequence for transcription factor binding sites utilized the web-based programs: PatSearch 1.1, utilizing the TRANSFAC 3.4 and TRRD 3.5 databases (Heinemeier et al. 1998, 1999); MatInspector Version 2.2 (Quandt et al. 1995), utilizing the TRANSFAC 3.5 database; TFSEARCH (Yutaka Akiyama, <http://www.rwcp.or.jp/papia/>), utilizing the TRANSFAC 3.3 database; and TESS (J. Schug and G. Christian Overton, <http://www.cbil.upenn.edu/tess/>), utilizing the TRANSFAC 3.3 database.

Multiple Sequence Alignments and Phylogenetic Analysis

Protein sequences were aligned for phylogenetic analysis using the ClustalX multiple alignment package (Thompson et al. 1997) with default values. Alignments were carried out using full length sequences or only conserved regions as indicated in the text. Pair-wise percent amino acid identity was calculated following local alignment by BLASTP (release 2.0) using the BLOSUM62 scoring matrix and gap opening/extension penalties of 8 and 2, and following global alignment with the ALIGN program (Genestream) using the codaa.mat scoring matrix and gap opening/extension penalties of 12 and 2. The sequence alignment shown in Figure 1 was manually edited for display but not for phylogenetic analysis or amino acid identity calculations. Phylogenetic relationships were inferred using the neighbor-joining method (Saitou and Nei 1987) of the ClustalX multiple alignment package (Thompson et al. 1997). The reliability of tree topology was evaluated using bootstrap analysis (Felsenstein 1985) with 10,000 iterations to provide confidence levels. Unrooted trees were plotted as rectangular cladograms using the Tree-View program (Page 1996).

Identification of CACNG5 cDNA

Oligonucleotide PCR primers predicted to amplify a 283-bp cDNA product were designed according to the sequence of the CACNG5 gene: HG5-F (exon 3): 5'-GATACTGGCCTTTGTCTCTGG-3'; HG5-R (exon 4): 5'-TTGTGGAATGTC-CCTTCTCC-3'. A single product of ~283 bp was amplified from a PCR reaction containing 1 μ l of phage suspension from a human fetal kidney cDNA library (Clontech, HL5004a) as template in a 50- μ l reaction volume, 50 mM KCl, 10 mM Tris-HCl, 1.5 mM MgCl₂, 0.1% Triton X-100, 25 mM dNTPs, 0.8 μ M

each oligonucleotide (HG5-F and HG5-R), and 1 Unit of *Taq* polymerase (Promega). The reaction was carried out using a PTC-100 thermocycler (MJ Research) with an initial denaturation of 94°C for 2 min, followed by 35 cycles of 94°C (30 sec), 55°C (60 sec), 72°C (60 sec), and a final extension of 72°C for 10 min. Following agarose gel electrophoresis, the PCR product was isolated using the QIAquick gel extraction kit (Qiagen) and sequenced by the Baylor College of Medicine DNA Sequencing Core Facility.

ACKNOWLEDGMENTS

This research was supported by an American Epilepsy Society postdoctoral fellowship and a Methodist Hospital Foundation (Houston, TX) grant to D.L.B. and National Institutes Health grant NS29709 to J.L.N. We thank T. Cormier in the laboratory of Dr. Huda Zoghbi for providing a sample of the Clontech fetal kidney cDNA library.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

REFERENCES

- Altschul, S.F., T.L. Madden, A.A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D.J. Lipman. 1997. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.* **25**: 3389–3402.
- Biel, M., P. Ruth, E. Bosse, R. Hullin, W. Stuhmer, V. Flockerzi, and F. Hofmann. 1990. Primary structure and functional expression of a high voltage activated calcium channel from rabbit lung. *FEBS Lett.* **269**: 409–412.
- Black, J.L. III and V.A. Lennon. 1999. Identification and cloning of putative human neuronal voltage-gated calcium channel γ_2 and γ_3 subunits: Neurologic implications. *Mayo. Clin. Proc.* **4**: 357–361.
- Burge, C. and S. Karlin. 1997. Prediction of complete gene structures in human genomic DNA. *J. Mol. Biol.* **268**: 78–94.
- . 1998. Finding the genes in genomic DNA. *Curr. Opin. Struct. Biol.* **8**: 346–354.
- Burgess, D.L. and J.L. Noebels 1999. Voltage-dependent Ca²⁺ Channel Mutations in Neurological Disease. *J.N.Y. Acad. Sci.* (in press).
- Castellano, A. and E. Perez-Reyes. 1994. Molecular diversity of Ca²⁺ channel β subunits. *Biochem. Soc. Trans.* **22**: 483–488.
- Chin, H.M., C.A. Kozak, H.L. Kim, B. Mock, and O.W. McBride. 1991. A brain L-type calcium channel α_1 subunit gene (CCHL1A2) maps to mouse chromosome 14 and human chromosome 3. *Genomics* **11**: 914–919.
- Collin, T., P. Lory, S. Taviaux, C. Courtieu, P. Guilbault, P. Berta, and J. Nargeot. 1994. Cloning, chromosomal location and functional expression of the human voltage-dependent calcium-channel β_3 subunit. *Eur. J. Biochem.* **220**: 257–262.
- Cribbs, L.L., J.-H. Lee, J. Yang, J. Satin, Y. Zhang, A. Daud, J. Barclay, M.P. Williamson, M. Fox, M. Rees et al. 1998. Cloning and characterization of α_{1H} from human heart, a member of the T-type calcium channel gene family. *Circ. Res.* **83**: 103–109.
- Deloukas, P., G.D. Schuler, G. Gyapay, E.M. Beasley, C. Soderlund, P. Rodriguez-Tomé, L. Hui, T.C. Matise, K.B. McKusick, J.S. Beckmann et al. 1998. A physical map of 30,000 human genes. *Science* **282**: 744–746.
- De Waard, M., C.A. Gurnett, and K.P. Campbell. 1996. Structural and functional diversity of voltage-activated calcium channels. In *Ion Channels* (ed. T. Narahashi), pp 41–87. Plenum Press, New York, NY.
- Diriong, S., P. Lory, M.E. Williams, S.B. Ellis, M.M. Harpold, and S. Taviaux. 1995. Chromosomal localization of the human genes for alpha 1A, alpha 1B, and alpha 1E voltage-dependent Ca²⁺ channel subunits. *Genomics* **30**: 605–609.
- Drouet, B., L. Garcia, D. Simon-Chazottes, M.G. Mattei, J.-L. Guenet, A. Schwartz, G. Varadi, and M. Pincon-Raymond. 1993. The gene coding for the α -1 subunit of the skeletal dihydropyridine

- receptor (*Cchl1a3=mdg*) maps to mouse chromosome 1 and human 1q32. *Mamm. Genome* **4**: 499–503.
- Eberst, R., S. Dai, N. Klugbauer, and F. Hofmann. 1997. Identification and functional characterization of a calcium channel γ subunit. *Pflügers Arch.* **433**: 633–637.
- Felsenstein, J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* **39**: 783–791.
- Ferraro, T.N., G.T. Golden, G.G. Smith, N.J. Schork, P. St Jean, C. Ballas, H. Choi, and W.H. Berrettini. 1997. Mapping murine loci for seizure response to kainic acid. *Mamm. Genome* **8**: 200–208.
- Fisher, S.E., A. Ciccodicola, K. Tanaka, A. Curci, S. Desicato, M. D'urso, and I.W. Craig. 1997. Sequence-based exon prediction around the synaptophysin locus reveals a gene-rich area containing novel genes in human proximal Xp. *Genomics* **45**: 340–347.
- Giles, R.H., H.G. Dauwerse, G.J. van Ommen, and M.H. Breuning. 1998. Do human chromosomal bands 16p13 and 22q11–13 share ancestral origins? *Am. J. Hum. Genet.* **63**: 1240–1242.
- Gregg, R.G., F. Couch, K. Hogan, and P.A. Powers. 1993a. Assignment of the human gene for the α_1 subunit of the skeletal muscle DHP-sensitive Ca^{2+} channel (*CACNLIA3*) to chromosome 1q31–q32. *Genomics* **15**: 107–112.
- Gregg, R.G., P.A. Powers, and K. Hogan. 1993b. Assignment of the human gene for the β subunit of the voltage-dependent calcium channel (*CACNLB1*) to chromosome 17 using somatic cell hybrids and linkage mapping. *Genomics* **15**: 185–187.
- Gyapay, G., K. Schmitt, C. Fizames, H. Jones, N. Vega-Czarny, D. Spillett, D. Muselet, J.F. Prud'Homme, C. Dib, C. Auffray et al. 1996. A radiation hybrid map of the human genome. *Hum. Mol. Genet.* **5**: 339–346.
- Heinemeyer, T., E. Wingender, I. Reuter, H. Hermjakob, A.E. Kel, O.V. Kel, E.V. Ignatieva, E.A. Ananko, O.A. Podkolodnaya, F.A. Kolpakov et al. 1998. Databases on Transcriptional Regulation: TRANSFAC, TRRD, and COMPEL. *Nucleic Acids Res.* **26**: 264–370.
- Heinemeyer, T., X. Chen, H. Karas, A.E. Kel, O.V. Kel, I. Liebich, T. Meinhardt, I. Reuter, F. Schacherer, and E. Wingender. 1999. Expanding the TRANSFAC database towards an expert system of regulatory molecular mechanisms. *Nucleic Acids Res.* **27**: 318–322.
- Hofmann, K. and W. Stoffel. 1993. TMBASE—A database of membrane spanning protein segments. *Biol. Chem. Hoppe-Seyler* **374**: 166.
- Iles, D.E., B. Segers, D.O. Weghuis, R. Suikerbuijk, and B. Wieringa. 1993. Localization of the gamma-subunit of the skeletal muscle L-type voltage-dependent calcium channel gene (*CACNLG*) to human chromosome band 17q24 by in situ hybridization and identification of a polymorphic repetitive DNA sequence at the gene locus. *Cytogenet. Cell. Genet.* **64**: 227–230.
- Jacob, J.C., F. Andermann, and J.P. Robb. 1961. Heredofamilial neuritis with brachial predilection. *Neurology* **11**: 1025–1033.
- Jay, S.D., S.B. Ellis, A.F. McCue, M.E. Williams, T.S. Vedvick, M.M. Harpold, and K.P. Campbell. 1990. Primary structure of the gamma subunit of the DHP-sensitive calcium channel from skeletal muscle. *Science* **248**: 490–492.
- Klugbauer, N., L. Lacinova, E. Marais, M. Hobom, and F. Hofmann. 1999. Molecular diversity of the calcium channel $\alpha_2\delta$ subunit. *J. Neurosci.* **19**: 684–691.
- Kozak, M. 1984. Compilation and analysis of sequences upstream from the translational start site in eukaryotic mRNAs. *Nucleic Acids Res.* **12**: 857–872.
- Lee, J.H., A.N. Daud, L.L. Cribbs, A.E. Lacerda, A. Pereverzev, U. Klockner, T. Schneider, and E. Perez-Reyes. 1999. Cloning and expression of a novel member of the low voltage-activated T-type calcium channel family. *Neuroscience* **19**: 1912–1921.
- Letts, V.A., R. Felix, G.H. Biddlecome, J. Arikath, C.L. Mahaffey, A. Valenzuela, F.S. Bartlett II, Y. Mori, K.P. Campbell, and W.N. Frankel. 1998. The mouse stargazer gene encodes a neuronal Ca^{2+} -channel gamma subunit. *Nat. Genet.* **19**: 340–347.
- Morita, K., M. Furuse, K. Fujimoto, and S. Tsukita. 1999. Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands. *Proc. Natl. Acad. Sci.* **96**: 511–516.
- Mount, S.M. 1982. A catalogue of splice junction sequences. *Nucleic Acids Res.* **10**: 459–472.
- Nadeau, J.H. and D. Sankoff. 1997. Comparable rates of gene loss and functional divergence after genome duplications early in vertebrate evolution. *Genetics* **147**: 1259–1266.
- Noebels, J.L., X. Qiao, R.T. Bronson, C. Spencer, and M.T. Davisson. 1990. Stargazer: A new neurological mutant on chromosome 15 in the mouse with prolonged cortical seizures. *Epilepsy Res.* **7**: 129–135.
- Ohno, S. 1970. *Evolution by gene duplication*. Springer Verlag, Heidelberg, Germany.
- Page, R.D.M. 1996. TREEVIEW: An application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**: 357–358.
- Pellegrino, J.E., R.A.V. George, J. Biegel, M.R. Farlow, K. Gardner, J. Caress, M.J. Brown, T.R. Rebeck, T.D. Bird, and P.F. Chance. 1997. Hereditary neuralgic amyotrophy: Evidence for genetic homogeneity and mapping to chromosome 17q25. *Hum. Genet.* **101**: 277–283.
- Perez-Reyes, E., L.L. Cribbs, A. Daud, A.E. Lacerda, J. Barclay, M.P. Williamson, M. Fox, M. Rees, and J.H. Lee. 1998. Molecular characterization of a neuronal low-voltage-activated T-type calcium channel. *Nature* **391**: 896–900.
- Powers, P.A., R.G. Gregg, P.A. Lalley, M. Liao, and K. Hogan. 1991. Assignment of the human gene for the α_1 subunit of the cardiac DHP-sensitive Ca^{2+} channel (*CCHLIA1*) to chromosome 12p12–pter. *Genomics* **10**: 835–839.
- Powers, P.A., S. Liu, K. Hogan, and R.G. Gregg. 1993. Molecular characterization of the gene encoding the γ subunit of the human skeletal muscle 1,4-dihydropyridine-sensitive Ca^{2+} channel (*CACNLG*), cDNA sequence, gene structure, and chromosomal location. *J. Biol. Chem.* **268**: 9275–9279.
- Powers, P.A., S.W. Scherer, L.C. Tsui, R.G. Gregg, and K. Hogan. 1994. Localization of the gene encoding the $\alpha_2\delta$ subunit (*CACNL2A*) of the human skeletal muscle voltage-dependent Ca^{2+} channel to chromosome 7q21–q22 by somatic cell hybrid analysis. *Genomics* **19**: 192–193.
- Quandt, K., K. Frech, H. Karas, E. Wingender, and T. Werner. 1995. MatInd and MatInspector—New fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acids Res.* **23**: 4878–4884.
- Saitou, N. and M. Nei. 1987. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406–425.
- Singer, D., M. Biel, I. Lotan, V. Flockerzi, F. Hofmann, and N. Dascal. 1991. The roles of the subunits in the function of the calcium channel. *Science* **253**: 1553–1557.
- Tanaka, O., H. Sakagami, and H. Kondo. 1995. Localization of mRNAs of voltage-dependent Ca^{2+} -channels: Four subtypes of α_1 and β -subunits in developing and mature rat brain. *Brain Res. Mol. Brain Res.* **30**: 1–16.
- Taviaux, S., M.E. Williams, M.M. Harpold, J. Nargeot, and P. Lory. 1997. Assignment of human genes for β_2 and β_4 subunits of voltage-dependent Ca^{2+} channels to chromosomes 10p12 and 2q22–q23. *Hum. Genet.* **100**: 151–154.
- Thompson, J.D., T.J. Gibson, F. Plewniak, F. Jeanmougin, and D.G. Higgins. 1997. The ClustalX windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **24**: 4876–4882.
- Walker, D. and M. De Waard. 1998. Subunit interaction sites in voltage-dependent Ca^{2+} channels: Role in channel function. *Trends Neurosci.* **21**: 148–154.
- Wei, X.Y., E. Perez-Reyes, A.E. Lacerda, G. Schuster, A.M. Brown, and L. Birnbaumer. 1991. Heterologous regulation of the cardiac Ca^{2+} channel α_1 subunit by skeletal muscle β and α subunits. Implications for the structure of cardiac L-type Ca^{2+} channels. *J. Biol. Chem.* **266**: 21943–21947.
- Williams, M.E., M.S. Washburn, M. Hans, A. Urrutia, P.F. Brust, P. Prodanovich, M.M. Harpold, and K.A. Stauderman. 1999. Structure and functional characterization of a novel human low-voltage activated calcium channel. *J. Neurochem.* **72**: 791–799.
- Yu, A.S. 1995. Identification and localization of calcium channel α_1 and β subunit isoforms in the kidney. *Kidney Int.* **48**: 1097–1101.

Received May 28, 1999; accepted in revised form October 14, 1999.