



Nature and function of insulator protein binding sites in the *Drosophila* genome

Yuri B Schwartz, Daniela Linder-Basso, Peter V Kharchenko, et al.

Genome Res. published online July 5, 2012

Access the most recent version at doi:[10.1101/gr.138156.112](https://doi.org/10.1101/gr.138156.112)

P<P	Published online July 5, 2012 in advance of the print journal.
Accepted Manuscript	Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.
Creative Commons License	This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see http://genome.cshlp.org/site/misc/terms.xhtml). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 3.0 Unported License), as described at http://creativecommons.org/licenses/by-nc/3.0/ .
Email Alerting Service	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here .

Comprehensive immune receptor profiling.
Discover the **DriverMap™ AIR Assay** difference.

LEARN
MORE



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

© 2012, Published by Cold Spring Harbor Laboratory Press

Nature and function of insulator protein binding sites in the *Drosophila* genome

Yuri B. Schwartz^{1,2,§}, Daniela Linder-Basso^{2*}, Peter V. Kharchenko^{3*}, Michael Y. Tolstorukov^{3*}, Maria Kim^{1,4}, Hua-Bing Li², Andrey A Gorchakov^{5,8}, Aki Minoda⁶, Gregory Shanower², Artyom A. Alekseyenko⁵, Nicole C. Riddle⁷, Youngsook L. Jung³, Tingting Gu⁷, Annette Plachetka⁵, Sarah C. R. Elgin⁷, Mitzi I. Kuroda⁵, Peter J. Park³, Mikhail Savitsky^{1,4}, Gary H. Karpen⁶, Vincenzo Pirrotta^{2,§}

#1 Department of Molecular Biology, Umeå University, Umeå, Sweden

#2 Department of Molecular Biology & Biochemistry, Rutgers University, Piscataway, NJ, USA

#3 Center for Biomedical Informatics, Harvard Medical School, Boston, MA, USA

#4 Group of Telomere Biology, Institute of Gene Biology, Moscow, Russia

#5 Division of Genetics, Dept. of Medicine, Brigham & Women's Hospital, and Dept. of Genetics, Harvard Medical School, Boston, MA, USA

#6 Department of Molecular and Cell Biology, University of California at Berkeley and Department of Genome Dynamics, Lawrence Berkeley National Lab, Berkeley, CA, USA

#7 Department of Biology, Washington University in St. Louis, St. Louis, MO, USA

#8 Institute of Molecular and Cellular Biology, Novosibirsk, Russia

running title: Genomics of *Drosophila* insulator proteins

key words: chromatin insulators, Polycomb, genomics, modENCODE

* these authors have contributed equally.

§ corresponding authors: yuri.schwartz@molbiol.umu.se ;

pirrotta@biology.rutgers.edu

ABSTRACT

Chromatin insulator elements and associated proteins have been proposed to partition eukaryotic genomes into sets of independently regulated domains. Here we test this hypothesis by quantitative genome-wide analysis of insulator protein binding to *Drosophila* chromatin. We find distinct combinatorial binding of insulator proteins to different classes of sites and uncover a novel type of insulator element that binds CP190 but not any other known insulator proteins. Functional characterization of different classes of binding sites indicates that only a small fraction act as robust insulators in standard enhancer-blocking assays. We show that insulators restrict the spreading of the H3K27me3 mark but only at a small number of Polycomb target regions and only to prevent repressive histone methylation within adjacent genes that are already transcriptionally inactive. RNAi knockdown of insulator proteins in cultured cells does not lead to major alterations in genome expression. Taken together these observations argue against the concept of a genome partitioned by specialized boundary elements and suggest that insulators are reserved for specific regulation of selected genes.

The data sets reported in this study are available from GEO database under the following accession numbers: GSE32775, GSE32773, GSE32774, GSE20811, GSE20812, GSE20760, GSE32816, GSE32777, GSE32780, GSE32776, GSE32778, GSE20815, GSE20766, GSE20814, GSE32781, GSE32783, GSE32782, GSE20767, GSE32749, GSE20768, GSE32750, GSE20802, GSE23489, GSE32808, GSE32812, GSE32813, GSE32810, GSE20808, GSE20833, GSE20809, GSE32853, GSE25373, GSE32791, GSE32788, GSE32789, GSE32790, GSE32792.

INTRODUCTION

Insulator elements were first discovered in *Drosophila melanogaster* by biochemical (Udvardy et al., 1985) and genetic approaches (Holdridge and Dorsett, 1991; Geyer and Corces, 1992) as specialized chromatin structures that appeared to define boundaries between different chromatin states. It was soon found that such insulator elements have the ability to block enhancer action when interposed between enhancers and promoters, and that this activity depended on specific DNA binding proteins that associate with the insulator element. In *Drosophila*, we now know of four well-defined insulator DNA binding proteins, SU(HW), ZW5 (also known as DWG), BEAF-32 and CTCF (Geyer and Corces, 1992; Zhao et al., 1995; Gaszner et al., 1999; Moon et al. 2005), of which only CTCF has an ortholog in mammals (Baniahmad et al. 1990; Lobanenko et al. 1990). Two other proteins, MOD(MDG4) and CP190, were found to associate with the SU(HW)-binding insulator element found in the *gypsy* transposon and are also required for its insulator function (Georgiev and Gerasimova, 1989; Gerasimova et al. 1995; Pai et al., 2004). Both MOD(MDG4) and CP190 contain a POZ/BTB structural motif, known to mediate homotypic and heterotypic protein-protein interactions, which may drive the association between multiple insulator elements. Subsequent work has shown that CP190 also shares some of its chromatin binding sites with BEAF-32 and CTCF and can interact directly with the latter (Mohan et al., 2007; Gerasimova et al., 2007; Bushey et al., 2009; N gre et al., 2010).

The mechanistic interdependencies between CP190 and the sequence-specific DNA binding proteins remain somewhat controversial. CP190 protein is recruited to *gypsy* insulator by SU(HW), but also binds directly to the endogenous SU(HW)-dependent

insulator DNA from the *y - achaete* locus (Pai et al., 2004). CTCF binding to chromosomes was variously claimed to be either strictly (Gerasimova et al., 2007) or partially (Mohan et al., 2007) dependent on CP190 or, more recently, completely independent of CP190 (Wood et al., 2011). Despite the uncertainties, an influential model proposes that CP190 acts as a universal “glue” protein that mediates interactions between insulator elements of different classes, thereby generating chromatin loops, whose properties are postulated to be such that regulatory elements located on one loop are hindered from interacting with promoters or other elements on the adjacent loop (Gerasimova et al., 2007; Bushey et al., 2009).

Previous immunolocalization and ChIP-chip/ChIP-seq analyses have shown that insulator proteins have numerous binding sites in the *Drosophila* genome (Zhao et al., 1995; Gerasimova and Corces, 1998; Mohan et al., 2007; Bartkuhn et al., 2009; Bushey et al., 2009; N  gre et al., 2010). Together with similar data for the mammalian CTCF protein (Kim et al., 2007; Cuddapah et al., 2009), these findings suggest that the genome is partitioned into domains delimited by boundaries that prevent spreading or influencing the chromatin state of flanking domains. According to this view, insulator elements would be expected to be very abundant in the genome and serve an essential function to protect genes from inappropriate action of enhancers, silencers and other chromatin modifying activities affecting gene function.

Another view of insulator function, not incompatible with the first, derives from the discovery that insulator protein binding sites play a critical role in several complex regulatory regions from mammalian and *Drosophila* genomes, where they bring together different components or juxtapose regulatory elements with the appropriate

promoters (Splinter et al., 2006; Kurukuti et al., 2006; Ling et al., 2006; Li et al., 2011). Although these elements were originally discovered for their ability to separate genomic units, the view that insulators may bring parts of the genome together is consistent with the realization that chromatin looping is an essential feature of genome architecture and gene regulation (Lancôt et al., 2007, Schoenfelder et al., 2010). In this view, however, such “linking” and folding is the basic function of “insulator” elements and, in principle, not every insulator protein binding site necessarily has an enhancer blocking insulator function.

Here we evaluate the two concepts by genome-wide analysis of insulator protein binding to *Drosophila* chromatin. We focus on the quantitative aspects of binding, which reveal classes of binding sites occupied by specific combinations of insulator proteins. We demonstrate that distinct rules govern the binding of an insulator protein to different classes of sites, which sometimes involve cooperation between several insulator proteins. We also describe a novel class of robust *Drosophila* insulator elements that in cultured cells bind CP190 but not any other known insulator proteins. We find that only a small fraction of insulator protein binding sites act as robust enhancer blockers *in vivo* and that significant depletion of insulator proteins in cultured cells has small effects on genome-wide expression or the spreading of the H3K27me3 mark. Our observations argue against the concept of a genome partitioned by specialized boundary elements and suggest, instead, that insulators are reserved for specific regulation of selected genes.

RESULTS

Using chromatin immunoprecipitation analyzed by hybridization to *Drosophila* genomic tiling arrays (ChIP-chip), we have mapped the distributions of SU(HW), CTCF, BEAF-32, ZW5, CP190 and MOD(MDG4)67.2 proteins in cultured S2-DRSC and ML-DmBG3-c2 cells (hereafter referred to as S2 and BG3). As has been reported previously (Bushey et al., 2009; N gre et al., 2010), the genomic distributions of insulator proteins overlap. To characterize the persistent co-binding groups we first used a relaxed threshold of $\log_2(\text{IP}/\text{INPUT}) > 0.7$ to define genomic regions bound by each protein and record all possible types of overlapping combinations. Each region was further examined for the strength of binding of associated proteins and only those regions in which all associated proteins bound with comparative strength were considered for further analysis (Figure 1, Tables S1-S18). The last step is critical to take into account differences in antibody strengths and discriminate between sites with robust co-binding of several proteins and sites at which one protein binds strongly but others are barely detectable. This approach shows clearly that some of the co-binding combinations reported earlier (N gre et al., 2010) are in fact at the edge of computational detection (Figure 1). For example class 14 sites that would appear to co-bind CP190 and SU(HW) in the absence of MOD(MDG4) display exceedingly weak CP190 signals, which is in stark contrast to the robust CP190 binding to class 3 sites in the presence of SU(HW) and MOD(MDG4).

Consistent with a broad role of CP190 in the insulator network, approximately 80% of robust CP190 binding sites are shared with either SU(HW), CTCF or BEAF-32 (Figure 1). In contrast, more than half of the SU(HW), CTCF and BEAF-32 sites are standalone, *i.e.* none of the other proteins tested are bound to these sites. This implies that the interaction of SU(HW), CTCF and BEAF-32 with CP190 and other co-

binding partners depends on additional factors. In addition, 83 robust standalone CP190 sites indicate that this protein can be recruited to chromatin independently of SU(HW), CTCF and BEAF-32.

As expected from polytene chromosome staining (Gerasimova and Corces, 1998) we detect ~ 300 sites with simultaneous robust binding of SU(HW), MOD(MDG4)67.2 and CP190, the combination of proteins associated with the *gypsy* insulator (Figure 1, class 3). We will refer to this class of binding sites as *gypsy*-like although none of them corresponds to *gypsy* retrotransposon insertions as all repetitive sequences were excluded from our analysis. We see no MOD(MDG4)67.2 binding in the absence of SU(HW) and CP190.

Cis-cooperation and motif coincidence govern the co-binding of CP190 with CTCF and SU(HW).

Since SU(HW) and CTCF interact directly with CP190 (Pai et al., 2004, Gerasimova et al., 2007), the large number of standalone sites for these two proteins requires explanation. The comparison of the genomic distributions of SU(HW), CTCF and CP190 in S2 and BG3 cells and in whole embryos (from Négre et al. 2010) shows that the distinction between the standalone and CP190 co-bound sites is well-preserved in all three sources of chromatin. We conclude that the co-binding of CP190 is an inherent property of a site rather than a product of tissue specific regulation.

The analysis of DNA sequences in standalone SU(HW) and CTCF regions (Class 2, 4) and those shared with CP190 (Class 3, 9) indicates that SU(HW) and CTCF bind to DNA directly and with the same sequence specificity irrespective of CP190 presence.

The most prominent motifs derived from the corresponding standalone and CP190 co-bound regions are essentially identical (Figure 2A, Figure S1) and match the reported binding sequences of SU(HW) and CTCF *in vivo* (Adryan et al., 2007; Holohan et al., 2007; N gre et al., 2010) and in *in vitro* (Spana and Corces, 1990; Golovnin et al., 2003; Moon et al., 2005). The number of SU(HW) or CTCF motifs in the standalone sites does not differ significantly from that in the CP190 co-binding sites.

In addition to the canonical CTCF motif, the sequence analysis reveals a new motif enriched in class 9 (CTCF+CP190) but not class 4 (standalone CTCF) regions (Figure 2A, Figure S1). Strikingly, this motif is also enriched at class 6 (standalone CP190) sites (Figure 2A, Figure S1), suggesting that CP190 can bind to DNA directly or through an unknown DNA-binding protein(s) (see Supplementary Results for more details) and that the binding of CTCF and CP190 to common sites results from the coincidence of the corresponding recognition sequences. It is clear, however, that at many sites CTCF is required for CP190 binding. RNAi depletion of CTCF that reduces its binding at class 9 (CTCF+CP190) sites (Figure 2B) also reduces CP190 binding at most of those same sites but not at class 3 (*gypsy*-like) or other sites where CP190 is not accompanied by CTCF (Figure S2). The converse knock-down of CP190 has very little effect on CTCF binding (Figure 2B, S2), indicating that CTCF is recruited to class 9 (CTCF+CP190) sites independently of CP190.

In contrast to class 9 (CTCF+CP190) sites, the sequence analysis of class 3 (*gypsy*-like) and class 2 (standalone SU(HW)) sites revealed no characteristic motifs other than the SU(HW) recognition sequence. The RNAi knock-down of SU(HW) results in its efficient depletion from chromosomes and also in depletion of CP190 from the

majority of *gypsy*-like sites, but not from other kinds of CP190 sites (Figure 2B, S2). Unexpectedly, CP190 depletion also causes loss of both proteins from class 3 (*gypsy*-like) sites (Figure 2B), indicating that the binding of SU(HW) and CP190 to these sites is mutually dependent. CP190 RNAi has little effect on the binding of SU(HW) to standalone sites (Figure S2), arguing that the dependence is direct. Although other explanations are possible, these results suggest that SU(HW) and CP190 proteins cooperate, allowing SU(HW) to bind to a recognition sequence of a quality or in an environment inadequate to recruit on its own. Supporting this notion, we find that the consensus scores of the SU(HW) motifs at class 3 (*gypsy*-like) sites are markedly lower than those at class 2 (standalone SU(HW)) sites (Figure S3).

Finally, although more than half of SU(HW) or CTCF sites are standalone, SU(HW) and CTCF never bind a common region unless together with CP190 and MOD(MDG4)67.2 (Figure 1; class 12 sites). The apparent co-binding of CTCF and SU(HW) to class 12 sites might be attributed to the crosslinking of complexes recruited to distinct insulator elements and bridged in *trans* by interactions between CP190 and MOD(MDG4)67.2 proteins. This model predicts that the DNA sequences recognized by CTCF and SU(HW) would rarely group together at class 12 sites. Contrary to this prediction, class 12 sites show a high coincidence of SU(HW) and CTCF motifs, a feature absent from sites that bind only SU(HW) or only CTCF (Figure 2D). This points to the DNA sequence as the primary determinant of the common binding to class 12 sites, and argues against their being the product of crosslinking of distinct trans-interacting regions (although such trans-interactions are not excluded).

BEAF-32 is dispensable for the recruitment of CP190 to chromatin

BEAF-32 was suggested to act as a DNA binding recruiter of CP190 (Bushey et al., 2009). Indeed the comparison of BEAF-32 and CP190 regions defined at low threshold ($\log_2(\text{IP}/\text{INPUT}) > 0.7$) shows extensive overlap (Figure 1). It is immediately obvious, however, that the binding of CP190 to these sites is often weak and disproportional to BEAF-32 (hence the relatively small number of regions listed as robustly bound by both proteins, classes 5 and 8). RNAi depletion of BEAF-32 causes a reduction of its binding to the majority of the sites shared with CP190 (class 5 sites, Figure 2C). However, it has no effect on the binding of CP190 to these sites. Conversely CP190 depletion reduces its binding to the majority of class 5 sites but does not affect BEAF-32 binding (Figure 2C). We conclude that BEAF-32 and CP190 bind chromatin independently of each other, and that their coincidence may result from a bias of both proteins towards active Transcription Start Sites (TSSs).

RNAi-knockdown discriminates between low- and high- affinity binding sites

The loss of a chromatin protein from its genomic binding sites upon RNAi knock-down is sometimes used to validate the genome-wide mapping. The ten-fold reduction of nuclear protein levels in the above RNAi experiments (Figure 3A, S4) results in the complete loss of binding at many chromosomal sites and reduction of binding to the majority of them (Figures 3B-E). Yet in all cases we see a number of strong sites that remain unaffected by RNAi. Immunoprecipitations using two antibodies independently raised against different parts of the proteins strongly suggest that these are genuine high affinity binding sites, and not false positives. For example, one of the sites with persistent BEAF-32 binding is *scs'*, the prototype BEAF-32 dependent insulator (Figure 3F). We conclude that immunoprecipitation with two independently

derived antibodies is a better validation criterion than the use of RNAi depletion, which in our case would reject sites with the highest affinity and therefore with the best functional potential.

Analysis of insulator function

A comparison of the genomic distributions of different classes of binding sites to genes and gene activity shows very clear differences. The majority of class 3 (*gypsy*-like), standalone SU(HW) (class 2) and CTCF (class 4) sites and about a half of class 9 (CTCF+CP190) sites are situated within introns of transcriptionally inactive genes or in intergenic regions (Figure 4A, B). In contrast, the other half of class 9 (CTCF+CP190), as well as ZW5, BEAF-32 and standalone CP190 sites tend to reside within 2kb of transcriptionally active TSSs (Figure 4A, B). None of the classes of binding sites have significant preference for positions situated between an active and a silent gene.

The distinct genomic location of different classes of binding sites raises the question of whether they have the same functions or insulating properties. To examine insulator function, we selected two representative 1kb DNA fragments from each major class and measured their ability to block the activation of the *yellow* reporter gene by the upstream wing- and body-specific enhancers when placed between these enhancers and the promoter. Unlike general repressors, insulators are expected to block the upstream enhancers without affecting the activation of the *yellow* promoter by the downstream bristle specific enhancer (Figure 4C, S5) (Geyer and Corces, 1992). Five randomly chosen 1kb genomic fragments that showed no association with any of the insulator proteins in our ChIP-chip experiments and the 680bp *gypsy*

insulator element were tested in the same reporter assay as negative and positive controls. Initially all reporter constructs were integrated in the same 51D landing site by targeted ϕ C31 *att* recombination (Bischof et al., 2007), which allowed direct comparison of the effects produced by different test fragments in the same chromosomal environment. Subsequently, the transgenes were mobilized from the 51D site using P-element mediated transposition to assess the robustness of the enhancer blocking effect in different chromosomal contexts.

As summarized in Figure 4D, and Tables 1, S19, S20, the transformation of flies lacking *yellow* function with negative control constructs restores their phenotype to nearly wild type. The pigmentation of the body and wings in these flies varies somewhat depending on the site of insertion, but is always much stronger than in flies transformed by the positive control construct that carries the *gypsy* insulator, which have black bristles but completely yellow wings and very light body cuticle. Of the 16 insulator protein binding sites tested, only two (BC1, class 5; CP1901, class 6) block the upstream enhancers to the same extent as the *gypsy* insulator construct. The enhancer block is robust, evident at all tested chromosomal locations and completely reversed upon FLP-mediated excision of tested fragments. In addition, we find four fragments (CTCFC1, class 9; B1, class 7; CP1902, class 6 and BC2, class 5) whose enhancer blocking ability is less strong than that of BC1 (class 5) and CP1901 (class 6) and varies depending on the surrounding chromatin context. Fragment BC2 (class 5) represents the most striking example of variability, with extremely good enhancer blocking at some insertion sites and complete lack of it at others.

None of the class 3 (*gypsy*-like) sites tested displayed enhancer blocking activity, in agreement with the results of N gre et al., (2011) who tested several other fragments of this class using a different enhancer-blocking assay based on the *eve* stripe 2 and 3 enhancers. On the other hand the class 3 (*gypsy*-like) binding sites from the *yellow - achaete* and 62D regions have been shown to robustly block *yellow* enhancers in transgenic tests (Golovnin et al., 2003; Parnell et al., 2003; Kuhn-Parnell et al., 2008). This suggests that the simple recruitment of CP190, MOD(MDG4)67.2 and SU(HW) to a chromosomal site is not sufficient for robust enhancer blocking, and that additional unknown factors or specific chromatin configurations are required for *gypsy*-like binding sites to have this function.

Overall we conclude 1) that, unlike the prototype insulators, the majority of insulator protein binding sites are not robust enhancer blockers, 2) that the complement of binding proteins at each class of sites is a poor predictor of whether a site can act as an enhancer blocker and 3) that a given site can act as enhancer blocker in one genomic context but not in another. The latter implies the possibility that a site that does not appear to act as enhancer blocker might become such if the chromatin environment changes. Some functional regulatory elements can be pinpointed based on their high DNA sequence conservation. This appears not to be the case for insulator protein binding sites. Thus the sequences of BC1 (class 5) and CP1901 (class 6) fragments, which show the best enhancer blocking in the transgenic test, and the sequences of the sites from these classes in general, show surprisingly low conservation (Figures S6-S8 and Supplementary results).

Unaddressed by our functional test is the question whether the sites occupied by a combination of DNA binding insulator proteins have properties markedly different from their simpler counterparts. Future experiments should reveal, for example, whether class 12 sites have poor enhancer-blocking ability like class 3 (*gypsy*-like) sites or can block enhancer-promoter communications as well or better than class 9 (CTCF+CP190) sites.

Finally we note that both fragments tested to represent class 6 (standalone CP190 sites) and class 5 (BEAF-32+CP190) sites display a degree of enhancer blocking and include the only robust enhancer blockers found in our tests (Table 1). Considering the fact that the CP190 binds to class 5 (BEAF-32+CP190) sites independently of BEAF-32, this underscores the importance of the novel pathway of CP190 recruitment to chromatin and suggests that it may be utilized at the majority of robust enhancer blocking elements in *Drosophila*.

Standalone SU(HW) binding sites act as general repressors

In these assays we found no evidence of enhancer blocking by class 2 and 4 (standalone CTCF or SU(HW)) binding sites, although we cannot exclude the possibility that some may be active in specific tissues where they acquire CP190, as recently proposed by Wood et al., (2011). Instead, at many chromosomal locations the two representative class 2 (standalone SU(HW)) binding fragments S1 and S2 cause loss of *yellow* expression not only in wings and body but also in bristles, indicative of general promoter repression rather than enhancer blocking activity (Tables S19, S20). Such behavior is reminiscent of the repressive properties of the *gypsy* insulator upon loss of *mod(mdg4)* function (Gerasimova et al., 1995), and suggests that

transcriptional repression is a general feature of SU(HW) protein when not associated with MOD(MDG4)67.2.

The impact of insulator proteins on gene expression

It is surprising that most of the insulator protein binding sites tested appear to lack robust enhancer blocking activity, raising the possibility that the transgenic assay may underestimate the fraction of functional binding sites because they require their native genomic context. Therefore, as an independent measure of an insulator protein impact on the genome, we evaluated genomic changes in gene expression after depletion of SU(HW), CTCF, BEAF-32 or CP190 in BG3 cells. Consistent with the notion that only a small fraction of insulator protein binding sites corresponds to functional insulators, significant depletion of any single insulator protein does not lead to major alterations in gene expression (Figure 5A). We see no wide-spread switching on of inactive genes by the adjacent “active” chromatin environment or repression of active genes by encroaching “repressive” chromatin states

The few changes in gene expression that we can detect are consistent with the results of the transgene tests (Figure 5A). From 39 genes affected by SU(HW) RNAi the expression of 24 genes is upregulated. The magnitude of the expression increase at these genes is much higher than the reduction at genes where the expression goes down, which fits well with the repressive properties of standalone SU(HW) sites seen in the transgenic assay. The transgenic assay also suggests that CP190 bound sites, especially standalone, tend to block enhancer-promoter communications. If they function as pure insulator elements one would expect that in some cases their loss might cause inappropriate activation of a gene, while in other cases it might lead to inappropriate repression. We find instead that the genes affected by CP190 knock-

down tend to reduce their expression. It is possible that this apparent stimulatory role of CP190 stems from a consistent bias for using CP190-dependent insulators to block long-range transcriptional repression. However, we favor an alternative explanation that the most frequent role of CP190 complexes is to aid chromatin folding to bring distant activators to their appropriate targets.

Insulators and Polycomb silencing

Broad domains of histone H3 trimethylated at K27 (H3K27me3) mark loci repressed by Polycomb group (PcG) proteins (Schwartz et al., 2006). In *Drosophila*, these proteins are recruited to the target genes by Polycomb Response Elements (PREs), from which the H3K27me3 mark spreads by a chromatin looping mechanism (Kahn et al., 2006; Comet et al., 2011). The *gypsy* insulator can interfere with the looping of PRE-bound complexes and block the spreading of H3K27me3 (Kahn et al., 2006, Comet et al., 2011). Visual inspection shows that half of the H3K27me3 domain edges (110 of 221) display a gradual decline to background level (exemplified by the edge of the *sens-2* domain in Figure 5B). The remaining 111 edges are sharp enough to define a distinct domain boundary (Figure 5C), which we will refer to as H3K27me3 domain ‘borders’. Two genomic features correlate with the presence of domain borders (Figure 5D); 54% of the domain borders coincide with robust insulator protein binding sites, and 78% of the borders coincide with the 5’ or 3’ ends of active transcripts. At least one of the two features is present at 97% (108 of 111) of definable borders, suggesting that both may contribute to limiting the spread of H3K27 tri-methylation.

44% of the borders are marked only by the presence of adjacent transcriptional activity, and 33% of the borders coincide with both the ends of active transcription and insulator protein binding sites (Figure 5D), consistent with the tendency of BEAF-32 and CP190 proteins to bind in the 5' region of active genes. Transcriptional activity itself may be sufficient to prevent the spread of H3K27me3 by the associated histone H3 replacement, H3K27 acetylation or inhibition of the histone methyltransferase activity of PcG complexes by H3K4me3 (Schmitges et al., 2011). This appears to be the case as only three H3K27me3 domains bordered by active transcripts are extended after RNAi knockdown of insulator proteins (3% of all domain borders associated with transcriptional activity). We conclude that, if putative insulator elements contribute to the establishment of the borders adjacent to active loci, they are dispensable for their maintenance in most cases.

In contrast, at 75% (18 out of 24) of domain borders that contain insulator protein binding sites but no active genes, RNAi knock-down of the corresponding insulator proteins results in expansion of the H3K27me3 domains. The affected boundaries coincide with class 12, class 3 (*gypsy*-like), class 9 (CTCF+CP190) and class 6 (standalone CP190) binding sites, consistent with the idea that such sites can act as functional insulators.

In ~5% (5 out of 110) of cases the knockdown of insulator proteins leads to the extension or changes in the shape of the gradually declining H3K27me3 domains. In these cases, exemplified by the left tail of the *sens-2* domain (Figure 5B), the affected domains contain class 3 (*gypsy*-like) and/or class 9 (CTCF+CP190) sites within their tails and change their length and/or shape upon RNAi knockdown of CP190, SU(HW)

or CTCF. We interpret this to indicate that the block imposed by class 3 (*gypsy*-like) or class 9 (CTCF+CP190) insulator elements is not always robust: some H3K27me3 may bleed through the insulator but, in its absence, spreading of the histone methylation is more efficient and longer-range. Finally, we note that standalone CTCF sites are never found at H3K27me3 domain borders, supporting the idea that these are not insulators and cannot block the spreading of histone methylation.

Overall, we conclude that insulators participate in shaping the genomic distribution of H3K27me3. However, in cultured cells their contribution is small and used primarily to prevent the extensive H3K27 tri-methylation of transcriptionally inactive genes adjacent to PcG target regions. This role may be most important when Polycomb repression is first established in the embryo.

DISCUSSION

The binding sites of insulator proteins are often taken to represent elements that partition the genome into independent regulatory domains and demarcate chromosomes into regions of “active” and “repressed” chromatin. The results presented here give little support to this view as a general principle of genome organization although it may be true in certain regions. Instead we would like to argue that: 1) insulator proteins bind to genomic sites in specific combinatorial patterns, 2) the properties of sites bound by key insulator proteins SU(HW) and CTCF are markedly different depending on whether the two co-bind with CP190, 3) many of the known insulator proteins sites do not function as robust enhancer blockers, and 4) at least in cultured cells the depletion of insulator proteins has a limited impact on genome-wide gene expression.

Combinatorial binding patterns

Classifications of combinatorial binding of insulator proteins have been described previously (Bushey et al., 2009; N gre et al., 2010). These classifications relied on the overlapping of bound regions defined according to arbitrary statistical thresholds and the position of these regions relative to TSSs. Because they did not take into account the relative strengths of binding, such classifications grouped together binding sites with very different biochemical and functional properties.

In contrast, we define the persistent co-binding patterns based on the strength of binding of the associated proteins, treating regions strongly bound by a combination of proteins differently from regions at which the same proteins are detected according to a statistical threshold but where the extent of their binding is disproportional. We argue that our approach retains the information on biochemical interrelations between the co-bound proteins and separates the sites with different functional properties. The strongest support for our argument comes from RNAi knock-down experiments which demonstrate that the effect of the loss of one insulator protein on the binding of another insulator protein is constrained to a specific class of co-bound regions. For example, the knock-down of SU(HW) results in the loss of CP190 from class 3 (*gypsy*-like) sites but not from class 9 (CTCF+CP190) or class 5 (BEAF-32+CP190) sites.

Our approach to select the sites representative of each co-binding class is conservative and inevitably excluded a fraction of binding sites from downstream analyses. For example, strong SU(HW) binding sites assigned to class 14 by initial overlap comparison (Figure 1) were not analyzed further due the uncertainty of their co-

binding by CP190. We therefore caution readers that our selection of representative binding sites (Tables S1-S18) is not a complete genomic catalogue and advise to use the ChIP-chip binding profiles, deposited to GEO and modMINE, to gauge whether their locus of interest has a strong insulator protein binding site.

The role of CP190 in insulation

The prevailing model in the field suggests that CP190 is recruited to different insulator elements by DNA binding proteins where it serves as a universal adapter that mediates interactions between different insulator elements (Bushey et al., 2009). Our results present a more complex picture. First, RNAi knock-down experiments demonstrate that the binding of SU(HW) protein to class 3 (*gypsy*-like) sites is dependent on CP190, indicating that CP190 is not passively tethered to common sites by SU(HW) and instead plays an active role in recruitment and/or stabilization of the bound complex. Second, the sequence analysis of class 9 (CTCF+CP190) sites suggests that the binding of both proteins to these sites is likely due to the coincidence of cognate recognition sequences. Third, RNAi knock-down experiments indicate that BEAF-32 is dispensable for CP190 binding at shared sites. Clearly CP190 plays an active role in the selection of sites shared with SU(HW), CTCF or BEAF-32. It is still possible that once it co-binds, or binds sufficiently close to another insulator protein, it may mediate the trans-interactions of the bound sites. However, such interactions would have to be rather transient, at least in cultured cells, as they are not easily detected in our ChIP-chip data.

The class of sites in which CP190 is not accompanied by any of the insulator proteins tested indicates the existence of a novel pathway of CP190 recruitment to chromatin.

Notably, our functional tests suggest that the sites employing this pathway (i.e. class 6 (standalone CP190) and class 5 (CP190+BEAF-32) sites may constitute the major pool of robust insulator elements in flies. This conclusion is supported by functional analyses of N gre et al., (2011) who found enhancer blocking activity by three DNA fragments that we would classify as class 6 (standalone CP190) sites and one fragment that we classify as class 5 (CP190+BEAF-32) site. Interestingly N gre et al (2011) found some degree of CTCF binding at these sites in embryos. Our results show that these sites bind no CTCF in S2 or BG3 cultured cells. Furthermore, unlike most CTCF binding sites (Figure 2D), these regions contain no CTCF recognition motif. Whether and how such sites can actually recruit CTCF in embryonic cells but not in cultured cells will require further investigation.

Is transcriptional repression the primary function of SU(HW) protein?

SU(HW) is not required for *Drosophila* viability, but mutant flies display defective oogenesis and female sterility (Parkhurst et al., 1988). As follows from the experiments presented here and previously (Golovnin et al., 2003; Parnell et al., 2003; Kuhn-Parnell et al., 2008), the class 3 (*gypsy*-like) sites do not have a direct impact on gene promoters and some can act as enhancer blockers. In contrast, our results show that standalone SU(HW) protein binding sites tend to repress transcription rather than insulate. Remarkably, a recent study indicates that neither CP190 nor MOD(MDG4)67.2 are required for oogenesis (Baxley et al., 2011), which suggests that the role of SU(HW) in the control of this process is distinct from its enhancer-blocking function. We hypothesize that the SU(HW) function critical for oogenesis is transcriptional repression exerted at standalone binding sites, consistent with the

upregulation of gene expression we observe after depletion of SU(HW) in cultured cells.

Insulator proteins and chromatin states

Previously we have shown that the fly genome can be partitioned based on nine combinatorial patterns of 18 histone modifications (Kharchenko et al., 2011). Contrary to initial expectations, we find little correlation between the positions of insulator protein binding sites and the boundaries of these combinatorial chromatin states (data not shown). In agreement with this result the transgenic tests suggest that only a small fraction of insulator protein binding sites can robustly block enhancer-promoter communication and we see no major changes in gene expression after RNAi knock-down of insulator proteins. Taken together these observations suggest that insulator proteins are unlikely to play a general role in partitioning of the fly genome into distinct domains of different chromatin states.

We realize that the incomplete loss of insulator proteins after RNAi knock-downs may cause an underestimate of the potential changes in gene expression and the extent of H3K27me3 domains. We believe the underestimated changes are likely to be few and their accounting would not influence our overall conclusions. Firstly, we note that even partial loss of insulator proteins from a site is sufficient to impair its ability to block the spreading of H3K27 methylation (Figure S9). Conversely, if all H3K27me3 domain borders at which the lack of expansion can be explained by the lack of significant reduction of insulator protein binding after corresponding RNAi are disregarded, the fraction of affected borders remains essentially the same. Secondly, it is clear that our statistical definition of the consistent reduction of binding to a site is

conserved. As illustrated by Figures 3B-E, most of the sites, even those not deemed to reduce the binding significantly (green dots on the scatter-plots), bind less insulator proteins after the corresponding RNAi. The sites that remain truly unaffected (data points on or above scatter-plot diagonals) are very few: 81 - SU(HW) binding sites; 27 – CTCF binding sites; 1 – CP190 binding site; 7 – BEAF-32 binding sites. These numbers are at least two orders of magnitude lower than the number of chromatin state partitions (Kharchenko et al., 2011) or active genes in BG3 cells (Cherbas et al., 2011). The disparity between the actual binding reduction and its statistical significance is greatest in the case of BEAF-32 and CTCF RNAi (Figures 3C, 3E) and is explained by the higher variability between the corresponding replicate experiments. Since our test to detect significant expression changes relies on replicate comparison it may have underestimated the number of changes in these two cases. In an attempt to account for this we have relaxed the detection criteria and looked for all measurable two-fold changes irrespective of their statistical significance. As illustrated in Figure S10 the numbers of expression changes increase but remain small (BEAF-32 RNAi – 16 up/36 down; CTCF RNAi – 28 up/34 down).

Perhaps not so important in the global scale, insulators may still be critical to restrict chromatin states at a limited set of sites. In fact, the ability of the *gypsy* insulator to shield reporter genes from Polycomb repression is well documented (Sigrist and Pirrotta, 1997; Kahn et al., 2006; Comet et al., 2011) and the extension of endogenous H3K27me3 domains in CTCF and CP190 mutants has been reported (Bartkuhn et al., 2009). The results of genome-wide assays presented here confirm that insulators restrict the spreading of H3K27me3, but only at a small number of Polycomb target regions, and only to prevent the repressive histone methylation of adjacent genes that

are already transcriptionally inactive. While this has no obvious consequences in cultured cells, it may be important in the context of the developing embryo to ensure that genes in the vicinity of Polycomb targets do not become permanently repressed.

METHODS

Cell culture conditions and RNAi

Cell lines were obtained from the Drosophila Genome Resource Center (DGRC) S2-DRSC cells (stock #181) and ML-DmBG3-c2 cells (DGRC, stock #68) and grown according to recommendations. The RNAi was performed as described by Schwartz et al. (2010). The sequences of PCR primers used to produce DNA template for dsRNA synthesis are indicated in the Table S21.

Genome-wide mapping

The mapping of each protein was initially done in the chromatin of S2 cells using two different independently raised antibodies when available (see Table S22, Figure S11 and Supplementary text for technical details). Because of the high congruence between the two independent antibodies (Figure S12) just one was used to map the corresponding proteins in BG3 cells. Chromatin preparation, immunoprecipitation, microarray hybridization and sequencing were done as described in (Kharchenko et al., 2011).

Additional details of experimental procedures and data analyses are indicated in Supplementary Methods.

DATA ACCESS

All data sets reported in this study are available from GEO (accessions: GSE32775, GSE32773, GSE32774, GSE20811, GSE20812, GSE20760, GSE32816, GSE32777,

GSE32780, GSE32776, GSE32778, GSE20815, GSE20766, GSE20814, GSE32781, GSE32783, GSE32782, GSE20767, GSE32749, GSE20768, GSE32750, GSE20802, GSE23489, GSE32808, GSE32812, GSE32813, GSE32810, GSE20808, GSE20833, GSE20809, GSE32853, GSE25373, GSE32791, GSE32788, GSE32789, GSE32790, GSE32792) and modMINE <http://intermine.modencode.org/> (Table S23).

ACKNOWLEDGEMENTS

We thank D. Acevedo, S. Gadel, C. Kennedy and Rutgers BRTC for excellent technical assistance. We are grateful to R. Renkawitz, V. Corces, F. Karch, R. Dorn and S. Small for generous gifts of antibodies, fly stocks and plasmids. This work was supported by NIH modENCODE grant U01HG004258 (GHK, MIK, PJP, SCRE, VP), Wenner-Gren Foundations (YS) and the Russian Academy of Sciences program for Molecular and Cellular Biology (MS).

FIGURE LEGENDS

Figure 1. The classes of insulator protein binding sites. The composition of sixteen co-binding groups detected by initial overlap comparison is indicated by the checkerboard pattern under the bar plot. The color code in $\log_2(\text{IP}/\text{INPUT})$ units (indicated to the right) is used to show the number of sites of different binding strength within each class. For the multi-protein classes the bars are divided from left to right corresponding to the top to bottom positions of the proteins in the chart below. The numbers of sites of each class that bind all corresponding proteins within 60% of their ChIP-chip signal dynamic range are indicated above the bars. Only those sites were used for further analysis. The dashed line on each bar indicates the position of the 60% cutoff. The classes representing robust co-binding combinations are numbered in red.

Figure 2. The sequence determinants and interdependence of the insulator protein binding to chromatin. **A.** The logo representations of sequence motifs characteristic of SU(HW), CTCF and CP190 binding sites defined by MEME algorithm and used in the analysis in (D). **B.** The effects of the RNAi knock-down on the target protein and its co-binding partners. The sites at which ChIP-chip signal was consistently reduced judged from the comparison of two replicate mock RNAi experiments and two specific RNAi experiments ($z\text{-scores} < -3$, unpaired t-test) were counted and their fractions plotted. Here and in panels C-D the error bars indicate the 95% confidence interval. The bar plots show that the binding of CP190 to some of the class 9 but not at *gypsy*-like sites depends on CTCF. However, the binding of CTCF to class 9 sites does not depend on CP190. In contrast, the binding of SU(HW) and CP190 to *gypsy*-like sites is interdependent. **C.** As illustrated by this bar plot BEAF-

32 and CP190 bind to common sites independently. **D.** The presence of SU(HW) and CTCF recognition sequences within indicated classes of sites demonstrates that the coincidence of the two motifs is responsible for the co-binding of SU(HW) and CTCF to Class 12 sites.

Figure 3. The effects of RNAi knock-down on the binding of insulator proteins to chromatin. BG3 cells were subjected to RNAi against key insulator proteins followed by ChIP-chip. **A.** Western blots of three-fold serial dilutions of nuclear protein from cells treated with specific and mock dsRNA (indicated above the panels) show 10-fold or greater knock-down of the corresponding proteins. The antibodies used for detection are indicated to the right and the loading controls are shown in Figure S4. The comparison of average binding for **B** – SU(HW), **C** – CTCF, **D** – CP190, **E** – BEAF-32 after mock and specific RNAi shows that the binding is reduced at the majority of sites (data points below red dashed line). The sites with consistent reduction in both replicate experiments (estimated conservatively with unpaired t-test; z-scores < -3) are shown as blue dots, others are shown as green dots. **F.** *scs'* is one of the BEAF-32 high-affinity binding sites resistant to RNAi. The BEAF-32 ChIP-chip signals after BEAF-32 and mock RNAi are plotted along the segment of Chromosome 3R. White circles mark the peaks affected by BEAF-32 knock-down, red circles mark the peaks that remain unchanged. The genes shown above the coordinate scale are transcribed from left to right, those below the scale from right to left.

Figure 4. Functional evaluation of classes of insulator protein binding sites. A-B. Sites bound by different combinations of insulator proteins show distinct biases in their distribution relative to genes and gene activity. Class 2-4 sites are rarely close to

TSS while class 5-7 sites are primarily TSS-proximal. In rare cases when class 2-4 sites are TSS-proximal these promoters tend to be inactive. In contrast, BEAF-32 (classes 5 and 7) binds predominantly next to active TSS. While many standalone CP190 sites are next to active TSS some are not. The proximity to TSS and genes in (A) is defined based on 2kb margin and the binding to TSS in (B) is defined on 1kb margin. The background distribution expected by chance is shown as the rightmost bar in (A) and is derived from 10 times the number of positions sampled randomly but with the same chromosome representation. C. The schematic of the transgenic enhancer blocking assay. A DNA fragment of interest (black rectangle) is cloned in the FRT cassette positioned between the upstream wing and body enhancers (green ovals) and the promoter of reporter *yellow* gene (yellow rectangle). The resulting construct is injected into *yellow* minus flies. DNA fragments capable of enhancer blocking (red rectangle) prevent the activation of reporter *yellow* gene by upstream enhancers but allow the activation of the gene by downstream bristle enhancer (“br” green oval). This yields transgenic flies with pigmented bristles but yellow body and wings. Ineffectual DNA fragments (green rectangle) allow activation of the reporter gene in all tissues and yield wild type transgenic flies. The fragments harboring repressive activity (blue rectangle) block the expression of transgenic *yellow* in all tissues which results in flies devoid of any pigmentation. The results of transgenic tests are summarized in D.

Figure 5. Functional effects of insulator protein withdrawal. A. Affymetrix GeneChip expression analysis of cells from the RNAi experiments described in Fig. 3. The average fold change between the two specific and two mock RNAi experiments (y-axes) was plotted against the highest average expression value detected in the

mock or specific RNAi samples (x-axes). Each graph point represents one transcript interrogated by the microarray. Transcripts robustly expressed before or after specific RNAi treatment are to the right of vertical dashed lines. Of these, those showing consistent 2-fold or greater change after specific RNAi treatment in both replicate experiments are circled. **B.** As evident from ChIP-chip of H3K27me3 from mock RNAi treated BG3 cells, the *sens-2* gene is repressed by PcG. The right border of the corresponding H3K27me3 domain is sharp and coincides with a standalone CP190 site (marked by vertical green dashed line) and with the *Rca1* transcript. The Chip-chip with H3K4me3 and H3K36me3 indicates that *Rca1* is transcriptionally active. The left side of H3K27me3 domain declines gradually with no obvious border. It harbors *gypsy*-like and CTCF+CP190 binding sites marked by orange and purple dashed lines respectively. The knock-downs of insulator proteins have no effect on the position of the right border of H3K27me3 domain but change the shape of its left tail. The changes in histone methylation profile are best seen on the relative difference browser tracks. **C.** *twi* is also repressed by PcG mechanisms in BG3 cells. The right border of the corresponding H3K27me3 domain is set by the presence of an active transcript. The left border is maintained by a *gypsy*-like (class 3) insulator (vertical orange dashed line) as evident from the extension of K27 tri-methylation after SU(HW) or CP190 knock-down. **D.** The pie chart shows the frequencies of various genomic features associated with definable H3K27me3 domain borders.

TABLES

Table 1. The results of enhancer blocking assay

Fragment class	Construct name	Pigmentation at 51D site*	Pigmentation after fragment excision	Mean after mobilization
Negative control	random1	4/5/5		4.7/4.7/5.0
Negative control	random2	5/4/5		4.7/3.8/5.0
Negative control	random3	4/3/5		4.0/3.1/5.0
Negative control	random4	4/3/5		4.0/3.2/5.0
Negative control	random5	5/3/5		4.6/3.4/5.0
2§	S1	4/2/5	5/4/5	3.4 ^Δ /2.8 ^Δ /3.2 ^Δ
2§	S2	4/2/5	5/4/5	3.9/3.0 ^Δ /3.4 ^Δ
3	SCM2	3/2/5		3.4 ^Δ /3.1/5.0
3	SCM3	5/4/5		4.1/3.6/5.0
4	CTCF1	4/3/5		4.3/3.6/5.0
4	CTCF2	4/3/5		4.3/3.6/5.0
9++	CTCFC1	3/3/5		3.4 ^Δ /2.9 ^Δ /5.0
9	CTCFC2	4/3/5		4.1/4.1/5.0
7++	B1	4/3/5		3.3 ^Δ /3.0 ^Δ /5.0
7	B2	3/3/5		3.6 ^Δ /3.3/5.0
5+++	BC1	1/1/5	5/4/5	2.0 ^Δ /1.8 ^Δ /5.0
5+	BC2	3/3/5		3.1 ^Δ /3.0/5.0
1	ZW3	4/3/5		4.2/3.0 ^Δ /5.0
1	ZW4	5/4/5		4.4/3.9/5.0
6+++	CP1901	2/2/5	5/4/5	1.9 ^Δ /1.4 ^Δ /5.0
6++	CP1902	3/3/5		3.6 ^Δ /2.9 ^Δ /5.0
Positive control	SuHw680 SmaI- ClaI	1/2/5	5/4/5	

* Wing, body, bristles scores are shown: 1 - no pigmentation; 5 – wild type pigmentation. § – context dependent repression; + rear context dependent insulation; ++ context dependent insulation; +++ robust insulation. The mean scores after mobilization marked with ^Δ are significantly different from control (p-value <0.05 in both unpaired t-test and Wilcoxon sum rank test; see Table S19 for scores at each insertion site)

REFERENCES

- Adryan B, Woerfel G, Birch-Machin I, Gao S, Quick M, Meadows L, Russell S, White R. 2007. Genomic mapping of Suppressor of Hairy-wing binding sites in *Drosophila*. *Genome Biol* **88**: R167.
- Baniahmad A, Steiner C, Köhne AC, Renkawitz R. 1990. Modular structure of a chicken lysozyme silencer: involvement of an unusual thyroid hormone receptor binding site. *Cell* **61**: 505-514.
- Bartkuhn M, Straub T, Herold M, Herrmann M, Rathke C, Saumweber H, Gilfillan GD, Becker PB, Renkawitz R. 2009. Active promoters and insulators are marked by the centrosomal protein 190. *EMBO J* **28**: 877-888.
- Baxley RM, Soshnev AA, Koryakov DE, Zhimulev IF, Geyer PK. 2011. The role of the Suppressor of Hairy-wing insulator protein in *Drosophila* oogenesis. *Dev Biol* **356**: 398-410.
- Bischof J, Maeda RK, Hediger M, Karch F, Basler K. 2007. An optimized transgenesis system for *Drosophila* using germ-line-specific phiC31 integrases. *Proc Natl Acad Sci U S A* **104**: 3312-3317.
- Bushey AM, Ramos E, Corces VG. 2009. Three subclasses of a *Drosophila* insulator show distinct and cell type-specific genomic distributions. *Genes Dev* **23**: 1338-1350.

Cherbas L, Willingham A, Zhang D, Yang L, Zou Y, Eads BD, Carlson JW, Landolin JM, Kapranov P, Dumais J, et al. 2011. The transcriptional diversity of 25 *Drosophila* cell lines. *Genome Res* **21**:301-314.

Comet I, Schuettengruber B, Sexton T, Cavalli G. 2011. A chromatin insulator driving three-dimensional Polycomb response element (PRE) contacts and Polycomb association with the chromatin fiber. *Proc Natl Acad Sci U S A* **108**: 2294-2299.

Cuddapah S, Jothi R, Schones DE, Roh TY, Cui K, Zhao K. 2009. Global analysis of the insulator binding protein CTCF in chromatin barrier regions reveals demarcation of active and repressive domains. *Genome Res* **19**: 24-32.

Gaszner M, Vazquez J, Schedl P. 1999. The Zw5 protein, a component of the scs chromatin domain boundary, is able to block enhancer-promoter interaction. *Genes Dev* **13**: 2098-2107.

Georgiev PG, Gerasimova TI. 1989. Novel genes influencing the expression of the yellow locus and mdg4 (gypsy) in *Drosophila melanogaster*. *Mol Gen Genet* **220**: 121-126.

Gerasimova TI, Gdula DA, Gerasimov DV, Simonova O, Corces VG. 1995. A *Drosophila* protein that imparts directionality on a chromatin insulator is an enhancer of position-effect variegation. *Cell* **82**: 587-597.

Gerasimova TI, Corces VG. 1998. Polycomb and trithorax group proteins mediate the function of a chromatin insulator. *Cell* **92**: 511-521.

Gerasimova TI, Lei EP, Bushey AM, Corces VG. 2007. Coordinated control of dCTCF and gypsy chromatin insulators in *Drosophila*. *Mol Cell* **28**: 761-772.

Geyer PK, Corces VG. 1992. DNA position-specific repression of transcription by a *Drosophila* zinc finger protein. *Genes Dev* **6**: 1865-1873.

Golovnin A, Biryukova I, Romanova O, Silicheva M, Parshikov A, Savitskaya E, Pirrotta V, Georgiev P. 2003. An endogenous Su(Hw) insulator separates the yellow gene from the Achaete-scute gene complex in *Drosophila*. *Development* **130**: 3249-3258.

Holdridge C, Dorsett D. 1991. Repression of hsp70 heat shock gene transcription by the suppressor of hairy-wing protein of *Drosophila melanogaster*. *Mol Cell Biol* **11**: 1894-1900.

Holohan EE, Kwong C, Adryan B, Bartkuhn M, Herold M, Renkawitz R, Russell S, White R. 2007. CTCF genomic binding sites in *Drosophila* and the organisation of the bithorax complex. *PLoS Genet* **3**: e112.

Kahn TG, Schwartz YB, Dellino GI, Pirrotta V. 2006. Polycomb complexes and the propagation of the methylation mark at the *Drosophila* *ubx* gene. *J Biol Chem* **281**: 29064-29075.

Kharchenko PV, Alekseyenko AA, Schwartz YB, Minoda A, Riddle NC, Ernst J, Sabo PJ, Larschan E, Gorchakov AA, Gu T, et al. 2011. Comprehensive analysis of the chromatin landscape in *Drosophila melanogaster*. *Nature* **471**: 480-485.

Kim TH, Abdullaev ZK, Smith AD, Ching KA, Loukinov DI, Green RD, Zhang MQ, Lobanenko VV, Ren B. 2007. Analysis of the vertebrate insulator protein CTCF-binding sites in the human genome. *Cell* **128**: 1231-1245.

Kuhn-Parnell EJ, Helou C, Marion DJ, Gilmore BL, Parnell TJ, Wold MS, Geyer PK. 2008. Investigation of the properties of non-gypsy suppressor of hairy-wing-binding sites. *Genetics* **179**: 1263-1273.

Kurukuti S, Tiwari VK, Tavoosidana G, Pugacheva E, Murrell A, Zhao Z, Lobanenko V, Reik W, Ohlsson R. 2006. CTCF binding at the H19 imprinting control region mediates maternally inherited higher-order chromatin conformation to restrict enhancer access to *Igf2*. *Proc Natl Acad Sci U S A* **103**: 10684-10689.

Lancôt C, Cheutin T, Cremer M, Cavalli G, Cremer T. 2007. Dynamic genome architecture in the nuclear space: regulation of gene expression in three dimensions. *Nat Rev Genet* **8**: 104-115

Li HB, Müller M, Bahechar IA, Kyrchanova O, Ohno K, Georgiev P, Pirrotta V. 2011. Insulators, not Polycomb response elements, are required for long-range

interactions between Polycomb targets in *Drosophila melanogaster*. *Mol Cell Biol* **31**: 616-625.

Ling JQ, Li T, Hu JF, Vu TH, Chen HL, Qiu XW, Cherry AM, Hoffman AR. 2006. CTCF mediates interchromosomal colocalization between *Igf2/H19* and *Wsb1/Nf1*. *Science* **312**: 269-272.

Lobanenkov VV, Nicolas RH, Adler VV, Paterson H, Klenova EM, Polotskaja AV, Goodwin GH. 1990. A novel sequence-specific DNA binding protein which interacts with three regularly spaced direct repeats of the CCCTC-motif in the 5'-flanking sequence of the chicken c-myc gene. *Oncogene* **5**: 1743-1753.

Mohan M, Bartkuhn M, Herold M, Philippen A, Heintz N, Bardenhagen I, Leers J, White RA, Renkawitz-Pohl R, Saumweber H, et al. 2007. The *Drosophila* insulator proteins CTCF and CP190 link enhancer blocking to body patterning. *EMBO J* **26**: 4203-4214.

Moon H, Filippova G, Loukinov D, Pugacheva E, Chen Q, Smith ST, Munhall A, Grewe B, Bartkuhn M, Arnold R, et al. 2005. CTCF is conserved from *Drosophila* to humans and confers enhancer blocking of the *Fab-8* insulator. *EMBO Rep* **6**: 165-170.

Nègre N, Brown CD, Shah PK, Kheradpour P, Morrison CA, Henikoff JG, Feng X, Ahmad K, Russell S, White RA, et al. 2010. A comprehensive map of insulator elements for the *Drosophila* genome. *PLoS Genet* **6**: e1000814.

Nègre N, Brown CD, Ma L, Bristow CA, Miller SW, Wagner U, Kheradpour P, Eaton ML, Loriaux P, Sealfon R, et al. 2011. A cis-regulatory map of the *Drosophila* genome. *Nature* **471**: 527-531.

Pai CY, Lei EP, Ghosh D, Corces VG. 2004. The centrosomal protein CP190 is a component of the gypsy chromatin insulator. *Mol Cell* **16**: 737-748.

Parkhurst SM, Harrison DA, Remington MP, Spana C, Kelley RL, Coyne RS, Corces VG. 1988. The *Drosophila* su(Hw) gene, which controls the phenotypic effect of the gypsy transposable element, encodes a putative DNA-binding protein. *Genes Dev* **2**: 1205-1215.

Parnell TJ, Viering MM, Skjesol A, Helou C, Kuhn EJ, Geyer PK. 2003. An endogenous suppressor of hairy-wing insulator separates regulatory domains in *Drosophila*. *Proc Natl Acad Sci U S A* **100**: 13436-13441.

Schoenfelder S, Clay I, Fraser P. 2010. The transcriptional interactome: gene expression in 3D. *Curr Opin Genet Dev* **20**: 127-133.

Schwartz YB, Kahn TG, Nix DA, Li XY, Bourgon R, Biggin M, Pirrotta V. 2006. Genome-wide analysis of Polycomb targets in *Drosophila melanogaster*. *Nat Genet* **38**: 700-705.

Schwartz YB, Kahn TG, Stenberg P, Ohno K, Bourgon R, Pirrotta V. 2010.

Alternative epigenetic chromatin states of polycomb target genes. *PLoS Genet* **6**: e1000805.

Schmitges FW, Prusty AB, Faty M, Stützer A, Lingaraju GM, Aiwazian J, Sack R, Hess D, Li L, Zhou S, et al. 2011. Histone methylation by PRC2 is inhibited by active chromatin marks. *Mol Cell* **42**: 330-341.

Sigrist CJ, Pirrotta V. 1997. Chromatin insulator elements block the silencing of a target gene by the *Drosophila* polycomb response element (PRE) but allow trans interactions between PREs on different chromosomes. *Genetics* **147**: 209-221.


Spana C, Corces VG. 1990. DNA bending is a determinant of binding specificity for a *Drosophila* zinc finger protein. *Genes Dev* **4**: 1505-1515.


Splinter E, Heath H, Kooren J, Palstra RJ, Klous P, Grosveld F, Galjart N, de Laat W. 2006. CTCF mediates long-range chromatin looping and local histone modification in the beta-globin locus. *Genes Dev* **20**: 2349-2354.



Udvardy A, Maine E, Schedl P. 1985. The 87A7 chromomere. Identification of novel chromatin structures flanking the heat shock locus that may define the boundaries of higher order domains. *J Mol Biol* **185**: 341-358.


Wood AM, Van Bortle K, Ramos E, Takenaka N, Rohrbaugh M, Jones BC, Jones KC, Corces VG. 2011. Regulation of chromatin organization and inducible gene expression by a *Drosophila* insulator. *Mol Cell* **44**: 29-38.




Zhao K, Hart CM, Laemmli UK. 1995. Visualization of chromosomal domains with boundary element-associated factor BEAF-32. *Cell* **81**: 879-889.









A**B****C****D**