



***In vitro*, long-range sequence information for *de novo* genome assembly via transposase contiguity**

Andrew Adey, Jacob O Kitzman, Joshua N Burton, et al.

Genome Res. published online October 19, 2014

Access the most recent version at doi:[10.1101/gr.178319.114](https://doi.org/10.1101/gr.178319.114)


P<P Published online October 19, 2014 in advance of the print journal.

Creative Commons License

This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Published by Cold Spring Harbor Laboratory Press

***In vitro*, long-range sequence information for *de novo* genome assembly via transposase contiguity**

Andrew Adey¹, Jacob O. Kitzman¹, Joshua N. Burton¹, Riza Daza¹, Akash Kumar¹, Lena Christiansen², Mostafa Ronaghi², Sasan Amini², Kevin Gunderson², Frank J. Steemers², Jay Shendure^{1,*}

¹ Department of Genome Sciences, University of Washington, Seattle, WA 98115, USA

² Illumina, Inc, Advanced Research Group, 5200 Illumina Way, San Diego, CA 92122, USA.

* To whom correspondence should be addressed.

Correspondence:

Jay A. Shendure, M.D., Ph.D.
Department of Genome Sciences
3720 15th Avenue NE Box 355065
Seattle, WA 98195-5065
Phone: 206-685-8543
Fax 206-685-7301
Email: shendure@uw.edu

Keywords: Genome Assembly, Transposase

Abstract

We describe a method that exploits contiguity preserving transposase sequencing (CPT-seq) to facilitate the scaffolding of *de novo* genome assemblies. CPT-seq is an entirely *in vitro* means of generating libraries comprised of 9,216 indexed pools, each of which contains thousands of sparsely sequenced long fragments ranging from 5 kilobases to over 1 megabase. These pools are “sub-haploid”, in that the lengths of fragments contained in each pool sums to approximately 5 to 10% of the full genome. The scaffolding approach described here, termed *fragScaff*, leverages coincidences between the content of different pools as a source of contiguity information. Specifically, CPT-seq data is mapped to a *de novo* genome assembly, followed by the identification of pairs of contigs or scaffolds whose ends disproportionately co-occur in the same indexed pools, consistent with true adjacency in the genome. Such candidate “joins” are used to construct a graph, which is then resolved by a minimum spanning tree. As a proof-of-concept, we apply CPT-seq and *fragScaff* to substantially boost the contiguity of *de novo* assemblies of the human, mouse, and fly genomes, increasing the scaffold N50 of *de novo* assemblies by 8 to 57 fold with high accuracy. We also demonstrate that *fragScaff* is complementary to Hi-C-based contact probability maps, providing mid-range contiguity to support robust, accurate chromosome-scale *de novo* genome assemblies without the need for laborious *in vivo* cloning steps. Finally, we demonstrate CPT-seq as a means of anchoring unplaced novel human contigs to the reference genome as well as for detecting misassembled sequences.

Introduction

The broad adoption of next-generation sequencing (NGS) has resulted in a proliferation of *de novo* genome assemblies (Pagani et al. 2012). For the most part, these assemblies are of far lower quality than the reference genomes produced by The International Human Genome

Consortium (The International Human Genome Consortium 2001; The International Human Genome Consortium 2004; Shemesh et al. 2013; Rong and Gardener 2013; Wang et al. 2013; Blanco-Ulate et al. 2013), largely secondary to a dearth of readily accessible NGS-based methods for generating mid-range and long-range contiguity information. Conventional NGS-based assemblies rely on deep sequencing of shotgun fragment and ~3 kbp mate-pair libraries to provide short-range contiguity, both of which are generated via straightforward *in vitro* protocols. Dilution pools of PCR amplicons can provide long virtual reads, but these are limited to 10 kbp by the use of PCR and the majority are ~8 kbp (Voskoboynik et al. 2013). Mid-range contiguity information requires still-expensive (but rapidly evolving) long sequencing reads (e.g. PacBio at ~18 kbp (Koren et al. 2012) or nanopore sequencing (Laszlo et al. 2014)) or labor-intensive fosmid or BAC clone libraries (Gnerre et al. 2011; Zhang et al. 2012). Alternatively, long-range contiguity information can be generated *in vitro* via contact probability maps, wherein Hi-C read-pairs are used for chromosome-scale scaffolding (Burton et al. 2013). However, the performance of Hi-C-based scaffolding depends heavily on input assembly scaffold size, with optimal results requiring an input N50 of ~200 kbp or greater, a level of contiguity that can be challenging to achieve with shotgun fragment and 3 kbp mate-pair libraries alone (Shemesh et al. 2013; Rong and Gardener 2013; Wang et al. 2013; Blanco-Ulate et al. 2013). As such, there remains a strong need for robust *in vitro* methods to capture mid-range contiguity information for *de novo* genome assembly.

Transposase-mediated library construction, or “tagmentation”, utilizes a hyperactive Tn5 transposase to both fragment and append universal adaptors in a single enzymatic step (Goryshin and Reznikoff 1997; Adey et al. 2010). In recent years, tagmentation has been applied in diverse ways including DNA-seq (Adey et al. 2010), stranded RNA-seq (Gertz et al. 2012), whole-genome bisulfite sequencing (Adey and Shendure 2012), chromatin profiling (Buenrostro et al. 2013) and *in situ* mate-pair library preparation directly on a sequencing

flowcell (Schwartz et al. 2012). We recently demonstrated a novel method, contiguity preserving transposase sequencing (CPT-seq) for haplotype-resolved genome sequencing (Amini et al. 2014). CPT-seq utilizes an inherent property of the Tn5 transposase in which the enzyme remains tightly bound to the target DNA after tagmentation, physically linking adjacent library molecules (Fig. 1A). Prior to PCR amplification, the high molecular weight products of linked templates are subjected to sub-haploid dilution and compartmentalization, followed by protein denaturation in order to free the templates for amplification. To increase the effective number of compartments, a two-tiered indexing approach (Erlich et al. 2009) is applied (Fig. 1B). The initial tagmentation is performed using 96 uniquely indexed transposase-adaptor complexes. The high molecular weight linked templates are then pooled, followed by limiting dilution into 96 indexed PCR reactions such that each PCR well contains templates from 96 originating transposase reactions, thus producing $96 \times 96 = 9,216$ distinct index combinations. Sequence reads corresponding to each of the 9,216 “virtual compartments” (also referred to as indexed pools) provide sparse, shotgun representation of the high molecular weight genomic DNA fragments that were transposed within that indexed pool (Fig. 1C), analogous to fosmid (Kitzman et al. 2011) or *in vitro* dilution pools (Kaper et al. 2013) but with a ~100x higher effective number of pools. With appropriate dilution of the input genomic DNA, these indexed pools are “sub-haploid”, in that the lengths of fragments contained in each pool sums to approximately 5 to 10% of full genome length. From an experimental perspective, the library preparation steps of CPT-seq are straightforward and scalable, with an overall processing time of less than 3 hours, relying on readily available equipment and requiring minimal hands-on time (Amini et al. 2014).

Results

We speculated that the large number of effective dilution pools in CPT-seq ($n = 9,216$), each of which contains thousands of long DNA fragments (Fig. 1D), might be an excellent source of

mid-range contiguity information for *de novo* genome assembly. Many contemporary mid-range contiguity methods involve mate-pair sequencing, in which each read in a sequenced pair is derived from the end of a long genomic DNA fragment, e.g. 3 kbp to 40 kbp in length. These methods can use existing software to perform scaffolding such as Bambus2 (Koren et al. 2011), GRASS (Gritsenko et al. 2012), SCARPA (Donmez and Brudno 2013), or SOAPdenovo2 (Luo et al. 2012), which were evaluated extensively against one another and several other tools by Hunt *et. al.* (2014) as well as other *de novo* assembly algorithms evaluated as part of the Assemblathon project (Earl et al. 2011; Bradnam et al. 2013). However, the data produced by CPT-seq is very different from that of mate-pair sequencing, consisting instead of sparse, shotgun representation of genomic DNA fragments that were transposed within each of 9,216 pools, wherein each pool derives from an essentially random mix of thousands of variably sized, high molecular weight genomic DNA fragments. We therefore developed a new algorithm, *fragScaff*, which leverages coincidences between the content of different pools as a source of contiguity information for linking proximally located sequences for *de novo* genome assembly (Fig. 2, Supplemental Fig. S1, Supplemental Note).

The input to *fragScaff* consists of an initial genome assembly (e.g. based on shotgun and mate-pair sequencing data) and data produced by CPT-seq. First, the input contigs' or scaffolds' (referred to here as input contigs) lengths are used to determine the "end-nodes", or sequence at the ends of each input contig to be utilized in the assembly process, thus producing an end-node pair for each input contig. The alignment of CPT-seq reads to the input assembly is then used to determine which of the 9,216 CPT-seq pools have quality alignments ($q \geq 10$) to each of the end-nodes. The distribution of the number of groups hitting the end-nodes (Supplemental Fig. S2) is then used to exclude those at the extremes which may contain repetitive sequence (high extreme) or misassembled sequence (low extreme).

In the next step, which makes up the core of the *fragScaff* algorithm, the shared fraction of pools between every possible node pair is calculated as the number of pools hitting both of the nodes divided by the total number of unique pools hit by either node. This results in a distribution of shared fractions for each individual node (Fig. 2B) for which a mean and standard deviation can be calculated and used to estimate the probability that the reciprocal node is an outlier. A p -value threshold is then manually set or automatically determined based on the number of outliers that would produce a pre-defined target mean number of outliers per node (Supplemental Table S1). In cases where two nodes are called as outliers with respect to the distribution of the other it is considered a “reciprocated link” and forms an edge in the subsequent graph manipulation steps with a weight corresponding to the outlier score defined as the $-\log_{10}(p\text{-value})$.

The third and final stage of *fragScaff* is the ordering of end-nodes based on the constructed graph (Fig. 2C, Supplemental Fig. S3). The maximum-weight minimum spanning tree (MST) is first identified for each connected component and the longest path (diameter) through the MST is identified as the trunk. An MST approach was chosen because the majority of connected components are already very close to their MST and only very few edges are removed for the purpose of trunk identification. After the trunk is established, the end-nodes not present in the trunk (branches) are placed using an iterative approach where the end-node with the highest weighted edge to a trunk end-node is inserted into the trunk path such that the new trunk path score is maximized. This method utilizes all edges as opposed to solely the MST edges and is iterated until every branch end-node is incorporated.

Run times and memory requirements of *fragScaff* are highly dependent on both the size of the input assembly as well as the number of input contigs present (Supplemental Note). For efficiency in optimizing the scaffolding process, each stage of *fragScaff* can be run independently to allow for multiple iterations of the fast graph manipulation stage using varying

cutoffs (Supplemental Table S1), without re-running the more time consuming I/O steps or the calculation of pool overlaps. For the determination of end-nodes and BAM file parsing, the number of reads is the primary constraint, with run times of 4 hours for 1,324 M reads (human) and 40 minutes for 334 M reads (fly) using a single core and requiring a <1 GB memory footprint. The all-by-all node calculation is generally the most computationally intensive step, requiring 22 minutes for the human assembly (N50 = 437 kbp, 18,922 input contigs, 50 threads executed via SGE (Sun Grid Engine)). This increases sharply with higher contig counts due to the $O(n^2)$ complexity, reaching 16 hours using 50 threads (via SGE) for the 25 kbp simulated human assembly which contained 115,162 input contigs. Finally, the graph manipulation steps are minimally intensive, requiring 2 minutes on a single core for the human assembly (N50 = 437 kbp, 18,922 input contigs).

We applied *fragScaff* to a human (GM12878) *de novo* assembly with an N50 scaffold size of 437 kbp generated by *ALLPATHS-LG* using only shotgun and 3 kbp mate-pair libraries (Gnerre et al. 2011). CPT-seq data was generated on GM12878 DNA (Amini et al. 2014), yielding 9,216 pools with a mean 2,513 fragments \geq 5 kbp per pool (23.2 M fragments in total with a mean fragment size of 33.9 kbp and 0.9 M fragments $>$ 100 kbp). The resulting *fragScaff* assembly using conservative parameters incorporated 97.1% of sum length of the input contigs, resulting in an N50 of 4.4 Mbp (10 \times increase) with a 97.1% join accuracy, a 92.4% orientation accuracy, and 98.5% of base pairs properly placed (Table 1, Supplemental Table S2, Supplemental Fig. S4-5) including a perfectly ordered scaffold of 23.9 Mbp. Additionally, we performed several less stringent iterations of *fragScaff* which resulted in N50 increases up to 42 \times (N50 = 18.2 Mbp) while still maintaining a 97.8% base pair placement accuracy.

It is important to note that unlike mate-pair sequencing, in which orientation information is captured via read pair alignment orientation, *fragScaff* has no such information and relies on the link scores of each input contig end-node. Therefore as input contig size decreases, the spacing

between the end-nodes decreases and so does the ability to infer orientation, reaching zero when the end-nodes of an input contig completely overlap (Supplemental Fig. S6). When excluding end-nodes for which there is no orientation information (i.e. where end-nodes completely overlap), the orientation accuracy increases to 96.8%. In situations where orientation information is limited, the quality scores generated by *fragScaff* are of particular importance to inform whether or not the reverse orientation should be considered in downstream analyses.

We also attempted to apply *fragScaff* to data generated with fosmid (Kitzman et al. 2011) or long fragment read (LFR) (Kaper et al. 2013) dilution pools, rather than CPT-seq. However, this resulted in low link counts of poor accuracy (N50 improvement: 1.3x, and 1.5x; join accuracy: 71.6% and 34.0%; sequence joined: 38.9% and 46.6% for fosmid and LFR respectively), predominantly due to the reduced pool counts which restrict the number of coincidences between pool content on which this method is based (fosmid: 288, LFR: 96 vs. CPT-seq: 9,216). Consistent with this, we observed markedly worse performance when down-sampling to a low number of CPT-seq pools (e.g. 47.9% (288 pools) vs 87.9% (9,216 pools) of sequence scaffolded for fly; see below).

As achieving a scaffold N50 of even 437 kbp with shotgun and 3 kbp libraries is quite challenging for complex genomes (Shemesh et al. 2013; Rong and Gardener 2013; Wang et al. 2013; Blanco-Ulate et al. 2013), we sought to evaluate this approach on less contiguous input assemblies. As a first analysis, we fragmented the human input *de novo* assembly at every gap to produce a contig-only assembly (N50 reduced from 437 kbp to 47 kbp) and then applied *fragScaff* with CPT-seq data. This resulted in an assembly that joined 97.7% of input contig sequence for a scaffold N50 of 2.7 Mbp (57x improvement), 97.9% join accuracy, and 99.2% of base pairs properly placed. However, the reduced contiguity of the input assembly and the effective exclusion of 3 kbp mate-pair data resulted in a substantial decrease in orientation accuracy to 71.4%. As a second analysis, we fragmented the human reference genome *in silico*

to sizes ranging from 15 to 300 kbp. The scaffolding of these simulated input assemblies with *fragScaff* resulted in N50 improvements ranging from 24x to 75x, with greater than 99.3% of base pairs properly placed and 96.0% properly oriented in every assembly (Table 1, Supplemental Table S2).

We also evaluated *fragScaff* and CPT-seq with *Mus musculus* (mouse) and *Drosophila melanogaster* (fruit fly), for which high quality reference genomes are available for comparison. For both organisms, we used assemblies generated using *ALLPATHS-LG* from either shotgun and mate-pair sequencing (mouse, N50 = 224 kbp) or just shotgun sequencing (fly, N50 = 68 kbp) (Gnerre et al. 2011; Burton et al. 2013). CPT-seq data and *fragScaff* were then used to increase the N50 to 2.9 Mbp for mouse (13x improvement) and 524 kbp for fly (7.7x improvement) with 98.1% and 99.7% of base pairs properly placed for mouse and fruit fly respectively (Table 1, Supplemental Table S2).

We recently reported the use of Hi-C data for chromosome-scale *de novo* assembly on human, mouse, and fruit fly genomes (Burton et al. 2013). For input assemblies with a large N50, Hi-C-based scaffolding produced high-quality scaffolds such as on human (input assembly with N50 = 437 kbp) where the resulting output clustered 98.2% of sequence into chromosome groups with 94.4% ordered. However, on the smaller fly input assembly (N50 = 68 kbp) Hi-C-based scaffolding was able to cluster only 81.2% and order 82.0% of sequence. Similarly, when we applied Hi-C-based scaffolding to our human contig assembly (N50 = 47 kbp), 89.7% of sequence was clustered with only 41.5% ordered, and even less for the simulated 15 kbp contig assembly (35.9% clustered, 0.2% ordered). However, when we perform CPT-seq and *fragScaff* prior to Hi-C-based scaffolding, the completeness and quality of the resulting chromosome-scale *de novo* assembly markedly improves for the human contig assembly (Table 2, Supplemental Table S3) with 99.4% of sequence clustered (gain of 9.7%) and more than doubling the amount of sequence ordered to 99.1% (gain of 57.6%). Similar improvements were

also observed for simulated contig assemblies of fly and human, most strikingly for the 15 kbp human simulated contig input wherein the proportion of the input contig assembly that is ultimately clustered and ordered increased from 35.9% to 91.9%, and from 0.2% to 88.2%, respectively. These results demonstrate that CPT-seq and *fragScaff* can improve the contiguity of assemblies generated from shotgun and short-range mate-pair libraries to a point suitable for chromosome-scale scaffolding (Supplemental Fig. S7).

In Kitzman *et al.* (2011) we described a method that utilized the sub-haploid content of each fosmid pool to anchor *de novo* assembled contigs from reads that did not align to the human reference genome, as well as a set of previously anchored sequences (hg18) described in Kidd *et al.* (2010) (Supplemental Fig. S8). This method worked on the premise that each window in the genome is hit by a discrete set of pools, as is each novel contig, and the window with maximum pool overlap with a novel contig is the most probable anchor location. We applied this approach to anchor the same set of contigs to hg18 (many of these have since been incorporated to the GRCh37 human reference assembly) but used CPT-seq data generated on GM12878, thus increasing the number of pools, and therefore anchoring power, from 115 to 9,216. We were able to place 1,816 of the 2,363 contigs (76.9%), though a number of unplaced contigs are likely population-specific and not present in GM12878 (Kidd *et al.* 2010; Genovese *et al.* 2013). 1,226 of our placed contigs were previously placed by Kidd *et al.* (2010), with 1,154 (94.1%) in agreement with our calls.

We next evaluated whether the same method used for anchoring unplaced genomic content could be applied to detect misassemblies in a *de novo* genome assembly, *i.e.* by examining the pool overlap fractions of immediately adjacent windows and windows one apart (Figure 3A). For example, using this method, 3,081 suspicious windows from 926 scaffolds were identified in the *de novo* assembly based on shotgun, 3 kbp mate-pair and CPT-seq data (N50 = 4.4 Mbp), of which 400 scaffolds (43% of those flagged) contained bona fide misassemblies upon

comparison with the GRCh37 reference assembly (Figure 3B). These misassemblies resulted in reduced pool overlap fractions over a span of several windows as opposed to false positive flaggings which were generally single window calls. Of course, the set of misassemblies identified in this example is biased towards those that were present in the input genome, and we have less power to detect misassemblies by *fragScaff* due to utilization of the same data that went into the scaffolding process.

Lastly, we sought to take advantage of the haplotype-specific nature of CPT-seq data (Amini et al. 2014) by haplotype-resolving the GM12878 *fragScaff* assembly (input N50 = 437 kbp, *fragScaff* N50 4.4 Mbp), *i.e.* to produce a diploid *de novo* genome assembly. We first aligned shotgun reads back to the assembly (56×) and performed variant calling to identify 902,905 heterozygous sites, substantially fewer than the 2,180,767 high quality GM12878 heterozygous sites previously described (Zook and Salit 2013). Furthermore, only 327,828 (36.3%) of variants could be uniquely assigned to reference genome coordinates (GRCh37) that matched the coordinates of known GM12878 variants. Of the 575,077 unassigned variant calls, 50.6% aligned to repeat elements and 34.8% aligned to segmental duplications, indicating that the majority are false positive variants secondary to collapsed regions in the original assembly.

We then sought out to perform haplotype resolution of either the full ($n = 902,905$) or validated ($n = 327,828$) sets of heterozygous sites using CPT-seq data and *ReFHap* (Duitama et al. 2010). For the full set, the high burden of false positive calls confounded the graph-based algorithm and was abandoned after 164 hours of run time. On a smaller subset of scaffolds containing only 8,437 variants, *ReFHap* completed but produced an essentially random result with a pairwise phasing accuracy of 50.1%. For the validated set, haplotype resolution was achieved in under an hour, with 82.4% of variants phased with a haplotype block N50 of 356 kbp (max 4.4 Mbp) and a pairwise phasing accuracy of 95.0%. Thus, while we were able to successfully achieve haplotype resolution concurrent with *de novo* genome assembly, this was

only possible by restricting the analysis to variants validated by alignments to the human reference genome. This result underscores the need for experimental methods and/or algorithms that improve repeat disambiguation in *de novo* genome assemblies.

Discussion

We demonstrate that a new, entirely *in vitro*, method of transposase-mediated library construction, CPT-seq, which preserves the contiguity of high molecular weight transposed molecules, can be used to provide mid-range sequence information. This library construction method results in 9,216 distinct groups of reads, each of which contains sparse sequence reads over a set of thousands of long 5 kbp to 1 Mbp fragments that sum up to approximately 5-10% of the target genome.

We emphasize that the quality of input genomic DNA is a critical parameter in the performance of this method (and more generally, methods based on sub-haploid complexity reduction (Lo et al. 2013)). The decreasing abundance of fragments at longer lengths (Fig. 1D) is largely due to fragmentation occurring at the transposase incorporation step and subsequent handling prior to PCR; however the maximum size of fragments is also inherently limited by the quality of the input material. As such, CPT-seq is best performed on DNA freshly isolated using protocols that minimize fragmentation (Methods).

In order to use this new form of sequence data to aid in *de novo* genome assembly, we designed and implemented a novel algorithm, *fragScaff*, which produces mid-range scaffolding comparable to 40 kbp fosmid-based mate-pair libraries, yet uses the entirely *in vitro* CPT-seq method. Furthermore, the algorithm can be tuned to allow for longer assemblies at a minor cost to accuracy depending on downstream requirements and provides detailed quality score information on the joins and sequence orientations present in the final assembly (Supplemental Note). Beyond the scaffolding of contigs, we were also able to utilize CPT-seq data to

haplotype-resolve a *de novo* genome assembly, thus producing a diploid assembly. However, this was only possible by restricting the analysis to validated heterozygous variants, as the large number of false positive variant calls due to collapsed repeat elements confounds algorithms for haplotype resolution. In future work, this may be circumvented by improving repeat disambiguation during the genome assembly process, or by improving methods for calling truly heterozygous variants within imperfect genome assemblies.

The primary limitation of CPT-seq and *fragScaff* at present is that for short input contigs (<3 kbp) are difficult to scaffold confidently due to a reduced number of pools covering the contig. In these situations, removing the shorter contigs prior to *fragScaff* followed by placement using the described anchoring method may be preferred to avoid potential misassemblies during the scaffolding process. A second limitation of the method is that CPT-seq does not provide orientation information that is inherent in more traditional mate pair sequencing approaches. As such, the only means of orientation is the difference in signal between the end-node pairs of each input scaffold or contig. For longer input contigs, in which the end-nodes do not overlap, orientation is possible with high accuracy; however input contigs shorter than twice the end-node size suffer from an increase in shared pools due to the overlapping end-nodes and therefore orientation accuracy rapidly decreases to random where the end-nodes overlap completely. An additional challenge that is shared by *fragScaff* and traditional mate-pair based scaffolding is that unrecognized repeat elements will lead to false joins. This will be particularly problematic for genomes that contain large numbers of closely related sequences. Indeed, the accuracies demonstrated here are dependent on robust repeat masking prior to scaffolding. Given that the termini of many or most contig/scaffold ends after the primary *de novo* assembly are repetitive, an advantage of *fragScaff*, relative to traditional mate-pair based scaffolding, is that the end-node can effectively be defined as the unique sequence just internal to this.

A key component of CPT-seq and its ability to generate useful data for either haplotype phasing as described in Amini *et. al.* (2014) or for use in *de novo* assembly scaffolding via *fragScaff* is the partitioning of fragments such that each pool contains 1-10% of the organism's haploid genome. The distribution of fragments occurs after the indexed tagmentation of the high molecular weight DNA via dilution; such that each of the 96 PCR reactions contains approximately 96 times the amount of desired long fragments per pool (e.g. a sum contiguous transposed template length of 32-320 Mbp per pool for roughly 0.3-3 Gbp per PCR reaction for human). On larger genomes, such as human and mouse, the size of the haploid genome allows for only modest dilution factors, with substantially more dilution required for fly. While CPT-seq performed well on the modest, ~170 Mbp fly genome, it is likely that the dilution factor required for genomes in the sub-10 Mbp range may be challenging or prohibitive. However, the pooling of genomes prior to the CPT-seq workflow may allow for a workaround with the added advantage of increased parallelization as long as special care is taken to insure the pooled genomes are sufficiently diverged (e.g. spiking in a bacterial genome into a primate CPT-seq prep). On a related note, we recently described the use of Hi-C data for the purposes of metagenome deconvolution and assembly (Burton *et. al.* 2014). In principle the contiguity provided by CPT-seq could be used in a similar fashion; however the varying abundances of species within a metagenome would likely render titration to 1-10% representation of each target genome very challenging or impossible for most samples. Nonetheless, this approach could be fruitful for subsets of species within a metagenome that present are at approximately equal abundances.

The ease of CPT-seq library preparation and the amount of information obtained per base of sequencing performed makes it a very appealing tool for routine use in *de novo* genome assembly. These methods bridge the gap between short-range assemblies and the scaffold sizes needed to perform chromosome-scale scaffolding with Hi-C-based methods. We envision

that the combination of four types of sequencing libraries – shotgun fragment, 3 kbp mate-pair, CPT-seq, and Hi-C – will provide a robust strategy to *de novo* assemble complex genomes to chromosome-scale contiguity using entirely *in vitro* methods that can be carried out cheaply and rapidly.

Methods

Library Construction

Genomic DNA was prepared using the Qiagen Genra Puregene Kit on 10 mg fresh liver (*mus musculus*, BL6) and 10 flies (*Drosophila melanogaster*, Canton S.) and analyzed on a pulsed field gel (Supplemental Fig. S10). CPT-seq libraries for human (GM12878) were those used in Amini *et. al.* (2014) are available from the Sequence Read Archive (SRA; <http://www.ncbi.nlm.nih.gov/sra>) under project accession PRJNA241346. For mouse and fly, CPT-seq libraries were prepared and sequenced using custom chemistry as described in Amini *et. al.* (2014) with a slight modification to PCR template amounts for the fly sample to account for substantially decreased genome size (1, 2, or 3 pg inputs were used in each PCR reaction as opposed to 10 pg for Human and Mouse, Supplemental Fig. S11). Fosmid libraries were previously generated for Adey *et. al.* (2013). Long fragment read (LFR) libraries were generated on CEU gDNA using methods outlined in Kaper *et. al.* (2013).

Input Assemblies

The human, mouse, and fly input assemblies were the same input assemblies generated using *ALLPATHS-LG* as described in Burton *et. al.* (2013) using reads accessed under Sequence Read Archive (SRA; <http://www.ncbi.nlm.nih.gov/sra>) accessions SRA024407 (Human, GM12878), SRA009956 (Mouse), and SRR516038 & SRR516001 (Fly). For an additional human assembly of lower quality, we broke up the original assembly into individual contigs by splitting scaffolds at any N base (N50 437 → 47 kbp). Simulated input assemblies were

generated by splitting the human reference sequence at regular intervals. In cases where repeat content was considered for the determination of input contig end-nodes, the entire Repbase version 19.03 database fasta (Jurka et al. 2005) was aligned to the input assembly using blast with default parameters followed by the merging of entries in the form of a bed file and including all segments ≥ 50 bp (half of the CPT-seq read length). End nodes were then adjusted to include additional unique sequence to compensate for the exclusion of repeat sequence. Reads in which $\geq 50\%$ of the read aligned to a repeat element were excluded from any further analysis.

Contig Scaffolding

Reads were aligned using *BWA* (Li and Durbin 2010) to references created from the input assemblies. The ends of each input contig (1 kbp fosmid and LFR, 5-10 kbp CPT-seq) were then used as end-nodes and the pools that had reads aligning to these end-nodes were considered hits (reads that aligned to flagged repeats were excluded, Supplemental Note). The distribution of the number of pools hitting each end-node was then used to exclude the top and bottom 5% of end-nodes. This exclusion does not necessarily remove the entire input contig, as the paired end-node at the other end of the input contig may allow scaffolding; however orientation information would be substantially weaker (Supplemental Fig. S2). Each end-node was then analyzed individually by calculating the fraction of pools that overlap with all other end-nodes ($\text{shared_pools} / (\text{total_node1_pools} + \text{total_node2_pools} - \text{shared_pools})$) and subsequently calculating the mean and standard deviation for the end-node. p -value cutoff thresholds were then determined by finding the $-\log_{10}(p\text{-value})$ score that would result the desired mean number of outliers per end-node. End-nodes (node 1) that have an outlier end-node (node 2) where node 1 is also an outlier in node 2 were then considered as potential reciprocated links with an edge weight of the combined $-\log_{10}(p\text{-value})$ score of each outlier, and stored as an included edge if that weight was above a set threshold based on a multiple of the $-\log_{10}(p\text{-value})$ cutoff score (Supplemental Table S1, Supplemental Note). End-nodes that

were at the ends of the same input contig (end-node pairs) were automatically given an edge score beyond the maximum score cutoff to prevent input contig splitting. Connected components in the graph were then determined followed by identification of the maximum weight Minimum Spanning Tree (MST) by implementing Prim's algorithm. The true longest path (trunk) was then determined by identifying every path between degree one end-nodes and taking the maximum length path. Since a typical MST in *fragScaff* has very few branches, the brute-force approach to finding the trunk can be implemented without excessive compute times. Non-trunk end-nodes (branches) were then placed by inserting the branch into the trunk at a position that allows for the maximum trunk path edge weight based on all edges (not only the MST edges).

fragScaff Parameters

fragScaff was run with varying parameters based on input assembly size and the desired output contiguity. The primary variable parameters are (i) end-node size, (ii) mean passing links per end-node, and (iii) link reciprocation factor (Supplemental Table S1); however a number of other options can be modified, including the use of a repeat-masking bed file to exclude reads in identified repeats (Supplemental Note). For the end-node size, the primary determining factor was the N90 of the input scaffolds. The end-node size was set to a minimum of 5 kbp with an optimal size at approximately $\frac{1}{2}$ of the input N90, up to a maximum of 10 kbp. A small N50 generally requires an increase to the mean passing links per end-node as well as modifications to the combined score cutoff. These parameters (j and u) are optimized in Supplemental Table S1 for each assembly. A detailed description of all parameters used in *fragScaff* is provided in the Supplemental Note.

Scaffolding Accuracy Measurement

Join accuracy was assessed only at locations where consecutive joined input scaffolds are properly mapped back to the reference assembly and were considered an accurate join if the scaffolds were within 5 of one another in the ordered rank. Orientation accuracy was also only assessed for consecutive, reference aligned scaffolds and considered accurate if the correct ends of the input scaffolds were joined. The fraction of base pairs properly placed was assessed by totaling the amount of correctly joined sequence in all scaffolds and then totaling the amount of sequence in all scaffolds that did not belong to the dominant locus of that scaffold. For example, if 3.5 Mbp of a scaffold was properly joined in correct order, and was fused with another set of properly joined scaffolds that was 2 Mbp in length, we would consider the 2 Mbp set of sequence as improperly placed bases. Scaffolds were considered fused if more than 10% of the sequence in the scaffold was considered improperly placed. A graphical representation of accuracy measurements can be found in Supplemental Fig. S4.

Contig Anchoring

CPT-seq reads were aligned to the human reference assembly using *BWA* (Li and Durbin 2010) (to hg18, as GRCh37 contains a number of the sequences we were anchoring). Fragments were then called in each pool using an alignment-distance cutoff of 15 kbp (Amini et al. 2014) on reads with a mapping quality ≥ 10 . Unaligned reads were then aligned using *BWA* (Li and Durbin 2010) to a reference of all of the unanchored contigs. Pools with at least one read aligned with a mapping quality ≥ 10 were then considered hits. Pools with fragments spanning windows of 1-5 kbp in the reference genome were then identified (Supplemental Fig. S8). For each contig, the fraction of window pools shared with the contig pools and the fraction of contig pools shared with the window were calculated for each window, sorted and ranked. The window with the top combined rank was then assigned as the anchor position. Agreement with published loci was determined by checking if our anchoring position was within 1 Mbp of the published position; however 96% of the agreement loci were within 100 kbp.

Misassembly Detection

CPT-seq reads were aligned to the resulting *fragScaff* assembly using *BWA* (Li and Durbin 2010) to assign a pool set for every 5 kbp window (excluding N's) in the assembly. The shared pool fraction for immediately adjacent windows as well as for windows one apart was calculated. Window junctions that had shared pool fractions in the bottom 5th percentile for both immediately adjacent windows and windows one apart were flagged as putative misassemblies. Regions flanking the misassembled regions were aligned to the human reference via BLAST and analyzed whether or not they should be adjacent as well as whether or not the detected misassembly falls at a *fragScaff* join.

Haplotype Resolution

GM12878 shotgun reads used in the original *ALLPATHS-LG* assembly (SRA024407, N50 = 437 kbp) were aligned onto the conservative *fragScaff* assembly (N50 = 4.4 Mbp) using *BWA* (Li and Durbin 2010). Variants were then called using Samtools (Li et. al. 2009) and filtered to remove indels and require coverage of at least 10, at least one read for each allele on each strand, and an allele balance > 0.25 and < 0.75 . The called heterozygous sites were then validated against a high quality reference set produced by the NIST (Zook and Salit 2013) on GM12878 by pulling the flanking 50 bp of assembled sequence around each called variant and aligning the 101 bp segments to the human reference genome (GRCh37) using *BWA* (Li and Durbin 2010). The variants that were uniquely aligned and matched the position and alleles of the reference set were then utilized for haplotype resolution. CPT-seq reads were then aligned to the reference and used to call fragments by a thresholding approach by which a distance between subsequent aligned reads in a given read group must be $\leq 15,000$ bp else it is determined to be a break between fragments (Amini et. al. 2014). During fragment calling, genotyping was performed on the fragments at the previously validated heterozygous sites. The

genotyped fragments were then subjected to haplotype phasing using *ReFHap* (Duitama et. al. 2010) and accuracy was determined by mapping the phasing of the validated sites back to the human reference genome positions and then compared to known haplotypes determined by pedigree information (Supplemental Fig. S9).

Data Access

Mouse and Fly CPT-seq data generated for this study have been submitted to the NCBI

Sequence Read Archive (SRA; <http://www.ncbi.nlm.nih.gov/sra>) under accession numbers

SRP041913 (mouse) and SRP041914 (fly).

Software Availability

The *fragScaff* software is available for download from Sourceforge:

<https://sourceforge.net/projects/fragscuff/files/>

Acknowledgements

We thank R. Patwardhan for generating *ALLPATHS-LG* assemblies and other members of the Shendure Lab for helpful discussion. We would also like to thank M. Wilken and the Reh lab for supplying a fresh mouse liver, and N. Peters and the Berg lab for supplying fly cultures. Our work was supported by grant HG006283 from the National Human Genome Research Institute (NHGRI; to J.S.); a graduate research fellowship DGE-0718124 from the National Science Foundation (to A.A. and J.O.K.); and grant T32HG000035 from the NHGRI (to J.N.B.).

Disclosure Declaration

Several authors of this manuscript are employed by Illumina Inc.

Figures and Tables

Figure 1. CPT-seq method and performance

A. High Molecular Weight (HMW) genomic DNA is reacted with hyperactive Tn5 transposase loaded with indexed adaptors. After the transposase complex fragments the DNA and appends the indexed adaptors, the enzyme remains tightly bound to the DNA, such that library molecules derived from the same HMW genomic DNA molecule remain physically linked. Once the transposase is removed by denaturation, PCR amplification of viable templates (gray boxes) can be performed. **B.** Schematic of two tier indexing. A 96-plex indexed tagmentation is performed (but without removing the transposase), followed by pooling, mixing, and redistribution to 96 wells. These new pools are subjected to removal of the transposase, 96-plex indexed PCR and then pooling to a single sequencing library. Individual molecules within the final library have indices corresponding to both the pool in which their originating HMW genomic DNA fragment was present during tagmentation (96 indices) as well as during PCR (96 indices), such that there are effectively $96 \times 96 = 9,216$ compartments. **C.** Representation of coverage profiles for indexed fragment pools, *i.e.* compartments, (top) and trimodal distribution of adjacently aligning reads within individual compartments. The first peak (~100 bp, red) corresponds to simple read pairs; the second peak (~3.2 kbp, green) corresponds to reads originating from the same HMW genomic DNA fragment; the third peak (~1 Mbp, blue) corresponds to reads originating from different HMW genomic DNA fragments. **D.** Distribution of estimated HMW genomic DNA fragment lengths for CPT-seq of GM12878. The mean fragment size is 33.9 Kb, but it is a broad distribution and nearly 1M fragments are >100 kbp.

Figure 2. *fragScaff* assembly method

A. The ends of contigs in a *de novo* genome assembly (gray boxes) are defined as nodes, and the subsets of the 9,216 CPT-seq compartments, *i.e.* indexed pools, containing reads that align to each node are identified. The fraction of shared compartments between every possible pair of nodes is calculated. Pairs of nodes that are truly adjacent to one another in the genome are expected to exhibit excess sharing with respect to CPT-seq compartments, as a result of HMW genomic DNA fragments that bridge the gap in the *de novo* genome assembly. Non-adjacent pairs of nodes will co-occur in a small fraction of compartments by chance, as each contains HMW genomic fragments that cover ~10% of the genome. **B.** The fraction of shared compartments is calculated for all possible pairs of nodes, and distributions are generated for each node. Outlier nodes in each distribution are identified assuming normality and using a *p*-value cutoff. If a link is reciprocated, *i.e.* if two nodes are each outliers in the other's distribution, it is stored as an edge. **C.** Sub-graphs are reduced to their Minimum Spanning Tree (MST) and the longest path (Trunk) is found. Branches (light nodes) are then placed to produce the final output scaffold. **D.** Size distribution of gaps between properly linked contigs. Boxes indicate joins spanning gaps just beyond the 2.5 kbp mate-pair library (red), ~6 kbp L1 repeat elements (green), and joins longer than 35 kbp which cannot be achieved via fosmid mate-pair libraries (blue; *n* = 664).

Figure 3. Misassembly detection using CPT-seq

A. Three regions of assembled scaffolds representing various misassembly detections are shown. For each region, the set of CPT-seq indexed pools that cover each 5 kbp window (x-axis) was determined. The shared fraction of indexed pools between immediately adjacent windows (blue) and for windows one apart (purple) is plotted. Sub-regions for which both shared fraction values were in the bottom 5th percentile overall were called as potential misassemblies. False positive misassembly calls (*i.e.* no misassembly is actually present) overwhelmingly consisted of an isolated window (green shading), while multiple consecutive windows with low shared fraction values corresponded to misassemblies by *fragScaff* (yellow shading) or in the initial input assembly (red shading). **B.** Breakdown of regions called as potentially misassembled by this approach (left) vs. a randomly selected set of windows for comparison (right).

Table 1. *fragScaff* performance summary**Table 1.** *fragScaff* performance summary

Organism	Input Assembly ^a	Input N50 (kbp)	Input Scaffold count	Library Method	<i>fragScaff</i> N50 (kbp)	N50 fold Improvement	<i>fragScaff</i> Scaffold count	Percent Bases Included ^b	Bases Properly Placed ^c
Human	S	47	127,088	CPT-seq	570 (2,720)	12 (57)	39,377 (24,251)	95.8 (97.8)	99.7 (99.2)
Human	S + 3Kb	437	18,921	CPT-seq	4,398 (18,193)	10 (42)	7,596 (5,514)	97.1 (99.0)	98.5 (97.8)
Human	S + 3Kb	437	18,921	Fosmid	567	1.3	15,303	38.9	88.0
Human	S + 3Kb	437	18,921	LFR	668	1.5	14,476	46.6	60.5
Human	R, 15Kb	15	191,312	CPT-seq	361	24	36,790	77.9 ^d	99.5
Human	R, 100Kb	100	28,817	CPT-seq	6,601	66	4,223	90.5 ^d	99.3
Mouse	S + 3Kb	224	25,964	CPT-seq	2,916	13	3,969	96.2	98.1
Fly	S	68	7,109	CPT-seq	524	7.7	3,779	87.3	99.7

^aInput Assembly. S=shotgun, 3kbp=3 kbp mate-pair, R=split reference with size of the fragments

^bPercentage of the sum length of input contigs included in the *fragScaff* assembled scaffolds

^cPercentage of bases in *fragScaff* assembly that are within scaffolds that appear to be correctly assembled

^dFor simulated assemblies the top and bottom 5% of end-nodes that are removed have a higher impact on reducing the percentage of bases included due to each contig having an equal amount of sequence, whereas in non-simulated assemblies the removed end-node contigs tend to be very small and generally sum to less than 2% of the sum contig length

Parentheses indicate values for runs of *fragScaff* with more lenient parameters

Table 2. Improvements to Hi-C based scaffolding with *fragScaff***Table 2.** Improvements to Hi-C based scaffolding with *fragScaff*

Organism	Input Assembly ^a	Input N50 (kbp)	Clustered ^b (Hi-C Only)	Ordered ^c (Hi-C Only)	Clustered ^d (<i>fragScaff</i> + Hi-C)	Ordered ^e (<i>fragScaff</i> + Hi-C)
Human	S	47	89.7	41.5	99.4	99.1
Human	S + 3kbp	437	98.2	94.4	98.8	96.0
Human	R, 15kbp	15	35.9	0.2	91.9	88.2
Human	R, 25kbp	25	28.9	0.4	92.5	93.1
Human	R, 50kbp	50	70.8	16.5	93.6	95.5
Mouse	S + 3kbp	224	98.0	86.7	99.8	98.6
Fly	S	68	81.2	82.0	96.2	93.0

^aInput Assembly. S=shotgun, 3kbp=3 kbp mate-pair, R=split reference with size of the fragments

^bPercent of sequence clustered into chromosome groups using Hi-C data only for *LACHESIS* based scaffolding

^cPercent of clustered sequence that can be ordered using Hi-C data only for *LACHESIS* based scaffolding

^dPercent of sequence clustered into chromosome groups, scaffolding with *fragScaff* prior to using Hi-C data

^ePercent of clustered sequence that can be ordered, scaffolding with *fragScaff* prior to using Hi-C data

References

- Adey A, Burton JN, Kitzman JO, Hiatt JB, Lewis AP, Martin BK, Qiu R, Lee C, Shendure J. 2013. The haplotype-resolved genome and epigenome of the aneuploid HeLa cancer cell line. *Nature* 500: 207-11.
- Adey A, Morrison HG, Asan, Xun X, Kitzman JO, Turner EH, Stackhouse B, MacKenzie AP, Caruccio NC, Zhang X, Shendure J. 2010. Rapid, low-input, low-bias construction of shotgun fragment libraries by high-density in vitro transposition. *Genome Biology* 11: R119.
- Adey A, Shendure J. 2012. Ultra-low-input, tagmentation-based whole-genome bisulfite sequencing. *Genome research* 22: 1139-43.
- Amini S, Pushkarev D, Christiansen L, Royce T, Turk C, Pignatelli N, Adey A, Kitzman JO, Ronaghi M, Shendure J, et al. Haplotype-resolved Whole Genome Sequencing using a Contiguity Preserving Transposition Assay. Under Review.
- Blanco-Ulate B, Rolshausen PE, Cantu D. 2013. Draft Genome Sequence of the Grapevine Dieback Fungus *Eutypa lata* UCR-EL1. *Genome announcements* 1.
- Bradnam KR, Fass JN, Alexandrov A, Baranay P, Bechner M, Birol I, Boisvert S, Chapman JA, Chapuis G, Chikhi R, et al. 2013. Assemblathon 2: evaluating de novo methods of genome assembly in three vertebrate species. *GigaScience* 2: 10.
- Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. 2013. Transposition of native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins and nucleosome position. *Nature methods* 10: 1213-8.
- Burton JN, Adey A, Patwardhan RP, Qiu R, Kitzman JO, Shendure J. 2013. Chromosome-scale scaffolding of de novo genome assemblies based on chromatin interactions. *Nature biotechnology* 13: 1119-25.
- Burton JN, Liachko I, Dunham MJ, Shendure J. 2014. Species-Level Deconvolution of Metagenome Assemblies with Hi-C-Based Contact Probability Maps. *G3 (Bethesda)*. 2014 May 22. pii: g3.114.011825. doi: 10.1534/g3.114.011825.
- Donmez N, Brudno M. 2013. SCARPA: scaffolding reads with practical algorithms. *Bioinformatics (Oxford, England)* 29: 428-34.
- Duitama J, Gayle M, Eun-Kyung S, Hoehe MR. 2010. ReFHap: A Reliable and Fast Algorithm for Single Individual Haplotyping. 2010. <http://dna.engr.uconn.edu/bibtexmgr/upload/Dal.10.pdf>
- Earl D, Bradnam K, John SJ, Darling A, Lin D, Fass J, Yu HO, Buffalo V, Zerbino DR, Diekhans M, et al. 2011. Assemblathon 1: a competitive assessment of de novo short read assembly methods. *Genome research* 21: 2224-41.

Erlich Y, Chang K, Gordon A, Ronen R, Navon O, Rooks M, Hannon GJ. 2009. DNA Sudoku—harnessing high-throughput sequencing for multiplexed specimen analysis. *Genome Research* 19: 1243-53.

Genovese G, Handsaker RE, Li H, Kenny EE, McCarroll SA. 2013. Mapping the human reference genome's missing sequence by three-way admixture in Latino genomes. *American Journal of Human Genetics* 93: 411-21.

Gertz J, Varley KE, Davis NS, Baas BJ, Goryshin IY, Vaidyanathan R, Kuersten S, Myers RM. 2012. Transposase mediated construction of RNA-seq libraries. *Genome research* 22: 134-41.

Gnerre S, Maccallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, et al. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proceedings of the National Academy of Sciences of the United States of America* 108: 1513-8.

Goryshin IY and Reznikoff WS. 1997. Tn5 in vitro transposition. *The Journal of Biological Chemistry*, 273, 7367-7374.

Gritsenko AA, Nijkamp JF, Reinders MJ, de Ridder D. 2012. GRASS: a generic algorithm for scaffolding next-generation sequencing assemblies. *Bioinformatics (Oxford, England)* 28: 1429-37.

Hunt M, Newbold C, Berriman M, Otto TD. 2014. A comprehensive evaluation of assembly scaffolding tools. *Genome biology* 15: R42.

The International Human Genome Consortium. 2001. Initial sequencing and analysis of the human genome. *Nature* 409: 860-921.

The International Human Genome Consortium. 2004. Finishing the euchromatic sequence of the human genome. *Nature* 431: 931-45.

Jia J, Zhao S, Kong X, Li Y, Zhao G, He W, Appels R, Pfeifer M, Tao Y, Zhang X, et al. 2013. *Aegilops tauschii* draft genome sequence reveals a gene repertoire for wheat adaptation. *Nature* 496: 91-5.

Jurka J, Kapitonov VV, Pavlicek A, Klonowski P, Kohany O, Walichiewicz J. 2005. Repbase Update, a database of eukaryotic repetitive elements. *Cytogenetic and genome research* 110: 462-7.

Kaper F, Swamy S, Klotzle B, Munchel S, Cottrell J, Bibikova M, Chuang H, Kruglyak S, Ronaghi M, Eberle MA, et al. 2013. Whole-genome haplotyping by dilution, amplification, and sequencing. *Proceedings of the National Academy of Sciences of the United States of America* 110: 5552-7.

Kidd JM, Sampas N, Antonacci F, Graves T, Fulton R, Hayden HS, Alkan C, Malig M, Ventura M, Giannuzzi G, et al. 2010. Characterization of missing human genome sequences and copy-number polymorphic insertions. *Nature methods* 7: 365-71.

Kitzman JO, Mackenzie AP, Adey A, Hiatt JB, Patwardhan RP, Sudmant PH, Ng SB, Alkan C, Qiu R, Eichler EE, et al. 2011. Haplotype-resolved genome sequencing of a Gujarati Indian individual. *Nature biotechnology* 29: 59-63.

Koren S, Schatz MC, Walenz BP, Martin J, Howard JT, Ganapathy G, Wang Z, Rasko DA, McCombie WR, Jarvis ED, et al. 2012. Hybrid error correction and de novo assembly of single-molecule sequencing reads. *Nature biotechnology* 30: 693-700.

Koren S, Treangen TJ, Pop M. 2011. Bambus 2: scaffolding metagenomes. *Bioinformatics (Oxford, England)* 27: 2964-71.

Laszlo AH, Derrington IM, Ross BC, Brinkerhoff H, Adey A, Nova IC, Craig JM, Langford KW, Samson JM, Daza R, Doering K, et al. 2014. Decoding long nanopore sequencing reads of natural DNA. *Nature biotechnology*. doi: 10.1038/nbt.2950.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R and 1000 Genome Project Data Processing Subgroup. 2009. The Sequence alignment/map (SAM) format and SAMtools. *Bioinformatics*, 25, 2078-9.

Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics (Oxford, England)* 26: 589-95.

Lo C, Liu R, Lee J, Robasky K, Byrne S, Lucchesi C, Aach J, Church G, Bafna V, Zhang K. 2013. On the design of clone-based haplotyping. *Genome biology*. 14(9):R100.

Luo R, Liu B, Xie Y, Li Z, Huang W, Yuan J, He G, Chen Y, Pan Q, Liu Y, et al. 2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *GigaScience* 1: 18.

Pagani I, Liolios K, Jansson J, Chen IA, Smirnova T, Nosrat B, Markowitz VM, Kyrpides NC. 2012. The Genomes OnLine Database (GOLD) v.4: status of genomic and metagenomic projects and their associated metadata. *Nucleic acids research* 40: D571-9.

Rong X, Gardener BB. 2013. Draft Genome Sequence of *Cryptococcus flavescens* Strain OH182.9_3C, a Biocontrol Agent against *Fusarium* Head Blight of Wheat. *Genome announcements* 1.

Schwartz JJ, Lee C, Hiatt JB, Adey A, Shendure J. 2012. Capturing native long-range contiguity by in situ library construction and optical sequencing. *Proceedings of the National Academy of Sciences of the United States of America* 109: 18749-54.

Shemesh M, Pasvolsky R, Sela N, Green SJ, Zakin V. 2013. Draft Genome Sequence of *Alicyclobacillus acidoterrestris* Strain ATCC 49025. *Genome announcements* 1.

Voskoboynik A, Neff NF, Sahoo D, Newman AM, Pushkarev D, Koh W, Passarelli B, Fan HC, Mantalas GL, Palmeri KJ, et al. 2013. *Elife* doi: 10.7554/eLife.00569.

Wang D, Wu R, Xu Y, Li M. 2013. Draft Genome Sequence of *Rhizopus chinensis* CCTCCM201021, Used for Brewing Traditional Chinese Alcoholic Beverages. *Genome announcements* 1: e0019512.

Zhang G, Fang X, Guo X, Li L, Luo R, Xu F, Yang P, Zhang L, Wang X, Qi H, et al. 2012. The oyster genome reveals stress adaptation and complexity of shell formation. *Nature* 490: 49-54.

Zook J, Salit M. 2013. NIST NA12878 High-Quality Variant Call Set. (ftp://ftp-trace.ncbi.nih.gov/giab/ftp/data/NA12878/variant_calls/NIST)

