




Inhibition of transcription leads to rewiring of locus-specific chromatin proteomes

Deepani W Poramba-Liyanage, Tessy Korthout, Christine E Cucinotta, et al.

Genome Res. published online March 18, 2020

Access the most recent version at doi:[10.1101/gr.256255.119](https://doi.org/10.1101/gr.256255.119)

| | |
|---------------------------------|--|
| P<P | Published online March 18, 2020 in advance of the print journal. |
| Accepted Manuscript | Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version. |
| Open Access | Freely available online through the <i>Genome Research</i> Open Access option. |
| Creative Commons License | This manuscript is Open Access. This article, published in <i>Genome Research</i> , is available under a Creative Commons License (Attribution 4.0 International license), as described at http://creativecommons.org/licenses/by/4.0/ . |
| Email Alerting Service | Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here . |



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Published by Cold Spring Harbor Laboratory Press

Inhibition of transcription leads to rewiring of locus-specific chromatin proteomes

Deepani W. Poramba-Liyanage^{1,*}, Tessy Korthout^{1,*}, Christine E. Cucinotta², Ila van Kruijsbergen¹, Tibor van Welsem¹, Dris El Atmioui³, Huib Ovaa³, and Toshio Tsukiyama², Fred van Leeuwen^{1,4,#}

¹Division of Gene Regulation, Netherlands Cancer Institute, 1066CX Amsterdam, The Netherlands

²Basic Sciences Division, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA

³Leiden Institute for Chemical Immunology, Leiden University Medical Center, 2333ZC Leiden, The Netherlands, and OncoCode Institute.

⁴Department of Medical Biology, Amsterdam UMC, University of Amsterdam, 1105 AZ Amsterdam, The Netherlands.

*These authors contributed equally

Running title:

Decoding dynamic transcription proteomes

Key Words:

Epi-Decoder, chromatin, transcription, 1,10-phenanthroline, proteomics, quiescence

Corresponding author

Fred.v.leeuwen@nki.nl; Division of Gene Regulation, Netherlands Cancer Institute, Plesmanlaan 121, 1066CX Amsterdam, The Netherlands. +31-20-5121973

Abstract

Transcription of a chromatin template involves the concerted interaction of many different proteins and protein complexes. Analyses of specific factors showed that these interactions change during stress and upon developmental switches. However, how the binding of multiple factors at any given locus is coordinated has been technically challenging to investigate. Here we employed Epi-Decoder in yeast to systematically decode, at one transcribed locus, the chromatin binding changes of hundreds of proteins in parallel upon perturbation of transcription. Taking advantage of improved Epi-Decoder libraries, we observed broad rewiring of local chromatin proteomes following chemical inhibition of RNA polymerase. Rapid reduction of RNA polymerase II binding was accompanied by reduced binding of many other core transcription proteins and gain of chromatin remodelers. In quiescent cells, where strong transcriptional repression is induced by physiological signals, eviction of the core transcriptional machinery was accompanied by the appearance of quiescent-cell specific repressors and rewiring of the interactions of protein-folding factors and metabolic enzymes. These results show that Epi-Decoder provides a powerful strategy for capturing the temporal binding dynamics of multiple chromatin proteins under varying conditions and cell states. The systematic and comprehensive delineation of dynamic local chromatin proteomes will greatly aid in uncovering protein-protein relationships and protein functions at the chromatin template.

Introduction

Many proteins assemble on to DNA to implement gene regulatory programs and ensure the expression of a subset of genes in agreement with the state and environmental cues of the cell. RNA polymerase II (Pol II) is an integral part of this assembly as it catalyzes the DNA dependent synthesis of messenger RNA (mRNA). Regulation of Pol II occurs at different stages of transcription and involves concerted actions of many proteins and protein complexes and dynamic post-translational modifications of histones and the carboxy-terminal domain (CTD) of Pol II (Sainsbury et al. 2015; Chen et al. 2018; Li et al. 2018).

The process of transcription involves distinct stages: initiation, elongation and termination (Sainsbury et al. 2015; Chen et al. 2018; Li et al. 2018). The wrapping of DNA by histones into nucleosomes presents a barrier for each of these steps. Chromatin remodelers recruited to promoters can displace nucleosomes to open up crucial recognition elements for transcription factor (TF) binding (Struhl and Segal 2013; Prasad et al. 2016). For initiation of transcription, Pol II assembles with basal TFs (TFIIB-H) at the promoter to form the pre-initiation complex, which opens up the DNA, initiates RNA synthesis, and stimulates the escape of Pol II from the promoter (Sainsbury et al. 2015; Feng et al. 2016; Hantsche and Cramer 2017). Productive elongation involves recruitment of Paf1C (Polymerase II Associated Factor complex), DSIF (DRB Sensitivity Inducing Factor; Spt4/5), Spt6 and FACT (Facilitator of Chromatin Transcription) (Xu et al. 2017; Ehara and Sekine 2018; Vos et al. 2018). Towards the end of genes transcription is terminated and the RNA is processed by the recruitment of cleavage and polyadenylation factors, which results in the release of Pol II and the nascent mRNA from the DNA template (Porrua and Libri 2015). Biochemical and genetic studies have provided a rich catalog of factors involved in the different stages of transcription. However, how the many different interactions are

coordinated at any given chromatin locus in time and under changing conditions is still poorly understood (Qiu and Kaplan 2019).

Unraveling the relationships between proteins and the hierarchies among them at the chromatin template will require measuring the chromatin interactome and monitoring the changes upon perturbation. A common and convenient strategy to perturb transcription is the use of inhibitors of Pol II (Bensaude 2011). Some of the commonly used inhibitors of transcription initiation or elongation are not active in live yeast cells because of uptake deficiency or because the target protein or target site on the protein is not conserved (Bensaude 2011). However, the addition of the metal chelators thiolutin and 1,10-phenanthroline (PH) leads to rapid loss of transcription in yeast. This has, for example, enabled the determination of mRNA half-lives and establishing the role of Pol II activity in nucleosome positioning (Adams and Gross 1991; Grigull et al. 2004; Sun et al. 2013; Hsieh et al. 2015; Vasseur et al. 2016; Lauinger et al. 2017). In addition, 6-Azauracil (AU), which perturbs the supply of ribonucleotides and thereby indirectly affects RNA polymerases, is frequently used to assess transcription-elongation phenotypes in yeast (Handschumacher and Welch 1956; Exinger and Lacroute 1992; Shaw and Reines 2000; Mason and Struhl 2005; Zhou et al. 2015). Although these chemicals and related compounds have been used extensively and are convenient and powerful tools to block transcription, little is known about how the direct and indirect inhibition of transcription affects the interactions of Pol II and its protein partners with the chromatin template.

Transcription is also regulated under physiological conditions during development and differentiation. In budding yeast, a global change in transcription is observed when cells enter quiescence, a survival mode that involves a G1-like cell-cycle arrest, with increased resistance to stress, metabolic rewiring, and large-scale reorganization of the genome and cellular machineries (Radonjic et al. 2005; Aragon et al.

2008; Broach 2012; Miles et al. 2013; Kuang et al. 2014; Mews et al. 2014; Guidi et al. 2015; McKnight et al. 2015; Rutledge et al. 2015; Young et al. 2017; Sagot and Laporte 2019b; Sagot and Laporte 2019a; Swygert and Tsukiyama 2019). Quiescence is central to many important biological processes and is conserved from unicellular eukaryotes to multicellular organisms (Cheung and Rando 2013; Sagot and Laporte 2019a). We recently demonstrated that quiescent cells show a 30-fold drop in mRNA levels and that this massive transcriptional shutoff is dependent on the conserved histone deacetylase (HDAC) Rpd3 (McKnight et al. 2015). Rpd3 is recruited to the majority of gene promoters in quiescent cells, leading to global hypoacetylation of chromatin and gene repression (McKnight et al. 2015). Further, transcriptional shutdown correlates with loss of Rpb3, one of the subunits of Pol II (McKnight et al. 2015; Young et al. 2017).

To systematically interrogate the chromatin changes at a transcriptionally active locus upon inhibition of transcription, we employed Epi-Decoder, a Tag-chromatin immunoprecipitation-Barcode-Sequencing technology in budding yeast (Korthout et al. 2018). Epi-Decoder enables the decoding of the proteome of a single barcoded genomic locus by DNA sequencing and barcode counting. It takes advantage of the power of cellular DNA barcoding (Yan et al. 2008; Fowler and Fields 2014; Vlaming et al. 2016; Chabbert et al. 2018; Keschull and Zador 2018; Roy et al. 2018) and yeast genetics (Duina et al. 2014) and provides a quantitative approach orthogonal to capture-mass-spectrometry efforts (Schmidtman et al. 2016; Wierer and Mann 2016; Myers et al. 2018). Here we describe an expanded library of yeast strains carrying a double-barcoded transcribed reporter gene integrated at the HO locus (Epi-Decoder-HO). Of this library, we used a dedicated subset of ~700 (putative) chromatin proteins (for more details see Supplemental Materials and Methods) of which the binding at the reporter locus can be assessed in parallel and in triplicate with three independent barcodes in a single sample. This Chrom-3×BC library

was used to determine the local chromatin-proteome rewiring at the barcoded transcribed locus in response to transcription inhibition by chemical perturbation and during quiescence.

Results

Generation of improved Epi-Decoder-HO libraries

To measure local proteome dynamics at a promoter and terminator region upon transcriptional inhibition, we first improved the previously reported Epi-Decoder-HO library to optimize the comparison of multiple time points and conditions. Epi-Decoder is a strategy for decoding the local proteome of a single genomic locus (Korthout et al. 2018). It relies on short (~16-20bp) DNA barcodes (BCs) integrated at a common locus in the genome. Here we used the constitutively expressed KanMX marker gene integrated at the HO locus and flanked by two barcodes BC_UP (promoter region) and BC_DN (terminator region) that are 1.5 kb apart (HO-Barcoders). The barcoded KanMX cassette is a Kanamycin gene controlled by the heterologous *AgTEF1* promoter and terminator from *Ashbya gossypii*, a yeast related to *S. cerevisiae*. This 1.5 kb reporter-gene cassette replaces the coding sequence of the *HO* gene and is therefore flanked by the endogenous *HO* promoter and terminator sequences as well as an origin of replication (ARS404) 53 bp downstream of BC_DN. This reporter locus has previously been used in screens for various chromatin- and transcription-related features (Verzijlbergen et al. 2011; Chen et al. 2013; Vlaming et al. 2016; Korthout et al. 2018; Vlaming et al. 2019). The library of HO-Barcoders was combined with a genome-wide library of proteins tagged with a common epitope tag, a Tandem Affinity Purification (TAP) tag (Figs. 1A, and S1A). Upon pooling, cross-linking, chromatin immunoprecipitation (ChIP), amplification of the barcoded regions, and counting the barcodes by massive parallel sequencing (Fig 1B), the abundance of each BC (ChIP/input) reports on the occupancy of each tagged protein at its barcoded locus (Verzijlbergen et al. 2011; Vlaming et al. 2016; Korthout et al. 2018). Integral for high-throughput assessments like this is the ability to multiplex many individual TAP-tag clones (~4800 in total) in the ChIP assays using unique BCs and indexing strategies. Here we expanded the previous set of HO-Barcoders from ~1100 to ~2500 (Douglas et al. 2012), enabling coverage of the full Epi-Decoder

library with only two separate subsets (Fig 1B and Supplemental Fig S1A). The expanded Barcoder-HO library was combined with the TAP-tag library in three different ways (see Methods) such that every TAP-tag is linked to three independent HO-Barcoders (I, II and III) and creating a triplicate ~700 Chromatin TAP-tag subset (Chrom-3×BC) that can be processed as one pool due to non-overlapping barcode pairs in the replicates.

Interrogation of protein binding with multiple DNA barcodes

Epi-Decoder uses DNA BC counting as a reporter for protein binding (Korthout et al. 2018). With the three independent Epi-Decoder-HO libraries (Fig 1B), we first assessed to what extent variation in quantification of protein binding is caused by the barcode sequences. The protein-binding patterns in each of the three Epi-Decoder-HO libraries strongly correlated between the replicates (Fig 1C) for both BC_UP and BC_DN (Supplemental Fig S1B, S1C and Supplemental Table S1) and this was further confirmed by the analysis of individual clones (see below). Although it is possible that specific barcode sequences could cause a bias for detection of certain proteins, in general our results show that the inferred protein binding scores are largely independent of the short barcode sequences. The use of multiple Tag-Barcode combinations further increases the confidence of single-barcode measurements. For example, we have previously shown that Ssl2 and Tfa2, two factors known for their roles in transcription initiation, also show high binding at the HO terminator region (Korthout et al. 2018). By replicating this finding with three independent barcodes, we can now negate the possibility that BC effects caused binding of Ssl2 and Tfa2 at the HO terminator region (Supplemental Table S1). The consistent and robust binding of Ssl2 and Tfa2 at the terminator region in the absence of other basal transcription factors suggests a non-canonical function of these factors.

A Chromatin-TAP-tag subset library for capturing dynamic local chromatin proteomes

Having confirmed that Epi-Decoder provides a robust assay to quantitatively measure protein-binding at a single locus of thousands of proteins in parallel, we focused on the Chromatin TAP-tag subset (Chrom-3×BC; Fig 2A and see Methods) to determine the dynamics of the local chromatin proteome following chemical inhibition of Pol II. The antifungal agent PH is a metal ion chelator that sequesters Zn^{2+} ions, which are essential for the activity of RNA polymerases (Scrutton et al. 1971; Markov et al. 1999; Cramer et al. 2000; McCann et al. 2012). Addition of PH to cells rapidly affects mRNA levels (Grigull et al. 2004; Miller et al. 2011; Wada and Becskei 2017). AU is an inhibitor of IMP dehydrogenase (IMPDH), the rate-limiting enzyme in *de novo* GTP synthesis (Grigull et al. 2004; Kaplan et al. 2012; Ljungdahl and Daignan-Fornier 2012). Treatment of cells with AU, results in depletion of intracellular nucleotide pools, thereby affecting transcription elongation (Handschumacher and Welch 1956; Exinger and Lacroute 1992; Shaw and Reines 2000; Shaw et al. 2001; Mason and Struhl 2005). To investigate the consequences of inhibition of Pol II at the chromatin template, the Chrom-3×BC Epi-Decoder library was incubated at with PH and AU and the local proteomes were determined at several subsequent time points (Fig 2A). All experiments were carried out at low temperature (16°C) to facilitate the capturing dynamic binding events. In addition, cells were arrested in G1 to avoid cell-cycle dependent effects (Supplemental Fig S2A) (O'Duibhir et al. 2014). Independent triplicate measurements in the Chrom-3×BC pool and sample multiplexing enabled efficient analysis of the dynamics of known chromatin proteins across multiple conditions and time points.

We confirmed that PH and AU are active under these conditions: both compounds inhibited cell growth (Supplemental Fig S2B) and addition of PH led to induction of *ZRT1* mRNA while addition of AU led to induction of *IMD2* mRNA (Fig 2B and Supplemental Fig S2C), which is in agreement with previous observations (Shaw and Reines 2000; Grigull et al. 2004) that inhibition of transcription by these drugs is not complete. Under these experimental conditions and time points, we observed little decay of mRNA

levels of the barcoded KanMX gene or other genes examined (Fig 2B and Supplemental Fig S2C), in agreement with the observed temperature effects on RNA decay kinetics (Lotan et al. 2005). In contrast, global inspection of binders at the barcoded KanMX gene showed that the addition of PH or AU led to many rapid and pronounced changes at the chromatin level at BC_UP and BC_DN (Fig 2C). Generally, changes in protein binding were stronger in PH than in AU, confirming that the observed effects are specific for the treatment.

Rewiring of the core transcription machinery at chromatin upon inhibition of Pol II

The global overview of the dynamic chromatin proteomes (average of three replicates, Fig 2C) revealed broad rearrangements, especially of protein complexes related to transcription. Other proteins were unaffected, such as the replication proteins specifically associated with BC_DN proximal to the origin of replication ARS404. This was further illustrated by the analysis of individual replicate measurements of proteins representing different aspects of chromatin biology (Fig 3A). Because these three replicates, were based on different barcode pairs, the reproducible results indicate that barcode sequences did not generally affect the protein-binding measurements in Epi-Decoder. As observed previously (Korthout et al. 2018), Pol II and transcription elongation factors were found at BC_UP and were more abundant at BC_DN, initiation factors and basal transcription factors (TFs) were more enriched at BC_UP, while transcription termination factors and replication proteins were specifically bound to BC_DN (Figs 2C, 3A, and Supplemental Table S2). To determine the consequences of transcription inhibition for the transcription machinery in more detail, we focused on Pol II and transcription elongation factors. All the Pol II subunits present in our Epi-Decoder library showed a pronounced and progressive loss of binding upon addition of PH at BC_UP and BC-DN. In AU, Pol II binding was also reduced but to a lesser extent (Fig 3B). Therefore, we here focused on the changes following treatment with PH. Loss of binding can be caused by lower levels of the protein due to protein degradation or by redistribution of the protein. To

distinguish between these two possibilities, we determined the protein level of Rpo21, the largest subunit of Pol II, by immunoblot analysis and found that treatment with PH did not lead to reduced Rpo21 protein levels (Fig 3C and Supplemental Fig S3A) while treatment with AU showed a modest decrease (Fig 3D and Supplemental Fig S3B). Therefore, the reduced binding of Pol II subunits in PH was not accompanied by increased Pol II protein degradation.

Finally, the strong reduction in Pol II binding observed in Epi-Decoder in PH at the barcoded KanMX gene under the control of the *AgTEF1* promoter and terminator could be validated by ChIP-qPCR at the 5' and 3' end of the endogenous *TEF1* gene (Figs 3E and Supplemental Fig S3C). Thus, in addition to the current knowledge that Pol II activity is inhibited directly by PH and indirectly by AU (Scrutton et al. 1971; Markov et al. 1999; Cramer et al. 2000; McCann et al. 2012), our results show that chemical inhibition of Pol II also leads to partial eviction of the core Pol II machinery from the chromatin template.

Epi-Decoder uncovers protein-protein relationships

Several proteins and protein complexes have been biochemically or genetically linked to the process of transcription elongation (Koerber et al. 2009; Venters and Pugh 2009; Kwak and Lis 2013; Pufall and Kaplan 2013; Cucinotta and Arndt 2016; Van Oss et al. 2017; Chen et al. 2018; Vinayachandran et al. 2018). However, how the interactions between the core transcriptional machinery and elongation factors are dynamically orchestrated on the chromatin template is still incompletely understood (Mayer et al. 2010; Vinayachandran et al. 2018). Inspection of the dynamic proteomes at the HO locus showed that elongation factors did not all respond equally to inhibition of Pol II by PH (Fig 4A) and AU (Supplemental Fig S4A). The conserved elongation factors Elf1, Spn1, Spt6, and DSIF (Spt4/5) were rapidly and progressively evicted from chromatin, closely resembling the dynamics of Pol II upon PH treatment (Figs 4A and 3B). This indicates that most elongation factors depend on Pol II for recruitment and maintenance. The treatment with AU, performed and processed in parallel, showed more modest

effects on binding of Pol II and the elongation factors (Supplemental Fig S4A-B). The FACT complex, composed of Spt16 and Pob3, is known to be recruited to chromatin by the act of transcription and its associating factors Nhp6a and Nhp6b, two High Mobility Group (HMG) proteins (Ruone et al. 2003; Kwak and Lis 2013; McCullough et al. 2015; Martin et al. 2018; Pathak et al. 2018).

At BC_UP, FACT eviction was delayed compared to eviction of Pol II upon PH treatment (Figs 4A-B), suggesting that FACT binding might depend on more stable factors at transcribed genes such as histones or histone modifications (Martin et al. 2018; Vinayachandran et al. 2018). At BC_DN, the two core FACT subunits Spt16 and Pob3 were rapidly lost upon addition of PH, closely resembling the behavior of Pol II and other elongation factors. In contrast, the HMG proteins Nhp6a and Nhp6b did not show reduced binding after Pol II inhibition (Figs 4A-B). This difference may reflect the ability of HMG proteins to bind DNA and nucleosomes directly, independently of other chromatin proteins.

Recent structural studies show that Paf1C, DSIF (Spt4/5), and Spt6 can form an intricate protein network around elongation-permissive Pol II (Vos et al. 2018; Xie et al. 2018). Indeed, like DSIF and Spt6, most Paf1C subunits showed reduced binding to chromatin when Pol II was inhibited and evicted by PH and AU (Fig 4A-B and Supplemental Fig S4A-B). However, binding of Ctr9, the largest subunit and a key scaffold protein of Paf1C (Deng et al. 2018; Vos et al. 2018; Xie et al. 2018), was largely unaffected in PH, with a small decrease only observed at later time points (Figs 4A-B). We confirmed the delayed loss of Ctr9 by ChIP-qPCR analysis of the barcoded promoter and terminator regions (Fig 4C) as well as the endogenous *TEF1* locus (Supplemental Fig S4C). The differences in kinetics were not caused by differences in protein abundance, because Ctr9 showed very similar global expression levels as Paf1 and Pol II subunits (Supplemental Fig S4D). We also tested whether the delayed loss of Ctr9 was caused by a difference in crosslinking sensitivity (Zaidi et al. 2017; De Jonge et al. 2019). However, Paf1 and Rpo21

were not more sensitive to shorter cross-linking conditions than Ctr9 (Supplemental Fig S4E). Finally, addition of PH did not lead to major changes in the expression level of Rpo21 and Paf1, compared to Ctr9, the latter remaining bound when Rpo21 and Paf1 had already been evicted upon PH treatment (Fig 4D). Together, our results show that binding of Ctr9 at transcribed chromatin is independent of Pol II and Paf1C subunits, suggesting that Ctr9 has a transcription elongation-independent mechanism to bind to DNA, perhaps mediated by its known interactions with DNA and nucleosomes (Musso et al. 2000; Deng et al. 2018; Vos et al. 2018). These binding activities may be responsible for anchoring Ctr9 to PH-perturbed chromatin in the absence of its complex partners and Pol II. Of note, under conditions of transcriptional repression in quiescent cells, Ctr9 followed the pattern of the other Paf1C members (see below), showing that its dynamics are context dependent.

In addition to uncovering protein-binding dynamics within well-known protein complexes, the parallel analysis of hundreds of proteins also offers the possibility to obtain more insight into proteins of which the chromatin functions have been less well annotated. Here we more closely examined Srm1 (also known as Prp20), a homologue of human Regulator of Chromosome Condensation 1 (RCC1). Srm1/RCC1 is a guanine nucleotide exchange factor that localizes to the nucleus, binds nucleosomes in a dynamic way, and occupies a large part of the yeast genome (Nemergut et al. 2001; Koerber et al. 2009; Makde et al. 2010; Wu et al. 2011; Bierbaum and Bastiaens 2013; McGinty and Tan 2016). In the Epi-Decoder analysis, all proteins are examined in one pool and with the same tag, thereby delivering qualitative as well as relative quantitative information. The high barcode counts observed for Srm1 show that it binds efficiently to chromatin, in the same range as the abundant transcription elongation factors mentioned above (Fig 4A and Supplemental Fig S4A). Analysis of Srm1 by ChIP-qPCR confirmed the efficient binding of this protein and an increase after PH treatment at the reporter locus as well as endogenous loci, including a non-transcribed region (Fig 4E). This, together with the genome-wide binding pattern

(Koerber et al. 2009), indicates that Srm1 should be considered as a common component of yeast chromosomes. However, the distribution and kinetic behavior of Srm1 did not mirror that of the transcription elongation factors: upon treatment with PH, Srm1 binding increased (Fig 4A,E) while binding of Pol II and most elongation proteins decreased. Although not quantitatively the same, the behavior of Srm1 is more similar to that of the chromatin remodelers INO80 and RSC, several metabolic enzymes (Fig 2C), the heat shock protein Ssa2 (Fig 3A and Supplemental Fig S4F) and high-mobility-group protein Hmo1 (Supplemental Fig S4F). In contrast, the high-mobility group-like protein Spt2, a chaperone involved in histone recycling over transcribed regions of active genes (Nourani et al. 2006; Chen et al. 2015), was decreased in active regions (Supplemental Fig S4F). The dynamic behavior of Srm1, Ssa2, Hmo1, and Spt2 after pH treatment was not caused by changes in global protein levels (Supplemental Fig S4G). Together, these results suggest that Srm1 is an abundant chromatin protein with the potential to affect chromatin structure and function but it that it might not act as a canonical transcription elongation factor.

Rearrangement of the chromatin proteome of the barcoded HO locus upon entry in quiescence

Finally, we investigated how the chromatin proteome of the barcoded HO locus was altered by the strong physiological transcriptional shut down during quiescence. Chrom-3×BC Epi-Decoder library pools were grown in YEPD at 30°C for seven days, which causes cells to arrest in saturation with G1 DNA content. Stationary phase cultures consist of two populations: quiescent (Q) cells, which are more uniform, long-lived, and stress-resistant, and nonquiescent (NQ) cells, which are more heterogeneous, short-lived, and stress-sensitive (Allen et al. 2006; Aragon et al. 2008; Li et al. 2013; Young et al. 2017; Sagot and Laporte 2019a; Sagot and Laporte 2019b). NQ cells were also isolated and processed as a reference.

Analysis of the RNA Pol II machinery, transcription elongation factors, basal transcription factors, and transcription termination and RNA processing factors showed a pronounced loss of the core transcription machinery in Q and NQ cells (Fig 5A-B, Supplemental Fig S5A-E; Supplemental Table S2). Overall, the magnitude of the changes was higher than that observed in the G1 cells treated with PH with several exceptions. The FACT-associating factors Nhp6a and Nhp6a showed increased chromatin binding in Q cells (Figs 5B and Supplemental Fig S5A), in contrast to PH treated cells (Figs 4 and Supplemental Fig S4). Tho1, a protein associated with transcribed chromatin, showed increased abundance in Q and NQ cells but not in PH (Figs 5B and Supplemental Fig S5A), suggesting that Tho1 might have Q/NQ-cell specific functions and that its recruitment is independent of ongoing transcription, in contrast to canonical transcription elongation proteins.

To gain more insight into the transcriptional repression of the barcoded HO locus, we inspected transcriptional regulators at the BC_UP promoter region (Figs 5C and Supplemental Fig S5B). Q cells showed binding of the quiescence-specific transcription repressor Xbp1, which was absent from mid-log cells and during PH treatment. This protein is known to recruit the histone deacetylase Rpd3, mediating the global shutdown of gene expression in Q cells (McKnight et al. 2015). Indeed, Rpd3 and members of the Rpd3L complex were mostly undetectable in mid-log but readily detected in Q cells. Members unique for the Rpd3S complex were not detected (Supplemental Table S3). This well-established repression pathway was accompanied by increased abundance of several TFs such as Mcm1, Mbf1, Tbf1, Abf1, and Swi6 (Fig 5C and Supplemental Fig S5B).

Besides the core transcriptional machinery, TFs, and Rpd3L, we observed several other changes in the chromatin proteomes (Figs 5C-E and Supplemental Fig S5B-G). Yeast Histone H1 (Hho1) showed increased abundance in Q and NQ cells (Figs 5C and Supplemental Fig S5B). High-mobility group protein 1 (Hmo1), which has been shown to function as a linker histone and promote chromatin compaction

(Panday and Grove 2016) showed the same trend (Figs 5C and Supplemental Fig S5B). In contrast, the negative regulator of transcription Spt2 left the chromatin in Q and NQ cells, as it did in PH treated cells. These changes were paralleled by the recruitment of the ATP-dependent chromatin remodeling complexes INO80 and RSC, as well as several heat shock proteins and other factors involved in protein folding (Figs 5E and Supplemental Fig S5C and S5F). In addition, changes were seen in the binding of metabolic enzymes (Supplemental Fig S5G). Future functional studies will be required to determine the role of these dynamic interactions in the global transcriptional repression occurring in Q cells.

Finally, quiescence also affected the organization of replication proteins at the origin of replication (ARS404) proximal to BC_DN (Fig 5D). In contrast to PH treatment, cells in quiescence showed a complete loss of the minichromosome maintenance complex (MCM), the replicative helicase. Q cells are arrested in a G1-like state but binding of MCM was maintained in cells arrested in G1 by mating pheromone (Liang and Stillman 1997) (Fig 5D). The loss of MCM was accompanied by increased abundance of the members of the origin-recognition complex (ORC) (Fig 5D). This altered balance is consistent with a competition model that has been proposed for MCM and ORC (Aparicio et al. 1997). Replication origin licensing, which involves the binding of ORC and subsequently the assembly of an inactive form of the replicative helicase MCM, occurs during late mitosis and G1 phase of the cell cycle. After formation of a replication complex, during S-phase the helicase is activated and the replication initiates (Siddiqui et al. 2013; Bell and Labib 2016). It is possible that clearing MCM from origins in Q cells is important for eliminating any inappropriate activation of replication origins during the prolonged state of arrest. The mechanism causing the absence of MCM and the role thereof in Q cell biology will require further study but it should be noted that similar dynamics have been observed in mouse and *S. pombe* (Madine et al. 2000; Sun et al. 2000; Namdar and Kearsey 2006). Our results suggest that replication origin licensing must be re-established when cells re-enter the cell cycle.

Discussion

Determining the changes to chromatin composition in response to cell signaling or stress can reveal insights into common mechanisms of gene regulation. Obtaining a comprehensive understanding of chromatin changes, although highly informative (Kim et al. 2010; Weiner et al. 2012; Vinayachandran et al. 2018), is often laborious and therefore not applicable to multiple conditions. Epi-Decoder provides a strategy for identifying and quantifying in an unbiased and systematic manner the proteome of an individual genomic locus by DNA sequencing. Here we used a dedicated comprehensive chromatin library (Chrom-3×BC) to determine the changes in local chromatin proteomes upon inhibition of Pol II by chemical means or by physiological signals during quiescence.

PH is a potent inhibitor of Pol II and has fungistatic activity against a broad range of pathogenic fungi and bacteria (McCann et al. 2012). PH and thiolutin, which acts in a similar way, are often used to determine consequences of perturbation of transcription in yeast. However, the consequences at the chromatin level have not been well characterized even though RNA processing, RNA export, and other downstream processes are linked to transcription and are influenced by chromatin modifying factors. Inhibition of Pol II with PH resulted in rapid remodeling of the local chromatin proteomes at the HO locus (Fig 2c). Pol II was evicted, probably reflecting a structural rearrangement caused by PH. This idea is in agreement with observations that mutants of Pol II with altered catalytic activity show reduced Pol II occupancy at genes (Malik et al. 2017). We did not observe changes in overall abundance of the evicted proteins Rpo21 and Paf1 and several other proteins, indicating that the factors evicted in PH were not subject to degradation. Besides inhibiting Pol II, PH - a metal chelator - is known to also affect the activity of metalloproteases, including a proteasome-associated deubiquitinating enzyme (Verma et al. 2002; McCann et al. 2012). Therefore, the stability of the proteins evicted by PH may at least in part

be explained by inhibition of proteasome-mediated degradation. The nature and scale of the changes observed after PH treatment, particularly the loss of the core transcription machinery, suggests that inhibition of Pol II can have widespread effects on many nuclear processes in the cell. This should be considered when interpreting downstream effects of chemical inhibitors of transcription, such as mRNA processing and stability.

During quiescence, global changes in histone modifications have been observed (Mews et al. 2014; Young et al. 2017) as well as global repression by Rpd3 (McKnight et al. 2015), chromatin compaction by condensin (Swygert et al. 2018), and reduced Pol II binding (Young et al. 2017). However, the extent to which the epigenome of quiescent cells is remodeled to support this important developmental state and to allow re-entry into vegetative growth remains poorly understood (Swygert and Tsukiyama 2019). Application of Epi-Decoder to the barcoded HO locus revealed that upon loss of transcription during quiescence, the chromatin proteome of a transcribed region undergoes a major rewiring that affects transcriptional proteins and replication complexes. In addition, we observed the recruitment of structural proteins, chromatin remodelers, and protein-folding machineries, as well as changes in metabolic enzymes. While many changes between mid-log and Q cells were also observed upon PH treatment, we observed Q-cell specific changes as well, including increased occupancy of other transcription-related proteins (Fig 5C), yeast histone H1 (Hho1), changes in specific metabolic enzymes (e.g. Tdh1, Arg1; Supplemental Fig S5E), and differential binding of several heat shock proteins (e.g. Ssa1 versus its paralog Ssa2; Fig 5E and Supplemental Fig 5E). We note that some of the observed dynamics in Q and NQ cells might, at least in part, be driven by altered expression of the proteins in the different cell states. Determining the functional roles of these Q- and NQ-cell specific changes will require further mechanistic studies.

While Q cells and NQ cells showed overlap in the changes compared to mid-log cells, we also observed important differences, in agreement with the different fates of the long-lived Q and short-lived NQ cells (Li et al. 2013) and the differences in global histone modifications (Young et al. 2017). NQ cells lacked efficient recruitment of Xbp1, which correlated with lower levels of Rpd3L and INO80. In addition, the loss of the MCM complex was incomplete in NQ cells. Together these changes may contribute to the poor fitness of these cells. Furthermore, the binding of Cdc48 at the BC_UP promoter region and BC_DN terminator region in NQ cells is a possible indicator of the general short-lived nature of NQ cells (Fig 5E). Cdc48, known as VCP in humans, is an essential and conserved AAA+ ATPase that functions as an unfoldase or segregase facilitating the extraction of proteins from macromolecular complexes, including chromatin, to enable subsequent degradation by the proteasome (Dantuma et al. 2014; Franz et al. 2016). The recruitment of protein quality-control factors to the chromatin agrees with recent observations that effective and diverse protein quality control mechanisms are active in the nucleus (Jones and Gardner 2016; Prasad et al. 2018; Frottin et al. 2019; Jones et al. 2019).

Our results show that Epi-Decoder provides a powerful strategy for capturing the temporal binding dynamics of chromatin proteins under varying conditions. We expect that future studies on local chromatin proteome maps under different conditions and at other genomic loci will offer powerful resources for detailed molecular and functional annotation of chromatin proteins and their interactions and relationships at the genome.

Methods

Yeast strains and libraries

Yeast strains used in this study are listed in Supplemental Table S4. Library manipulations on solid media were performed using synthetic genetic array (SGA) technology (Tong and Boone 2006) and a RoToR instrument (Singer Instruments, Watchet, UK). Yeast media were prepared as previously described (Tong and Boone 2006; Korthout et al. 2018). Details of library construction and growth conditions are provided in the Supplemental Materials and Methods.

RNA isolation and reverse transcription

RNA was isolated using the RNeasy Mini Kit (QIAGEN) using the protocol for yeast cells, with a few modifications, essentially as in (Korthout et al. 2018) and with a spike-in reference (A8545; a gift from R. Allshire) as described in the Supplemental Materials and Methods. RT-qPCR was performed with the primers described in (Supplemental Table S5). Each sample was measured in two technical duplicates and the average value was taken when combining biological replicates.

Epi-Decoder

Epi-Decoder was performed and analyzed essentially as described previously (Korthout et al. 2018). Protein-binding at BC_UP and BC_DN was analyzed separately with specific primers (Supplemental Table S5). Libraries of PCR products were mixed in an equimolar fashion and sequenced (single read, >50 bp) on a HiSeq2500/MiSeq platform (Illumina, San Diego, CA), using one or a mix of custom sequencing primers (Supplemental Table S5). Details of the Epi-Decoder protocol and analysis and the barcode counting can be found in the Supplemental Materials and Methods.

ChIP-qPCR

ChIP-qPCR experiments were performed with IgG Sepharose 6 Fast Flow beads or epoxy-activated Dynabeads as described previously (Korthout et al. 2018; Vlaming et al. 2019) and in the Supplemental Materials and Methods. Each ChIP was performed in triplicate.

Protein detection by immunoblot and antibodies

For immunoblotting, strains were grown to mid-log phase or arrested in G1 and processed as described previously (Korthout et al. 2018; Vlaming et al. 2019). Antibodies used and detailed protocols can be found in the Supplemental Materials and Methods.

Data access

All processed data are within the paper and the Supplemental Material. All raw sequencing data generated in this study and corresponding reference tables (Supplemental Tables S6 - S11) have been submitted to the NCBI BioProject database (<https://www.ncbi.nlm.nih.gov/bioproject/>) under accession number PRJNA610036, and to the NCBI BioSample database (<https://www.ncbi.nlm.nih.gov/biosample/>) under accession numbers SAMN14271070, SAMN14271071, and SAMN14271072.

Acknowledgements

We thank Brenda J Andrews and Charles Boone for kindly providing the expanded yeast Barcode Collection. The authors thank the RHPC facility of the Netherlands Cancer Institute for providing computational resources, the NKI Genome Core Facility for barcode-sequencing, Ben Morris of the NKI high-throughput center for help with rearraying yeast libraries by robotics. We thank Craig D. Kaplan, Thom Molenaar, Eliza Mari Kwesi-Maliepaard, Bas van Steensel, Tineke Lenstra, Marlize van Breugel and

Anne Kappert for insightful discussions and valuable suggestions. FvL was supported by the Dutch Research Council (NWO-VICI-016.130.627). TT was supported by the National Institutes of Health (R01 GM111428). CEC was supported by the National Cancer Institute (T32CA009657) and National Institutes of Health (F32GM131554). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

DWP-L, TK, FvL: Conception and design, Acquisition of data, Analysis and interpretation of data, Drafting and revising the article; CEC and TT: Q cell library preparation; DWP-L: Methodology and Validation; TvW: Acquisition, analysis and interpretation of data; IvK: data analysis; DEA, HO: Generation of reagents.

References

- Adams CC, Gross DS. 1991. The yeast heat shock response is induced by conversion of cells to spheroplasts and by potent transcriptional inhibitors. *J Bacteriol* **173**(23): 7429-7435.
- Allen C, Büttner S, Aragon AD, Thomas JA, Meirelles O, Jaetao JE, Benn D, Ruby SW, Veenhuis M, Madeo F et al. 2006. Isolation of quiescent and nonquiescent cells from yeast stationary-phase cultures. *J Cell Biol* **174**(1): 89-100.
- Aparicio OM, Weinstein DM, Bell SP. 1997. Components and dynamics of DNA replication complexes in *S.cerevisiae*: redistribution of MCM proteins and Cdc45p during S phase. *Cell* **91**393: 59-69.
- Aragon AD, Rodriguez AL, Meirelles O, Roy S, Davidson GS, Tapia PH, Allen C, Joe R, Benn D, Werner-Washburne M. 2008. Characterization of Differentiated Quiescent and Nonquiescent Cells in Yeast Stationary-Phase Cultures. *Mol Biol Cell* **19**: 1271-1280.
- Bell SP, Labib K. 2016. Chromosome Duplication in *Saccharomyces cerevisiae*. *Genetics* **203**(3): 1027-1067.
- Bensaude O. 2011. Inhibiting eukaryotic transcription: Which compound to choose? How to evaluate its activity? *Transcr* **2**(3): 103-108.
- Bierbaum M, Bastiaens PI. 2013. Cell cycle-dependent binding modes of the ran exchange factor RCC1 to chromatin. *Biophys J* **104**(8): 1642-1651.
- Broach JR. 2012. Nutritional Control of Growth and Development in Yeast. *Genetics* **192**: 73-105.
- Chabbert CD, Adjalley SH, Steinmetz LM, Pelechano V. 2018. Multiplexed ChIP-Seq Using Direct Nucleosome Barcoding: A Tool for High-Throughput Chromatin Analysis. *Methods Mol Biol* **1689**: 177-194.
- Chen FX, Smith ER, Shilatifard A. 2018. Born to run: control of transcription elongation by RNA polymerase II. *Nat Rev Mol Cell Biol* **19**(7): 464-478.
- Chen M, Licon K, Otsuka R, Pillus L, Ideker T. 2013. Decoupling epigenetic and genetic effects through systematic analysis of gene position. *Cell Rep* **3**(1): 128-137.
- Chen S, Rufiange A, Huang H, Rajashankar KR, Nourani A, Patel DJ. 2015. Structure-function studies of histone H3/H4 tetramer maintenance during transcription by chaperone Spt2. *Genes Dev* **29**(12): 1326-1340.
- Cheung TH, Rando TA. 2013. Molecular regulation of stem cell quiescence. *Nat Rev Mol Cell Biol* **14**(6): 329-340.
- Cramer P, Bushnell DA, Fu J, Gnatt AL, Maier-Davis B, Thompson NE, Burgess RR, Edwards AM, David PR, Kornberg RD. 2000. Architecture of RNA polymerase II and implications for the transcription mechanism. *Science* **288**(5466): 640-649.
- Cucinotta CE, Arndt KM. 2016. SnapShot: Transcription Elongation. *Cell* **166**(4): 1058-1058 e1051.
- Dantuma NP, Acs K, Luijsterburg MS. 2014. Should I stay or should I go: VCP/p97-mediated chromatin extraction in the DNA damage response. *Exp Cell Res* **329**(1): 9-17.
- De Jonge WJ, Brok M, Kemmeren P, Holstege FCP. 2019. An extensively optimized chromatin immunoprecipitation protocol for quantitatively comparable and robust results. *bioRxiv*: doi.org/10.1101/835926.
- Deng P, Zhou Y, Jiang J, Li H, Tian W, Cao Y, Qin Y, Kim J, Roeder RG, Patel DJ et al. 2018. Transcriptional elongation factor Paf1 core complex adopts a spirally wrapped solenoidal topology. *Proc Natl Acad Sci USA* **115**(40): 9998-10003.
- Douglas AC, Smith AM, Sharifpoor S, Yan Z, Durbic T, Heisler LE, Lee AY, Ryan O, Gottert H, Surendra A et al. 2012. Functional analysis with a barcoder yeast gene overexpression system. "G3: Genes, Genomes, Genetics" **2**(10): 1279-1289.
- Duina AA, Miller ME, Keeney JB. 2014. Budding Yeast for Budding Geneticists: A Primer on the *Saccharomyces cerevisiae* Model System. *Genetics* **197**: 33-48.

- Ehara H, Sekine SI. 2018. Architecture of the RNA polymerase II elongation complex: new insights into Spt4/5 and Elf1. *Transcr* **9**(5): 286-291.
- Exinger F, Lacroute F. 1992. 6-Azauracil inhibition of GTP biosynthesis in *Saccharomyces cerevisiae*. *Curr Genet* **22**(1): 9-11.
- Feng Y, Zhang Y, Ebright RH. 2016. Structural basis of transcription activation. *Science* **352**(6291): 1330-1333.
- Fowler DM, Fields S. 2014. Deep mutational scanning: a new style of protein science. *Nat Methods* **11**(8): 801-807.
- Franz A, Ackermann L, Hoppe T. 2016. Ring of Change: CDC48/p97 Drives Protein Dynamics at Chromatin. *Front Genet* **7**: 73.
- Frottin F, Schueder F, Tiwary S, Gupta R, Körner R, Schlichthaerle T, Cox J, Jungmann R, Hartl FU, Hipp MS. 2019. The nucleolus functions as a phase-separated protein quality control compartment. *Science* **365**: 342-347.
- Grigull J, Mnaimneh S, Pootoolal J, Robinson MD, Hughes TR. 2004. Genome-wide analysis of mRNA stability using transcription inhibitors and microarrays reveals posttranscriptional control of ribosome biogenesis factors. *Mol Cell Biol* **24**(12): 5534-5547.
- Guidi M, Ruault M, Marbouty M, Loïodice I, Cournac A, Billaudeau C, Hocher A, Mozziconacci J, Koszul R, Taddei A. 2015. Spatial reorganization of telomeres in long-lived quiescent cells. *Genome Biol* **16**: 206.
- Handschumacher RE, Welch AD. 1956. Microbial studies of 6-azauracil, an antagonist of uracil. *Cancer Res* **16**(10 Part 1): 965-969.
- Hantsche M, Cramer P. 2017. Conserved RNA polymerase II initiation complex structure. *Curr Opin Struct Biol* **47**: 17-22.
- Hsieh TH, Weiner A, Lajoie B, Dekker J, Friedman N, Rando OJ. 2015. Mapping Nucleosome Resolution Chromosome Folding in Yeast by Micro-C. *Cell* **162**(1): 108-119.
- Jones RD, Enam C, Ibarra R, Borrer HR, Mostoller KE, Fredrickson EK, Lin J, Chuang E, March Z, Shorter J et al. 2019. The extent of Ssa1/Ssa2 Hsp70 chaperone involvement in nuclear protein quality control degradation varies with the substrate. *Mol Biol Cell*: mbcE18020121.
- Jones RD, Gardner RG. 2016. Protein quality control in the nucleus. *Curr Opin Cell Biol* **40**: 81-89.
- Kaplan CD, Jin H, Zhang IL, Belyanin A. 2012. Dissection of Pol II trigger loop function and Pol II activity-dependent control of start site selection in vivo. *PLoS Genet* **8**(4): e1002627.
- Krebschull JM, Zador AM. 2018. Cellular barcoding: lineage tracing, screening and beyond. *Nat Methods* **15**(11): 871-879.
- Kim TS, Liu CL, Yassour M, Holik J, Friedman N, Buratowski S, Rando OJ. 2010. RNA polymerase mapping during stress responses reveals widespread nonproductive transcription in yeast. *Genome Biol* **11**(7): R75.
- Koerber RT, Rhee HS, Jiang C, Pugh BF. 2009. Interaction of transcriptional regulators with specific nucleosomes across the *Saccharomyces* genome. *Mol Cell* **35**(6): 889-902.
- Korthout T, Poramba-Liyanage DW, van Kruijsbergen I, Verzijlbergen KF, van Gemert FPA, van Welssem T, van Leeuwen F. 2018. Decoding the chromatin proteome of a single genomic locus by DNA sequencing. *PLoS Biol* **16**(7): e2005542.
- Kuang Z, Cai L, Zhang X, Ji H, Tu BP, Boeke JD. 2014. High-temporal-resolution view of transcription and chromatin states across distinct metabolic states in budding yeast. *Nat Struct Mol Biol* **21**.
- Kwak H, Lis JT. 2013. Control of transcriptional elongation. *Annu Rev Genet* **47**: 483-508.
- Lauinger L, Li J, Shostak A, Cemel IA, Ha N, Zhang Y, Merkl PE, Obermeyer S, Stankovic-Valentin N, Schafmeier T et al. 2017. Thiolutin is a zinc chelator that inhibits the Rpn11 and other JAMM metalloproteases. *Nat Chem Biol* **13**(7): 709-714.

- Li L, Miles S, Melville Z, Prasad A, Bradley G, Breeden LL. 2013. Key events during the transition from rapid growth to quiescence in budding yeast require posttranscriptional regulators. *Mol Biol Cell* **24**: 3697-3709.
- Li X, Egervari G, Wang Y, Berger SL, Lu Z. 2018. Regulation of chromatin and gene expression by metabolic enzymes and metabolites. *Nat Rev Mol Cell Biol* **19**(9): 563-578.
- Liang C, Stillman B. 1997. Persistent initiation of DNA replication and chromatin-bound MCM proteins during the cell cycle in *cdc6* mutants. *Genes Dev* **11**(24): 3375-3386.
- Ljungdahl PO, Daignan-Fornier B. 2012. Regulation of amino acid, nucleotide, and phosphate metabolism in *Saccharomyces cerevisiae*. *Genetics* **190**(3): 885-929.
- Lotan R, Bar-On VG, Harel-Sharvit L, Duek L, Melamed D, Choder M. 2005. The RNA polymerase II subunit Rpb4p mediates decay of a specific class of mRNAs. *Genes Dev* **19**(24): 3004-3016.
- Madine MA, Swietlik M, Pelizon C, Romanowski P, Mills AD, Laskey RA. 2000. The roles of the MCM, ORC, and Cdc6 proteins in determining the replication competence of chromatin in quiescent cells. *J Struct Biol* **129**(2-3): 198-210.
- Makde RD, England JR, Yennawar HP, Tan S. 2010. Structure of RCC1 chromatin factor bound to the nucleosome core particle. *Nature* **467**(7315): 562-566.
- Malik I, Qiu C, Snavely T, Kaplan CD. 2017. Wide-ranging and unexpected consequences of altered Pol II catalytic activity in vivo. *Nucl Acids Res* **45**(8): 4431-4451.
- Markov D, Naryshkina T, Mustaev A, Severinov K. 1999. A zinc-binding site in the largest subunit of DNA-dependent RNA polymerase is involved in enzyme assembly. *Genes Dev* **13**(18): 2439-2448.
- Martin BJE, Chruscicki AT, Howe LJ. 2018. Transcription Promotes the Interaction of the FAcilitates Chromatin Transactions (FACT) Complex with Nucleosomes in *Saccharomyces cerevisiae*. *Genetics* **210**(3): 869-881.
- Mason PB, Struhl K. 2005. Distinction and relationship between elongation rate and processivity of RNA polymerase II in vivo. *Mol Cell* **17**(6): 831-840.
- Mayer A, Lidschreiber M, Siebert M, Leike K, Soding J, Cramer P. 2010. Uniform transitions of the general RNA polymerase II transcription complex. *Nat Struct Mol Biol* **17**(10): 1272-1278.
- McCann M, Kellett A, Kavanagh K, Devereux M, Santos AL. 2012. Deciphering the antimicrobial activity of phenanthroline chelators. *Curr Med Chem* **19**(17): 2703-2714.
- McCullough L, Connell Z, Petersen C, Formosa T. 2015. The Abundant Histone Chaperones Spt6 and FACT Collaborate to Assemble, Inspect, and Maintain Chromatin Structure in *Saccharomyces cerevisiae*. *Genetics* **201**(3): 1031-1045.
- McGinty RK, Tan S. 2016. Recognition of the nucleosome by chromatin factors and enzymes. *Curr Opin Struct Biol* **37**: 54-61.
- McKnight JN, Boerma JW, Breeden LL, Tsukiyama T. 2015. Global Promoter Targeting of a Conserved Lysine Deacetylase for Transcriptional Shutoff during Quiescence Entry. *Mol Cell* **59**: 732-743.
- Mews P, Zee BM, Liu S, Donahue G, Garcia BA, Berger SL. 2014. Histone methylation has dynamics distinct from those of histone acetylation in cell cycle reentry from quiescence. *Mol Cell Biol* **34**: 3968-3980.
- Miles S, Li L, Davison J, Breeden LL. 2013. Xbp1 directs global repression of budding yeast transcription during the transition to quiescence and is important for the longevity and reversibility of the quiescent state. *PLoS Genet* **9**: e1003854.
- Miller C, Schwalb B, Maier K, Schulz D, Dumcke S, Zacher B, Mayer A, Sydow J, Marcinowski L, Dolken L et al. 2011. Dynamic transcriptome analysis measures rates of mRNA synthesis and decay in yeast. *Mol Syst Biol* **7**: 458.
- Musso M, Bianchi-Scarra G, Van Dyke MW. 2000. The yeast CDP1 gene encodes a triple-helical DNA-binding protein. *Nucl Acids Res* **28**(21): 4090-4096.

- Myers SA, Wright J, Peckner R, Kalish BT, Zhang F, Carr SA. 2018. Discovery of proteins associated with a predefined genomic locus via dCas9-APEX-mediated proximity labeling. *Nat Methods* **15**(6): 437-439.
- Namdar M, Kearsley SE. 2006. Analysis of Mcm2-7 chromatin binding during anaphase and in the transition to quiescence in fission yeast. *Exp Cell Res* **312**(17): 3360-3369.
- Nemergut ME, Mizzen CA, Stukenberg T, Allis CD, Macara IG. 2001. Chromatin docking and exchange activity enhancement of RCC1 by histones H2A and H2B. *Science* **292**(552121269703): 1540-1543.
- Nourani A, Robert F, Winston F. 2006. Evidence that Spt2/Sin1, an HMG-like factor, plays roles in transcription elongation, chromatin structure, and genome stability in *Saccharomyces cerevisiae*. *Mol Cell Biol* **26**(4): 1496-1509.
- O'Duibhir E, Lijnzaad P, Benschop JJ, Lenstra TL, van Leenen D, Groot Koerkamp MJ, Margaritis T, Brok MO, Kemmeren P, Holstege FC. 2014. Cell cycle population effects in perturbation studies. *Mol Syst Biol* **10**: 732.
- Panday A, Grove A. 2016. The high mobility group protein HMO1 functions as a linker histone in yeast. *Epigen Chrom* **9**: 13.
- Pathak R, Singh P, Ananthkrishnan S, Adamczyk S, Schimmel O, Govind CK. 2018. Acetylation-Dependent Recruitment of the FACT Complex and Its Role in Regulating Pol II Occupancy Genome-Wide in *Saccharomyces cerevisiae*. *Genetics* **209**(3): 743-756.
- Porrua O, Libri D. 2015. Transcription termination and the control of the transcriptome: why, where and how to stop. *Nat Rev Mol Cell Biol* **16**(3): 190-202.
- Prasad R, D'Arcy S, Hada A, Luger K, Bartholomew B. 2016. Coordinated Action of Nap1 and RSC in Disassembly of Tandem Nucleosomes. *Mol Cell Biol* **36**(17): 2262-2271.
- Prasad R, Xu C, Ng DTW. 2018. Hsp40/70/110 chaperones adapt nuclear protein quality control to serve cytosolic clients. *J Cell Biol* **217**: 2019-2032.
- Pufall MA, Kaplan CD. 2013. Mechanisms of eukaryotic transcription. *Genome Biol* **14**(9): 311.
- Qiu C, Kaplan CD. 2019. Functional assays for transcription mechanisms in high-throughput. *Methods* **159-160**: 115-123.
- Radonjic M, Andrau JC, Lijnzaad P, Kemmeren P, Kockelkorn TT, van Leenen D, van Berkum NL, Holstege FC. 2005. Genome-wide analyses reveal RNA polymerase II located upstream of genes poised for rapid response upon *S. cerevisiae* stationary phase exit. *Mol Cell* **18**(2): 171-183.
- Roy KR, Smith JD, Vonesch SC, Lin G, Tu CS, Lederer AR, Chu A, Suresh S, Nguyen M, Horecka J et al. 2018. Multiplexed precision genome editing with trackable genomic barcodes in yeast. *Nat Biotechnol* **36**(6): 512-520.
- Ruone S, Rhoades AR, Formosa T. 2003. Multiple Nhp6 molecules are required to recruit Spt16-Pob3 to form yFACT complexes and to reorganize nucleosomes. *J Biol Chem* **278**(46): 45288-45295.
- Rutledge MT, Russo M, Belton J-M, Dekker J, Broach JR. 2015. The yeast genome undergoes significant topological reorganization in quiescence. *Nucl Acids Res* **43**: 8299-8313.
- Sagot I, Laporte D. 2019a. The cell biology of quiescent yeast - a diversity of individual scenarios. *J Cell Sci* **132**(1).
- . 2019b. Quiescence, an individual journey. *Curr Genet* **65**(3): 695-699.
- Sainsbury S, Bernecky C, Cramer P. 2015. Structural basis of transcription initiation by RNA polymerase II. *Nat Rev Mol Cell Biol* **16**(3): 129-143.
- Schmidtman E, Anton T, Rombaut P, Herzog F, Leonhardt H. 2016. Determination of local chromatin composition by CasID. *Nucleus* **7**(5): 476-484.
- Scrutton MC, Wu CW, Goldthwait DA. 1971. The presence and possible role of zinc in RNA polymerase obtained from *Escherichia coli*. *Proc Natl Acad Sci USA* **68**(10): 2497-2501.

- Shaw RJ, Reines D. 2000. Saccharomyces cerevisiae transcription elongation mutants are defective in PUR5 induction in response to nucleotide depletion. *Mol Cell Biol* **20**(20): 7427-7437.
- Shaw RJ, Wilson JL, Smith KT, Reines D. 2001. Regulation of an IMP dehydrogenase gene and its overexpression in drug-sensitive transcription elongation mutants of yeast. *J Biol Chem* **276**(35): 32905-32916.
- Siddiqui K, On KF, Diffley JFX. 2013. Regulating DNA replication in eukarya. *Cold Spring Harbor perspectives in biology* **5**(9): a012930-a012930.
- Struhl K, Segal E. 2013. Determinants of nucleosome positioning. *Nat Struct Mol Biol* **20**(3): 267-273.
- Sun M, Schwalb B, Pirkl N, Maier Kerstin C, Schenk A, Failmezger H, Tresch A, Cramer P. 2013. Global Analysis of Eukaryotic mRNA Degradation Reveals Xrn1-Dependent Buffering of Transcript Levels. *Mol Cell* **52**(1): 52-62.
- Sun W, Hola M, Pedley K, Tada S, Blow JJ, Todorov IT, Kearsey SE, Brooks RF. 2000. The replication capacity of intact mammalian nuclei in Xenopus egg extracts declines with quiescence, but the residual DNA synthesis is independent of Xenopus MCM proteins. *J Cell Sci* **113** (Pt 4): 683-695.
- Swygert SG, Kim S, Wu X, Fu T, Hsieh T-H, Rando OJ, Eisenman RN, Shendure J, McKnight JN, Tsukiyama T. 2018. Condensin-Dependent Chromatin Compaction Represses Transcription Globally during Quiescence. *Mol Cell* **73**: 533-546.
- Swygert SG, Tsukiyama T. 2019. Unraveling quiescence-specific repressive chromatin domains. *Curr Genet* **65**: 1145-1151.
- Tong AH, Boone C. 2006. Synthetic genetic array analysis in Saccharomyces cerevisiae. *Methods Mol Biol* **313**: 171-192.
- Van Oss SB, Cucinotta CE, Arndt KM. 2017. Emerging Insights into the Roles of the Paf1 Complex in Gene Regulation. *Trends Biochem Sci* **42**(10): 788-798.
- van Welsem T, Korthout T, Ekkebus R, Morais D, Molenaar TM, van Harten K, Poramba-Liyanage DW, Sun SM, Lenstra TL, Srivas R et al. 2018. Dot1 promotes H2B ubiquitination by a methyltransferase-independent mechanism. *Nucl Acids Res* **46**(21): 11251-11261.
- Vasseur P, Tonazzini S, Ziane R, Camasses A, Rando OJ, Radman-Livaja M. 2016. Dynamics of Nucleosome Positioning Maturation following Genomic Replication. *Cell Rep* **16**(10): 2651-2665.
- Venters B, Pugh BF. 2009. How eukaryotic genes are transcribed. *Crit Rev Biochem Mol Biol* **44**(2-3): 117-141.
- Verma R, Aravind L, Oania R, McDonald WH, Yates JR, 3rd, Koonin EV, Deshaies RJ. 2002. Role of Rpn11 metalloprotease in deubiquitination and degradation by the 26S proteasome. *Science* **298**(5593): 611-615.
- Verzijlbergen KF, van Welsem T, Sie D, Lenstra TL, Turner DJ, Holstege FCP, Kerkhoven RM, van Leeuwen F. 2011. A barcode screen for epigenetic regulators reveals a role for the NuB4/HAT-B histone acetyltransferase complex in histone turnover. *PLoS Genet* **7**(10): e1002284.
- Vinayachandran V, Reja R, Rossi MJ, Park B, Rieber L, Mittal C, Mahony S, Pugh BF. 2018. Widespread and precise reprogramming of yeast protein-genome interactions in response to heat shock. *Genome Res* **28**(3): 357-366.
- Vlaming H, McLean C, Korthout T, Alemdehy MF, Hendriks S, Lancini C, Palit S, Klarenbeek S, Kwesi-Maliepaard EM, Molenaar TM et al. 2019. Conserved crosstalk between histone deacetylation and H3K79 methylation generates DOT1L-dose dependency in HDAC1-deficient thymic lymphoma. *EMBO J* **38**(14): e101564.
- Vlaming H, Molenaar TM, van Welsem T, Poramba-Liyanage DW, Smith DE, Velds A, Hoekman L, Korthout T, Hendriks S, Altelaar AM et al. 2016. Direct screening for chromatin status on DNA barcodes in yeast delineates the regulome of H3K79 methylation by Dot1. *eLife* **5**: 10.7554/eLife.18919.

- Vos SM, Farnung L, Boehning M, Wigge C, Linden A, Urlaub H, Cramer P. 2018. Structure of activated transcription complex Pol II-DSIF-PAF-SPT6. *Nature* **560**(7720): 607-612.
- Wada T, Becskei A. 2017. Impact of Methods on the Measurement of mRNA Turnover. *Int J Mol Sci* **18**(12).
- Weiner A, Chen HV, Liu CL, Rahat A, Klien A, Soares L, Gudipati M, Pfeffner J, Regev A, Buratowski S et al. 2012. Systematic Dissection of Roles for Chromatin Regulators in a Yeast Stress Response. *PLoS Biol* **10**(7): e1001369.
- Wierer M, Mann M. 2016. Proteomics to study DNA-bound and chromatin-associated gene regulatory complexes. *Hum Mol Genet* **25**(R2): R106-R114.
- Wu F, Liu Y, Zhu Z, Huang H, Ding B, Wu J, Shi Y. 2011. The 1.9A crystal structure of Prp20p from *Saccharomyces cerevisiae* and its binding properties to Gsp1p and histones. *J Struct Biol* **174**(1): 213-222.
- Xie Y, Zheng M, Chu X, Chen Y, Xu H, Wang J, Zhou H, Long J. 2018. Paf1 and Ctr9 subcomplex formation is essential for Paf1 complex assembly and functional regulation. *Nat Commun* **9**(1): 3795.
- Xu Y, Bernecky C, Lee CT, Maier KC, Schwalb B, Tegunov D, Plitzko JM, Urlaub H, Cramer P. 2017. Architecture of the RNA polymerase II-Paf1C-TFIIS transcription elongation complex. *Nat Commun* **8**: 15741.
- Yan Z, Costanzo M, Heisler LE, Paw J, Kaper F, Andrews BJ, Boone C, Giaever G, Nislow C. 2008. Yeast Barcoders: a chemogenomic application of a universal donor-strain collection carrying bar-code identifiers. *Nat Methods* **5**(8): 719-725.
- Young CP, Hillyer C, Hokamp K, Fitzpatrick DJ, Konstantinov NK, Welty JS, Ness SA, Werner-Washburne M, Fleming AB, Osley MA. 2017. Distinct histone methylation and transcription profiles are established during the development of cellular quiescence in yeast. *BMC Genomics* **18**(1): 107-107.
- Zaidi H, Hoffman EA, Shetty SJ, Bekiranov S, Auble DT. 2017. Second-generation method for analysis of chromatin binding with formaldehyde-cross-linking kinetics. *J Biol Chem* **292**(47): 19338-19355.
- Zhou H, Liu Q, Shi T, Yu Y, Lu H. 2015. Genome-wide screen of fission yeast mutants for sensitivity to 6-azauracil, an inhibitor of transcriptional elongation. *Yeast* **32**(10): 643-655.

Figure 1. Outline of expanded and optimized Epi-Decoder analysis

(A) Three Epi-Decoder-HO libraries were generated by crossing the TAP-tag library to an expanded HO-BC library in three different ways to combine each TAP-tag protein with three independent barcodes (see also Supplemental Fig S1a). The HO-BC locus consists of a constitutively expressed 1.5 kb KanMX resistance gene integrated at the *HO* locus, controlled by the heterologous *AgTEF1* promoter and terminator from *Ashbya gossypii*, and flanked by a promoter-proximal BC_UP and a terminator-proximal BC_DN. Downstream of BC_DN lies an origin of replication (ARS404). **(B)** Clones of each Epi-Decoder-HO library are combined and processed in two separate pools and used for ChIP of TAP-tagged proteins (spheres with black handle). The BCs (colored lines), which flank the KanMX reporter gene (grey box) at the HO locus are amplified from ChIP and Input and indexed, allowing for the pools to be combined and counted by massive parallel sequencing. The relative barcode count (IP/input) reports on protein abundance of each TAP-tagged protein (~4250) at the barcoded locus. **(C)** Comparison of the binding scores (IP/input) of both BC_UP and BC_DN of chromatin binders (as determined previously in (Korthout et al. 2018)) in the three Epi-Decoder-HO libraries. Indicated are the Spearman correlation coefficients and the diagonal line represent $x=y$. Density plots show the distribution of the barcode counts in each of the three replicates. See Supplemental Table S1 for counts of all proteins examined. The results for BC_UP and BC_DN separately are shown in Supplemental Fig S1B-C.

Figure 2. Chromatin-proteome dynamics at the Barcoded-HO locus upon chemical inhibition of transcription

(A) Three versions of the Chromatin-TAP-tag subset, each containing a unique set of barcode pairs, were combined in one pool (Chrom-3×BC). The culture was incubated with a-factor pheromone to synchronize the cells in G1, after which PH or AU was added and samples were collected at the time points indicated for Epi-Decoder analysis, RNA analysis, and flow cytometry. A sample treated with

vehicle was collected at the last time point and a control sample was taken of a G1-arrested culture without treatment. All experiments were carried out at 16°C to facilitate capturing dynamic binding events. **(B)** Analysis of mRNA changes over time by RT-qPCR confirmed the decay of most transcripts and an increase in the PH-responsive ZRT1 gene under conditions shown in panel A. RNA levels are relative to untreated (Pre) and normalized to a transcript from a spike-in of untreated *S. pombe* cells (see Materials and Methods) to correct for global changes (mean of three biological replicates +/- SD). **(C)** Heatmap of the binding scores (mean IP/input of three biological replicates) of selected proteins with a binding score >0.5 at any of the four local proteome time series indicated. See Supplemental Table S2 for mean binding scores of all proteins examined in the Chrom-3xBC library. Proteins were manually clustered and ranked in functional subcategories.

Figure 3. Treatment with phenanthroline leads to rapid loss of Pol II and transcription-associated proteins

(A) Differential and dynamic binding behavior of proteins representing different chromatin processes upon treatment with PH and AU. The lines indicate the three different barcode pairs of the indicated TAP-tagged proteins in the Chrom-3xBC library (Log_2 IP/input at time points indicated in Fig 2). **(B)** Zoom in of heat map of Fig 2 showing the Pol II subunits present in the library. **(C-D)** Immunoblot analysis of the largest subunit of Pol II (Rpo21-TAP) with and without PH and AU treatment in G1-arrested cells at 16°C. Pgk1, Hmo1, and a non-specific band (*) were used as loading controls. **(E)** ChIP-qPCR analysis of Rpo21 binding at the BC_UP and BC_DN regions in G1-arrested cells treated with (15 and 60 min) and without (Pre) PH at 16°C (average of three biological replicates +/- SD).

Figure 4. Differential response of transcription elongation factors to chemical inhibition of Pol II

(A) Zoom in on the heat map of Fig 2 (PH treatment), showing proteins annotated to transcription elongation. **(B)** Independent replicates of proteins related to FACT and Paf1C (Log_2 IP/input at time points as in Fig 2). The lines show the three different barcode pairs of the indicated TAP-tagged proteins in the Chrom-3 \times BC library. **(C)** ChIP-qPCR analysis of Ctr9 binding at the BC_UP and BC_DN regions in G1-arrested cells treated with (15 and 60 min) and without (Pre) PH at 16°C (average of three biological replicates \pm SD). **(D)** Immunoblot analysis of Ctr9-TAP and Paf1-TAP with and without PH treatment in G1-arrested cells at 16°C. P $gk1$ and H3 were used as loading controls. **(E)** ChIP-qPCR analysis of Srm1 binding in G1-arrested cells at 16°C, treated for 15 min with vehicle (V) or phenanthroline (PH) (average of three biological replicates \pm SD). Analyzed loci are the BC_UP and BC_DN regions, the 5' and 3' end of the endogenous *TEF1* gene, the *ADH1* promoter, the *PMA1* open reading frame, and a non-transcribed locus (van Welsem et al. 2018) (see Supplemental Table S5 for more details).

Figure 5. Chromatin proteome rewiring upon transcriptional repression in quiescence

(A) Heatmap of the HO Epi-Decoder binding scores in mid-log, Q, and NQ cells (mean IP/input of three biological replicates). See Supplemental Table S3 for mean binding scores of all proteins examined in the Chrom-3 \times BC library. Proteins were manually clustered and ranked in functional subcategories as in Fig 2C. **(B-E)** Zoom in of heat map of panel A (promoter region BC_UP, except replication for which terminator BC_DN next to the origin of replication was used) showing proteins in the indicated annotated clusters. PH treatment during G1 arrest (vehicle, 5 and 12 minutes) is shown for comparison.




Figure 1**A****B****C**






Figure 2**A****B****C**

Figure 3


A




B



C



D



E




Figure 4





Figure 5



$\text{Log}_2(\text{IP}/\text{input})$

