



Chromosome-specific telomere lengths and the minimal functional telomere revealed by nanopore sequencing

Samantha L. Sholes, Kayarash Karimian, Ariel Gershman, et al.

Genome Res. published online October 26, 2021

Access the most recent version at doi:[10.1101/gr.275868.121](https://doi.org/10.1101/gr.275868.121)

P<P	Published online October 26, 2021 in advance of the print journal.
Accepted Manuscript	Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.
Creative Commons License	This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see https://genome.cshlp.org/site/misc/terms.xhtml). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at http://creativecommons.org/licenses/by-nc/4.0/ .
Email Alerting Service	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here .

An advertisement banner with a teal background. On the left, the text reads "CRISPR and RNAi Genetic Screening. Your new superpower." in white. In the center, there is a white-bordered box containing the words "LEARN MORE" in teal. On the right, there is a photograph of a woman wearing a red superhero mask and a red cape over a grey shirt. To the right of the photo is the Cellecta logo, which consists of a cluster of green dots of varying sizes, with the word "CELLECTA" in white capital letters below it.

To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Published by Cold Spring Harbor Laboratory Press

Chromosome specific telomere lengths and the minimal functional telomere revealed by nanopore sequencing

Samantha L. Sholes^{1,2}, Kayarash Karimian^{1,2}, Ariel Gershman^{1,2}, Thomas J. Kelly⁴, Winston Timp^{1,3}, and Carol W. Greider^{1,5*}

¹Department of Molecular Biology and Genetics, Johns Hopkins University School of Medicine, Baltimore, MD, 21205, USA

²Biochemistry, Cellular and Molecular Biology Graduate Program, Johns Hopkins University School of Medicine, Baltimore, MD, 21205, USA

³Department of Biomedical Engineering, Johns Hopkins University School of Medicine, Baltimore, MD, 21205, USA

⁴Program in Molecular Biology, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY 10065

⁵Department of Molecular Cell and Developmental Biology, University of California, Santa Cruz, CA, 95064, USA

* To whom correspondence should be addressed. Tel: (831) 502-7367
Email: cgreider@ucsc.edu

Keywords: Telomere length regulation, Nanopore sequencing, *TEL1*, *RIF1*

ABSTRACT

We developed a method to tag telomeres and measure telomere length by nanopore sequencing in the yeast *S. cerevisiae*. Nanopore allows long read sequencing through the telomere, subtelomere and into unique chromosomal sequence, enabling assignment of telomere length to a specific chromosome end. We observed chromosome end specific telomere lengths that were stable over 120 cell divisions. These stable chromosome specific telomere lengths may be explained by slow clonal variation or may represent a new biological mechanism that maintains equilibrium unique to each chromosome end. We examined the role of *RIF1* and *TELI* in telomere length regulation and found that *TELI* is epistatic to *RIF1* at most telomeres, consistent with the literature. However, at telomeres that lack subtelomeric Y' sequences, *tel1Δ rif1Δ* double mutants had a very small, but significant, increase in telomere length compared to the *tel1Δ* single mutant, suggesting an influence of Y' elements on telomere length regulation. We sequenced telomeres in a telomerase-null mutant (*est2Δ*) and found the minimal telomere length to be around 75 bp. In these *est2Δ* mutants there were apparent telomere recombination events at individual telomeres before the generation of survivors, and these events were significantly reduced in *est2Δ rad52Δ* double mutants. The rate of telomere shortening in the absence of telomerase was similar across all chromosome ends at about 5 bp per generation. This new method gives quantitative, high resolution telomere length measurement at each individual chromosome end, and suggests possible new biological mechanisms regulating telomere length.

INTRODUCTION

Maintenance of the telomere length equilibrium is critical for cell survival. Telomeres shorten with each cell division while a few are stochastically elongated by the enzyme telomerase (Greider and Blackburn 1985; Teixeira et al. 2004). This balance of shortening and lengthening is exquisitely regulated to generate a normal distribution of lengths that is established around a mean or “set point”. Different species have very different telomere length set points; for example *Oxytricha* has 20-30 bp, yeast have 300 bp, humans have 10 kb, and some mouse strains have up to 80 kb mean telomere lengths (Greider 1996). How this distribution is established and maintained is not fully understood. When telomerase levels are low or absent, the distribution shifts toward shorter lengths and the shortest telomeres signal a DNA damage response. These cells undergo a permanent cell cycle arrest, termed senescence or apoptosis, depending on the cell type (Enomoto et al. 2002; d'Adda di Fagagna et al. 2003; Ijma and Greider 2003).

Loss of telomere length maintenance in humans leads to disease. Individuals with inherited mutations in telomerase, or other telomere regulatory genes, have Short Telomere Syndrome, which includes bone marrow failure, immunodeficiency, pulmonary fibrosis and other diseases (Armanios 2013). In contrast, activation of telomerase (Counter et al. 1992; Greider 1998) or inheritance of long telomeres allows cell immortalization which can predispose to cancer (Stanley and Armanios 2015; McNally et al. 2019). Thus, deviation from the mean telomere length in either direction affects cellular lifespan and plays a critical role in disease.

The yeast *S. cerevisiae* serves as an excellent model for probing mechanisms of telomere length regulation. Many genes and regulatory mechanisms that affect human telomere length were first discovered in yeast (Wellinger and Zakian 2012). Yeast telomere sequences are more variable than the consistent TTAGGG repeats found in vertebrates (Moyzis et al. 1988). They are composed of a mixture

of GT, GGT or GGGT sequence motifs, usually abbreviated, $G_{1-3}T$. The subtelomeric region in yeast contains specific families of repeated DNA.

Deletion of any component of telomerase in yeast leads to progressive telomere shortening, termed ever-shorter telomere (EST) phenotype (Lundblad and Szostak 1989; Lendvay et al. 1996). Previous work suggests that telomeres shorten by about 3-5 bp per cell division in the absence of telomerase and that this value is constant and independent of telomere length (Marcand et al. 1999; Wellinger and Zakian 2012; Xu et al. 2013). There is initially no effect of telomere shortening on cell growth and survival, but when telomeres become very short after passaging in culture for several days, most cells arrest at G2/M (Enomoto et al. 2002; IJpma and Greider 2003). A few cells can rescue their short telomeres through recombination and will generate “survivors” that grow in the absence of telomerase (Lundblad and Blackburn 1993). Survivors are generated through a break induced replication (BIR) pathway (Bosco and Haber 1998; McEachern and Haber 2006) that can initiate within the telomere repeats (Type II survivors) or in the subtelomere repeated elements (Type I survivors) (Teng and Zakian 1999). Deletion of the recombination gene *RAD52* significantly reduces production of both types of survivors (Lundblad and Blackburn 1993) although very rare survivors have been found in *est2Δ rad52Δ* double mutants (Lebel et al. 2009).

The normal distribution of the telomere length equilibrium is due primarily to sequence loss through replication and sequence addition by telomerase. Shampay and Blackburn showed that independent chromosome ends can slightly shift their mean length due to clonal variability (Shampay and Blackburn 1988). If a cell that happens to have a short telomere at Chromosome 8L, for example, seeds a new colony on a plate, the descendants of that cell will initially have a new, shorter telomere length distribution until enough divisions occur in the population to homogenize all ends towards a population mean (Shampay and Blackburn 1988).

The “protein counting” model for telomere length regulation established in *S. cerevisiae* (Marcand et al. 1997; Marcand et al. 1999) proposes that proteins bound to the telomeric repeats negatively regulate telomere elongation by telomerase, so that the probability of telomere elongation decreases with telomere length. Mean telomere length is stable when the average rate of length-dependent elongation is equal to the average rate of length-independent shortening. (Bianchi and Shore 2008). This model has received significant attention and is now the accepted model for telomere length regulation in organisms as distantly related as mammals, plants and fission yeast (Smogorzewska and de Lange 2004; Watson and Riha 2010; Wellinger and Zakian 2012; Armstrong and Tomita 2017; Maciejowski and de Lange 2017). Sequence analysis of individual telomeres showed that telomerase added between 3 bp to 179 bp (average 44bp) onto chromosome ends in one cell cycle; however only a few telomeres were elongated at each cycle. Further, shorter telomere repeat tracts were preferentially elongated compared to those with longer repeat tracts (Marcand et al. 1999; Teixeira et al. 2004). This work further supported the protein counting model by providing evidence that telomerase preferentially elongates short telomere repeat tracts. How preferential elongation of short telomeres allows the robust maintenance of the telomere length equilibrium over many cell divisions is not yet clear.

The standard technique for visualizing telomere length in yeast is by Southern blot. Genomic DNA from yeast cells is cut with a restriction enzyme near the telomere and hybridized with a probe to the telomere repeats or to the subtelomere, generating smeared bands visible on the autoradiogram (Shampay and Blackburn 1988). The smears represent the heterogenous distribution of telomere lengths across both the different chromosomes and between cells. Since restriction sites on different chromosomes are at different locations, the absolute size of each smear differs. Frequently, the restriction enzyme XhoI is used to cut genomic DNA because many telomeres have a conserved XhoI restriction site in the Y' subtelomere element. Multiple telomeres are then visualized in a heterogeneous band centered around 1.2 kb. Measuring median length of this 1.2 kb sized band can allow large changes in bulk telomere length to be

measured in various mutant backgrounds, but it is not possible to deconvolute the length of individual telomeres from each other.

To provide a more quantitative analysis of telomere length variation, we developed a nanopore sequencing method to measure telomere length. Nanopore sequencing allowed us to capture a full telomere sequence as well as the subtelomere and unique chromosomal sequence in one long read. Using this quantitative method for measuring telomere length, we re-evaluated fundamental questions about telomere length distribution and changes in mutant backgrounds that alter telomere length.

RESULTS

Telomere tagging and bioinformatic analysis

To accurately measure telomere length and retain the telomeric 3' overhangs, we modified a method (Teixeira et al. 2004) to tag the molecular end of the chromosome. We prepared high molecular weight DNA (Denis 2018) and added poly(A) to 3' ends with terminal transferase. We next annealed an Oligo(dT) primer that also contained a unique adapter sequence, termed TeloTag. The addition of *Sulfolobus* DNA polymerase IV, which lacks strand displacement and exonuclease activity, was used to generate a double-stranded blunt end, and T4 DNA ligase to seal the nicks. Nanopore library adaptors were added to allow sequencing (Fig. 1A).

Tagged genomic DNA was run on nanopore MinION or Flongle flow cells to generate Whole Genome Sequencing (WGS) data (Fig 1B). For optimal use of a nanopore MinION flow cell, samples were multiplexed, and only reads with barcodes on both ends were retained. Raw reads with telomeric sequence were extracted using TideHunter. To ensure that reads had sequenced to the end of the telomere, we retained reads only if they contained the TeloTag (Gao et al. 2019). Adapter sequences were then bioinformatically removed using Porechop to ensure that they are not calculated as part of the telomere length (<https://github.com/rrwick/Porechop>). These filters were employed on the raw reads before

alignment. The resulting reads were aligned using minimap2 to a *S. cerevisiae* reference genome, sacCer3 (Foury et al. 1998). To ensure that the reads mapped only to the correct chromosome end, we removed all reads that aligned in multiple places in the genome using SAMtools to select only reads with a map quality score greater than twenty and remove all secondary and supplementary alignments. Finally, the position where the subtelomere ends and telomere repeats begins was identified for each chromosome end, and the telomere sequence length was calculated using the aligned reads (Fig. 1C) (see Methods).

Figure 1. Telomere length measurement by nanopore sequencing. A) Schematic of method to tag the molecular end of the Chromosome. 1. Terminal transferase adds poly A sequence to the 3' overhang of the telomere. 2. A complementary Oligo(dT) sequence on the TeloTag oligonucleotide end containing a unique TeloTag sequence is annealed to the poly(A) sequence. 3. Polymerase fill-in generates double stranded blunt ends. 4. Ligase is used to seal the nick. Nanopore adaptors are then added using the manufacture's protocol. B) Bioinformatic pipeline to generate telomere length information (see methods). C) Telomere length is calculated using the chromosomal position of each telomere sequence start and calculating the distance to the TeloTag sequence at the chromosome end. A random sample of 10 sequence reads aligned at TEL01R is presented as an example.

Telomere length by nanopore sequencing recapitulates length measurement by Southern blot

To test the accuracy of our nanopore method for telomere length measurement, we compared telomere fragment length analysis by nanopore and Southern blot. We first looked at strains with expected clear differences in telomere length profiles, wildtype and *rif1Δ*. Deletion of the *RIF1* gene results in very long telomeres on Southern blots (Hardy et al. 1992), allowing us to probe a greater distribution of telomere length profiles and test whether these length differences are captured with nanopore sequencing. We measured telomere fragment length after XhoI digest using a probe against the common Y' subtelomere element; this will measure telomere lengths in bulk, probing 17 of 32 subtelomeres (Fig. 2A, Supplemental Fig. S1C). We ran all samples in duplicate on a Southern blot and included an additional probe to the unique sequence at CEN4 that runs at 1.4k b as an internal molecular marker for densitometry plots (Fig. 2A, Supplemental Fig. S1C). We used densitometry scanning to quantify the signal on the Southern blot (see Methods).

On the same samples, we performed our nanopore sequencing method as described (Fig. 1; Methods). Nanopore sequencing data statistics are summarized in Supplemental Table S1. To directly compare to Southern blot telomere length, we included the internal chromosomal sequence up to the XhoI site for Y' on the nanopore telomere read file. The histogram of the nanopore telomere sequence read lengths aligned well with the Southern blot densitometry for both wildtype telomeres and the much longer *rif1Δ* telomeres in two different strains (Fig. 2B, Supplemental Fig. S2A). The differences at the edges of the telomere length distribution between the Nanopore and noisier Southern densitometry data highlight the increased ability to detect subpopulations in Nanopore that are not detected Southern blotting densitometry scans (Fig. 2D, Supplemental Fig. S2B). We used WALTER, a tool for Southern blot densitometry that generates summary length statistics (Lycka et al. 2021), to compare Southern blot and nanopore telomere length profiles, and again saw comparable results (Supplemental Fig. S3A) (Levy et al. 1992). We then generated Q-Q plots to examine the distribution of nanopore and Southern density data against a normal distribution (Supplemental Fig. S4A&B) and against each other (Supplemental Fig. S4C). The linearity of the Q-Q plots of both the Southern and nanopore data confirm that these profiles resemble a normal distribution (See Supplemental Fig S4 for details), and we assumed a normal distribution for our statistical analysis throughout this work, including calculating correlation coefficients between data sets, performing *t*-tests, and performing analysis of the means testing. The sample standard deviations of nanopore and Southern density distributions were similar and there was no significant difference in the means by Welch's *t*-test (Supplemental Fig. S3A).

Because the Y' probe measures 17 telomere length distributions simultaneously, it is challenging to detect potential small effects and chromosome specific telomere lengths with this probe on a Southern blot. To assess the length distribution of single chromosome ends, we generated strains in which the subtelomere was replaced by a unique sequence. We used two different strains (Shubin et al. 2021) with unique sequence at either Chromosome end TEL01L Unique End 1 (strain UE1), or at Chromosome end TEL11R: Unique End 11 (strain UE11) (Supplemental Fig. 1A/B). We examined the telomere lengths of

these single TEL01L and TEL11R chromosome ends in both wildtype and *rif1Δ* backgrounds, using a probe to the unique ends (Fig. 2C, Supplemental Fig. S1D). Again, we saw good agreement between the single telomere length distributions determined by nanopore sequencing and Southern blotting (Fig. 2D, Supplemental Fig. 2B). In both genetic backgrounds, we observed no significant differences between the sample means (Supplemental Fig. S3B). We conclude that nanopore sequencing can accurately measure telomere fragment length in a bulk population or single telomere population across different telomere length profiles.

Figure 2. Telomere fragment length measured by Nanopore sequencing accurately reflects length measurements by Southern blot. A) Southern blot probed with Y' subtelomeric probe and with CEN4 probe (arrow) as a marker at 1.4kb to calibrate for densitometry. Two replicates of wildtype (WT) cells and two replicates of *rif1Δ* cells are shown for each strain, UE1 and UE11. B) Comparison of densitometry (dotted lines) of Southern blot shown in A) and Nanopore sequencing (histogram fill bars) for UE1 strain. The light orange bars represent nanopore read counts for WT cells, the light purple bars represent nanopore read counts for *rif1Δ* cells. The spike in densitometry data at 1400 bp is due to the CEN4 internal control signal. C) Southern blot probed with the Unique single telomere TEL01L probe and CEN4 probe (arrow). D) Comparison of densitometry (dotted lines) of Southern blot in C) and Nanopore sequencing (bars) of UE1 TEL01L. The light orange bars represent nanopore read counts for WT cells, the light purple bars represent nanopore read counts for *rif1Δ* cells.

To determine the most effective flow cell type to achieve the minimum number of reads required for telomere length measurement using nanopore sequencing, we compared the MinION flowcells to the lower cost, but lower yield, Flongle flowcells. Multiplexing two samples per Flongle and MinION flowcell yielded many fewer reads from the Flongle (~150-250 per barcode) vs the MinION (~2000 reads per barcode), but we found no statistically significant differences between the mean and distributions of bulk telomere lengths across technical replicates and flow cell types (Supplemental Fig. S5). Thus, while the lower cost Flongle flow cells yield reliable bulk telomere length data, the low number of reads does not allow analysis of individual chromosome ends (see below) and thus may not provide as useful a platform for investigating telomere biology.

Each chromosome end shows reproducible chromosome-specific telomere length distributions

Probing telomere lengths on unique chromosome ends is challenging with Southern blots, but nanopore sequence reads are long enough to span the subtelomeric elements and extend into unique chromosomal sequence. To examine individual chromosome ends, we sequenced three independent wildtype clones and three *rif1Δ* clones (Supplemental Fig. S6B). For the wildtype, we used two independent clones of UE1 and one clone of UE11. Each clone was isolated by picking an independent single colony from a freshly streaked plate of cells (Supplemental Fig. 6B). We analyzed both the bulk telomere length of all telomeres and individual chromosome ends (Fig. 3A).

Figure 3. Nanopore sequencing mapping of the telomere length distribution of each individual chromosome end. A) Bulk telomere length (all) and telomere length for each chromosome end were mapped by aligning to internal unique sequences. Three independent clones were sequenced: two from UE1 (UE1-1 and UE1-2) and one from UE11 (see supplemental figure S6). Telomere lengths are reported as violin plots with a line at the mean. A dotted line shows the mean of the total population at 343.3 for WT cells. Analysis of the means (ANOM) multiple contrast test of each telomere against the grand mean of all telomere length profiles was performed (see Supplemental Figure S7A) and p values adjusted by the Bonferroni method are reported as $* < 0.05$, $** < 0.01$, $*** < 0.001$, $**** < 0.0001$, or not significant (n.s.). To note, TEL01L is engineered (as described in the text) in the UE1 strains and TEL11R is engineered in the UE11 strains. B) Pairwise correlations of mean telomere length across each of the three replicates for WT. A linear model was fit to the data (dotted line) to calculate the correlation constant, reported as the r value. Marginal density of each dataset is plotted as solid black lines across the top and right of the plots to demonstrate the spread of telomere lengths across each population. C) Clonal variation was observed at TEL10R, TEL03L, and TEL08R in WT cells. Individual telomere length reads are represented as single points within the violin plots. The significance of the difference between two samples was tested using a multiple *t*-test (methods) and p values adjusted by the Bonferroni method are reported as above. D) Model of clonal variation where the telomere length of the majority of clones remains at the parental average, but occasional length outliers may seed new distributions represented by Clone 1 and Clone 100. E). Three independent clones from the W303-0 parent were passages for 120 population doublings in independent cultures. The resulting clones W303-1, W303-3, and W303-3 were sequenced along with the parent W303-0. (See Supplemental Figure S8A). Telomeres for each chromosome end were mapped by aligning to internal unique sequence. Analysis of the means (ANOM) multiple contrast test of each telomere against the grand mean of all telomere length profiles was performed (see Supplemental Figure S7B) and p values adjusted by the Bonferroni method are reported as $* < 0.05$, $** < 0.01$, $*** < 0.001$, $**** < 0.0001$, or not significant (n.s.).

We tested alignment to the S288C ((Foury et al. 1998) as well as the W303 (Berlin et al. 2015; Matheson et al. 2017) references genomes (Supplemental Fig. S7) and found the genome assembly in the subtelomeres was more complete for S288C and allowed reads to correctly map to more specific telomeres, so we used S288C as the reference. We were able to individually map 20 of 32 total

chromosome ends with unique subtelomere and upstream genic sequences in our UE1 and UE11 strains. However, we were not able to map 12 chromosome ends uniquely (TEL01R, TEL02R, TEL03R, TEL06L, TEL07L, TEL09L, TEL12L, TEL12R, TEL13R, TEL14R, TEL16L, and TEL16R) due to variation in subtelomeric structure from the reference genome. Subtelomeric elements are known to recombine and generate end diversity (Louis and Haber 1990; Louis and Haber 1992; Maxwell et al. 2004), and we therefore excluded these 12 difficult to map telomeres in our analysis.

We were surprised to find that some chromosome ends showed telomere length distributions that were consistent across three independent clones, but distinctly longer or shorter from the population mean of all telomeres (Fig. 3A). To quantify the differences, we applied analysis of the means (ANOM) to compare the mean length of individual chromosome ends to the population mean length of all the telomeres, and found that eight chromosome ends (TEL04L, TEL04R, TEL08L, TEL10L, TEL11L, TEL11R, TEL13L, and TEL14L) had significantly shorter telomeres than the population mean, while six chromosome ends (TEL02L, TEL03L, TEL05L, TEL05R, TEL07R, and TEL09R) had significantly longer telomeres than the population mean (Supplemental Fig. S8A) (Pallmann and Hothorn 2016). A pairwise comparison of the three datasets suggests that telomere length of individual chromosome ends was consistent across replicates with correlation coefficients (r) of 0.96, 0.91, and 0.91 (Fig. 3B.) We note that for chromosomes for which both ends could be analyzed, there was a trend toward concordance with both ends being long like TEL05L/5R and TEL015L/15R, or both ends being short like TEL04L/4R, TEL08L/8R, and TEL11L/11R but there was no correlation between the length of chromosomes and the mean length of telomeres.

To determine whether chromosome end specific lengths were maintained in the absence of *Rif1*, we analyzed two *rif1* Δ clones from UE1-1 and UE11 (Supplemental Fig. 6). Each clone exhibited telomeres whose mean lengths were significantly different from the clonal population mean (Supplemental Fig. S10A,B,C). However, in contrast to the wildtype cells, we did not see consistent length differences at

individual chromosome ends in the two clones, evidenced by the correlation coefficient of 0.039 (Supplemental Fig. S10D). Because of the lack of correlation, we performed ANOM separately for the two clones and found significant deviations from the mean as expected from the broader length distribution seen in Figure 2 (Supplemental Fig. S10B,C). These results suggest that deletion of *RIFI* may disrupt the mechanism responsible for maintenance of telomere-specific lengths, or that the very broad length distribution obscures the differences between specific telomeres.

The apparent chromosome end specific telomere lengths in wildtype cells could be due to stochastic clonal variation as previously proposed (Shampay and Blackburn 1988). That is, when a cell with a short telomere on Chromosome 4L forms a new colony, it will seed a new telomere distribution that will center around a shorter midpoint (Fig. 3D). We found a few cases of statistically significant variation among the three clones analyzed. In particular, TEL03L was significantly shorter in UE11 than in UE1-1 and UE1-2, and TEL08R was significantly longer in UE1-1 than UE-1 and UE11, and TEL10R in UE1-1 and UE1-2 than in UE11 (Fig. 3C). This rate of clonal variation was infrequent compared to the rate of clonal variation in the previous study (Shampay and Blackburn 1988), although this may be due to different number of cell divisions in propagating clones.

Shampay et al. introduced the concept of the telomere length ‘set point’, in which all telomeres will gradually move back toward a population mean length distribution and suggested that differences in mean telomere length between clones might be erased during further propagation with all telomeres evolving toward a common mean length. Therefore, given enough cell divisions in the presence of telomerase, we might expect all chromosome ends to approach a mean value. In fact, Shampay et al. showed that continued growth in liquid culture led to a broadening of the length distribution, which they interpreted as homogenization of all end lengths (Shampay and Blackburn 1988). The “protein counting model” for telomere length regulation (Marcand et al. 1997) suggests that telomere length is actively regulated around a mean based on the number of Rap1 proteins bound and, thus the length of telomere sequence.

These length regulation models are supported by the observation that telomerase apparently preferentially elongates short telomeres while ignoring long ones (Teixeira et al. 2004, Marcand, 1999). To test whether the telomere lengths would homogenize with more cell divisions, we isolated a clone of W303 cells (W303-0) and propagated it in liquid culture for 120 generations. Strain W303-0 is the parent of the clones UE1-1, UE1-2, and UE11 analyzed in figure 3A (Supplemental Figure S6A) and has a similar pattern of end specific telomere lengths with some exceptions, such as that at TEL03L, which could be due to clonal variation. After 120 generations in liquid culture, three independent clones were isolated on plates and sequenced (W303-1, W303-2, and W303-3) (Supplemental Figure S9A). We expected, based on previous models, that the telomere length distributions at individual chromosome ends would revert towards a grand population mean with more cell divisions. However, we saw that most chromosome specific telomere lengths were maintained across all W303 clones, for example: TEL02L, TEL04L, TEL04R, TEL05L, TEL05R, TEL07R, TEL08L, TEL08R, TEL09R, TEL10L, TEL10R, TEL11L, TEL11R, TEL13L, and TEL14L (Fig. 3E, Supplemental Fig. S8B). Although there were some examples of significant deviations from the initial length, for example at TEL03L, TEL06R, TEL15L and TEL15R, the major trend was reproducibility of the parents' chromosome end specific lengths (Supplemental Fig. S8A and B). To quantify the variation between different clones, we examined pairwise correlations between the W303-0 parent and the W303-1, W303-2, and W303-3 subclones, and found correlation coefficients (r) of 0.92, 0.82, 0.89, 0.82, 0.86, and 0.75 (Supplemental Fig. S9B). Our data indicate that after 120 doublings, the chromosome end specific telomere lengths are largely maintained. This implies a very slow rate of reversion to a common mean length distribution, if it occurs at all, and implies new models for telomere length regulation are needed to explain this data.

Y' elements have small effect on telomere length regulation by *TEL1* and *RIF1*

The possibility of chromosome end specific telomere lengths raised the idea that subtelomeric sequence might influence telomere length. A similar suggestion was made by Craven and Petes who proposed that the subtelomeric X and Y' elements influenced the effects of *TEL1* and *RIF1* on telomere length (Craven

and Petes 1999). Specifically, they suggested that *TEL1* is epistatic to *RIF1* only at telomeres with Y' elements, but not at telomeres with only X elements. That is, telomeres are long in *rif1Δ* mutants, short in *tel1Δ* mutants, and also short in the *tel1Δ rif1Δ* double mutant at Y' telomeres.

We reexamined this question using both Southern blots and nanopore sequencing. We generated *tel1Δ*, *rif1Δ*, and *tel1Δ rif1Δ* mutants in the in UE1 background (see Methods). As expected, in the Southern blot probed with Y' sequence, *rif1Δ* mutants had long telomeres and *tel1Δ* mutants had short telomeres. The double mutant, *tel1Δ rif1Δ*, had short telomeres very similar to the single *tel1Δ* mutant (Fig. 4A), indicating that *TEL1* is epistatic to *RIF1* at telomeres with Y' elements. Nanopore sequencing showed the same result at chromosomes with Y' elements; *tel1Δ rif1Δ* double mutants had short telomeres indistinguishable from *tel1Δ* telomere lengths, supporting the previous finding (Fig. 4B). However, for chromosomes ends that lacked Y' elements and thus had only X elements, telomere length in *tel1Δ rif1Δ* was slightly longer (~65 bp on average) and statistically different from *tel1Δ* (Fig. 4D). This trend also was seen at the unique Chromosome TEL01L where the subtelomeric sequence was removed entirely. Both nanopore sequencing (Fig. 4D, TEL01L) and Southern blot (Fig. 4C) showed that telomeres in the *tel1Δ rif1Δ* double mutant were longer than the *tel1Δ* alone. While the difference in length was not as large as seen by Craven and Petes, it is reproducible and suggests the need for further study.

Figure 4. Y' element subtelomeric sequences have a small effect on telomere length regulation by *TEL1* and *RIF1*. A) Southern blot of WT, *tel1Δ*, *rif1Δ*, and *tel1Δ rif1Δ* cells with bulk telomere length hybridized with the Y' and CEN4 probes (arrow). B) Individual telomere lengths determined by nanopore for telomere ends containing Y' subtelomeric elements reported for *tel1Δ*, *rif1Δ*, and *tel1Δ rif1Δ* cells. C) Southern blot of WT, *tel1Δ*, *rif1Δ*, and *tel1Δ rif1Δ* samples hybridized with Unique probe to TEL01L and CEN4 probe (arrow) to examine unique TEL01L with no subtelomeric elements. D) Individual telomere lengths determined by nanopore for telomere ends containing only X subtelomeric elements or no telomeric elements (TEL01L) reported for *tel1Δ*, *rif1Δ*, and *tel1Δ rif1Δ* cells. A multiple *t*-test was used to assess the significance of telomere length differences between genotypes, and p values are reported as * <0.05 , ** <0.01 , *** <0.001 , and **** <0.0001 , or not significant (n.s.).

Telomere shortening in the absence of telomerase reveals the minimal functional telomere

To examine the dynamics of telomere loss in cells without telomerase, we examined telomerase null cells. We sporulated an *EST2/est2Δ RAD52/rad52Δ* heterozygous diploid and selected individual haploid colonies of *est2Δ*, *est2Δ rad52Δ*, and wildtype (*EST2*). We grew these genotypes from a single colony and passaged each three times in liquid culture (Supplemental Fig. S11). To determine population doubling numbers, we estimated growth from a single cell to colony formation to be about 20 cell doublings. Both *est2Δ* and *est2Δ rad52Δ* cells grew slowly after 55 population doublings, and *est2Δ rad52Δ* cells exhibited significantly slower growth than *est2Δ*. Telomere shortening in the absence of telomerase leads to cell cycle arrest (Enomoto et al. 2002; Ijima and Greider 2003), and also to telomeric elongation by Break Induced Replication (BIR) (Bosco and Haber 1998) that ultimately allows the outgrowth of “survivors” (Lundblad and Blackburn 1993). This implies there must be a competition between arrest and BIR, and presumably all chromosomes must be elongated for phenotypic survivors to emerge.

The ability to measure the length distributions of individual telomeres using nanopore sequencing suggested that it might be possible to detect possible BIR events in the progeny of the *est2Δ* cells. Such events would not be visible by traditional Southern blotting of telomeres in bulk (Fig. 5A/B). In fact, we observed some telomere length distributions that were highly skewed, unlike any in the wild type populations that we studied (see for example TEL03L and TEL08R in Fig. 5C). To quantify this effect, we calculated the sample standard deviation of each telomere distribution in wildtype cells (P0, see Methods) and in *est2Δ* cells. For most telomeres we observed, as expected, that the sample standard deviation decreased relative to that of the parental wildtype telomere during passaging of the *est2Δ* cells. This observation is consistent with the fact that in the absence of Est2, telomeres progressively shorten, but their lengths are bounded by the minimum telomere length consistent with cell survival (see below). We found seven telomeres for which the sample standard deviation actually increased relative to that of the wildtype parent: TEL02R, TEL03L, TEL04L, TEL07R, TEL08R, TEL11R, and TEL15R (Fig. 5C).

These telomere samples contain subpopulations with unexpectedly long telomeres, which we refer to below as outliers. Qualitative inspection of the distributions confirmed that these telomere distributions as highly skewed (Fig. 5C). While we did not passage *est2Δ* cells for enough generations to observe outgrowth of survivors, our data suggest that telomere elongation events due to BIR may occurring in these cells at an early stage, albeit at a relatively low frequency.

Figure 5. Telomere shortening and outlier elongation events in a telomerase null, *est2Δ*, background at individual telomeres over time. Freshly generated *est2Δ* and *est2Δ rad52Δ* haploids were grown up (P1, 35 doublings) and passaged twice (P2, 55 doublings and P3, 75 doublings) to allow for telomere shortening (see Methods). Telomere length was measured by Southern blot hybridized with the Y' probe and by nanopore sequencing on the same samples. A) Southern blot of WT, P1, P2, and P3 *est2Δ* hybridized with the Y' probe. B) Southern blot of WT, P1, P2, and P3 *est2Δ rad52Δ* hybridized with the Y' probe. C/D) Individual telomere length of C) *est2Δ* and D) *est2Δ rad52Δ* cells determined by nanopore sequencing of P1, P2, and P3. We calculated the sample standard deviations of (σ) of each telomere at every passage and in the wildtype parental strain. Telomeres where the standard deviation was greater than that of the wildtype are considered outliers and are marked with a red carrot. At P2, 55 doublings, in the *est2Δ* sample, not enough data was collected at TEL07R and TEL15R to plot and at P3, 75 doublings, in the *est2Δ rad52Δ* sample, not enough data was collected at TEL04L and TEL05L to plot.

To further examine the role of BIR in the generation of outlier telomere lengths, we examined the *est2Δ rad52Δ* cells, which are recombination deficient (Fig. 5B/D). If the long telomere outliers in *est2Δ* cells are due to BIR, we would expect to see fewer of these outlier length increases in *est2Δ rad52Δ* cells. Indeed, there was a trend toward fewer outliers as we saw only five examples of long outlier telomeres at TEL09L and TEL15R in the *est2Δ rad52Δ* cells. (Fig. 5D). These length increases may be due to rare events that occur even in *est2Δ rad52Δ* cells that allow telomere elongation (Lebel et al. 2009; Claussin and Chang 2016). Nanopore sequencing will allow detailed mechanistic insight into telomere elongation by BIR and the generation of survivors.

To determine the rate of telomere shortening in the absence of telomerase before cell division arrest, we compared the rate of shortening at individual chromosome ends in *est2Δ* and *est2Δ rad52Δ* cells over three outgrowths or passages (P1, 2, 3) (Supplemental Fig. S11). We exclude data where the telomere

length increases likely due to BIR over the 75 population doublings at TEL02R and TEL11L in *est2Δ* cells, as it would skew the analysis. To establish the initial rate of shortening, we need to establish the initial telomere length before *EST2* deletion. Since this was not possible in the newly dissected haploid *est2Δ* strain, we instead used the wildtype telomere length for each chromosome as a proxy for the initial length and designated this P0 (Supplemental Fig. S11). The rate of shortening for each chromosome end was very similar from P0 to P1; there was a loss of 5.33 bp/population doubling in *est2Δ* and 5.13 bp/population doubling in *est2Δ rad52Δ* cells (Fig. 6A/B). This telomere shortening rate is similar to other estimates of 3-4 bp/population doubling (Marcand et al. 1999; Wellinger and Zakian 2012; Xu et al. 2013). If we include the later passages P2 and P3 in the rate, there is a continuous decline in the apparent rate because cells with very short telomeres do not divide and thus are lost from the calculated rate. To compare individual ends we separately examined the rate of shortening from P1 to P3, and found all ends had a similar loss rate of 0.76 bp/population doubling in *est2Δ* cells and 0.39 bp/population doubling in *est2Δ rad52Δ* cells (Fig. 6A/B). The loss rates of the P1 to P3 *est2Δ* cells were higher than those of *est2Δ rad52Δ* cells on average perhaps due to the more severe growth defects we observed in *est2Δ rad52Δ* compared to *est2Δ* cells. The rates of telomere shortening at each individual chromosome end were very similar in both *est2Δ* and *est2Δ rad52Δ*. Our quantitative single telomere data support previous analysis (Marcand et al. 1999) that telomere shortening rates are similar across chromosome ends.

Figure 6. Telomere shortening rate in the absence of telomerase. A) The progression of mean telomere length at individual chromosome ends in *est2Δ* cells over 3 passages. TEL01-08 and TEL09-16 are shown on separate graphs for clarity. The initial telomere length, P0, is estimated from the wildtype parental strain for each chromosome, and so it is not the exact genotype (See results). The line of best fit to the average of the shortening rate is shown as a black dotted line. The rate of shortening for P0 to P1 and P1 to P3 are plotted separately because the two rates differ and each rate is reported above the line. B) The rate of telomere shortening for the *est2Δ rad52Δ* strain was measured as described above in A.

To determine the minimum telomere length in telomerase-null cells, we examined telomeres at early passages before survivor events were predicted to occur (Lundblad and Blackburn 1993). We calculated

the bottom 99th length percentile at each chromosome end and identified critically short telomeres in both the *est2Δ* and *est2Δ rad52Δ* ranging from 70 to 80 bp. The shortest three values are highlighted in each genotype at each passage (Table 1). This minimal telomere length is also consistent with the minimal length we found in telomerase-positive wildtype and *tel1Δ* cells of ~75 bp. (Table 1). Previous work has suggested that telomeres shorter than 100 bp could be considered “critically short” (Abdallah et al. 2009). Our sequence analysis now more precisely identified a minimum of ~75 bp at individual chromosome ends.

Table 1. Minimum telomere length is approximately 75bp. The bottom 99th percentile of telomere length at the three shortest chromosome ends per genotype at each passage in *est2Δ*, *est2Δ rad52Δ*, as well as for *wildtype* and *tel1Δ* are shown.

DISCUSSION

Nanopore telomere sequencing uncovered the unexpected finding of chromosome end specific telomere length differences that were preserved across multiple independent clones over 120 population doublings.

The stable maintenance of telomere-specific length differences, with mean values differing by as much as 200 bp, implies that current models for telomere length homeostasis require some revision (Marcand et al. 1997; Marcand et al. 1999; Teixeira et al. 2004; Greider 2016). Both the protein counting, and the replication fork models suggest that the average rate of telomere elongation by telomerase decreases with telomere length because of increased binding of inhibitory proteins to the telomeric repeats. In the context of this model, our new data indicates that the functional relationship between the number of bound proteins and the decrease in the average elongation rate is different for different telomeres. Thus, the balance point between length-dependent elongation and length-independent shortening occurs at different lengths for different telomeres. This finding implies the existence of yet undiscovered telomere-specific influences on length homeostasis. It also suggests that the observed preferential elongation of short telomeres (Marcand et al. 1999; Teixeira et al. 2004) must be applied in a telomeric-specific sense,

i.e. preferential elongation of telomeres that are short relative to their specific equilibrium length. In addition, our new data indicates that the rate of shortening in the absence of telomerase is similar for all telomeres, confirming and extending previous studies (Marcand et al. 1999; Wellinger and Zakian 2012; Xu et al. 2013). It will be of interest to determine whether in the absence of telomerase, telomeres with a shorter equilibrium length may have a higher probability of becoming critically short and triggering cell cycle arrest.

Our work adds to other evidence showing that counting Rap1 binding proteins does not fully explain telomere length regulation. Alteration of the telomerase RNA template to generate TTAGGG repeats that do not bind Rap1 has little effect on telomere length (Henning et al. 1998; Alexander and Zakian 2003; Brevet et al. 2003; Bah et al. 2011). In addition, recent work suggests the apparent preferential elongation of short telomeres may be due to replication fork collapse in telomere repeats with subsequent elongation by telomerase (Paschini et al. 2020). Current models of length regulation need to be modified to account for end specific equilibrium distributions and for Rap1 independent length maintenance.

What mechanism might regulate these end specific telomere lengths? One obvious hypothesis is that sequences adjacent to the telomere repeats might affect length regulation, either through the sequence itself, through chromatin modifications or transcription, or the non-coding RNA TERRA (Feuerhahn et al. 2010). In fact, there were small differences in response to *RIF1* length regulation at chromosomes that had Y' elements compared to those that did not (Figure 4). Subtelomeric DNA-binding proteins such as Tbf1 and Reb1 that have binding sites in X and Y' are thought to play some role in length regulation (Berthiau et al. 2006). However, the small effect of these proteins cannot explain the large length differences of up to 200 bp on different chromosome ends observed in wildtype cells. The hint that there may be concordance between telomeres on the same chromosome (Figure 3A and D) suggests that perhaps chromosome territories within the nucleus (Noma 2017) have an effect on length regulation. Alternatively, topologically associated domains (TADs) (Noma 2017) or localization to the nuclear

periphery (Sobecki et al. 2018) may play a role in chromosome end-specific telomere length differences. The fact that deletion of *RIFI* disrupted telomere-specific lengths might provide a hint at the mechanism, however it might be that that deletion of *RIFI* simply broadens the length distribution obscuring any differences between telomeres. Nanopore sequencing of telomeres provides a method to further interrogate these interesting new questions.

In the absence of telomerase, all telomeres shortened at a similar rate as predicted by the end replication problem. A shortening rate of about 5 bp/population doubling is similar to what has been seen previously with other methods (Marcand et al. 1999; Abdallah et al. 2009; Wellinger and Zakian 2012; Xu et al. 2013). The shortest telomeres in telomerase null cells, as well as in wildtype and *tell1* Δ cells, were around 75 bp. This is shorter than previously suggested (Abdallah et al. 2009), and allows specific hypotheses to be made for what might cause loss of telomere function. For instance, Rap1 binds telomeric DNA approximately every 20 bp (Gilson et al. 1993; Wahlin and Cohn 2000), so telomere dysfunction might be due to loss of Rap1 binding, or perhaps loss of the terminal nucleosome binding. However, it was observed previously that a yeast chromosome with human telomere repeats can be elongated by telomerase, even though Rap1 cannot bind human telomere repeats (Henning et al. 1998; Alexander and Zakian 2003; Brevet et al. 2003). This suggests that there may be distinct factors, independent of Rap1, that negatively regulate telomere length. We note that even at a minimal length of 75 bp Ku's should be able to bind DNA through its end binding function (Gravel et al. 1998; Chen et al. 2018). Given this new data on end specific telomere lengths, it will be of interest to re-evaluate the role of specific protein binding in determining length regulation.

This novel nanopore sequencing method allows for the analysis of single molecules and provides the ability to assign telomere length measurement to individual chromosome ends. Leveraging this, we were able to uncover new aspects of telomere biology. Nanopore sequencing can be done in any laboratory and requires relatively little capital investment. Even a smaller, less expensive Flongle flowcell provides a

suitable number of independent reads for bulk telomere length measurement. With further improvements and the use of targeted enrichment (Gilpatrick et al. 2020; Kovaka et al. 2021), the cost and yield of this method can be even further improved. Telomere length measurement by nanopore sequencing will allow the telomere field to study new questions and revisit past unanswered questions in telomere biology. A different method of long read sequencing, PacBio sequencing, has recently been applied to human telomere sequences, (Grigorev et al. 2021) indicating long read sequencing technologies are widely useful in telomere research. It will be of interest to determine whether chromosome end-specific telomere length differences are generalizable to other organisms, and perhaps even humans, and understanding how they are established and maintained will be a fascinating new area of telomere biology to explore.

METHODS

Biological Resources

Yeast strains and plasmid vectors are listed in (Supplemental Table S4). Yeast strains are available upon request and plasmid vectors are available upon request or from Addgene (Supplemental Table S4/S6).

Molecular Cloning

Yeast molecular cloning, culturing, sporulation and transformation were performed as previously described in (Green and Sambrook 2012; Keener et al. 2019; Shubin et al. 2021).

The UE1 (previous named 1L 5×UAS landing pad) and UE11 (previously named 11R 5×UAS landing pad) constructs were designed *in silico* using SnapGene software and constructed as described in (Shubin et al. 2021). UE11 was generated in the same manner as UE1, but with homology at Chromosome arm 11R, rather than Chromosome arm 1L. The subtelomere was removed in these strains using homology to the unique gene region directly upstream of the telomere and ensuring recombination with a digested product with a 3' overhang that is endogenously elongated by telomerase.

Yeast strains culturing and transformation

CVy61 (W303-0 WT haploids), UE1 (previously CALy117, *1L::leu2*), UE11 (previously CALy119, *11R::leu2*). UE1 *rif1* Δ (previously CALy202, *1L::leu2*, *rif1* $\Delta::kanMX$), UE11 *rif1* Δ (previously CALy204, *11R::leu2*, *rif1* $\Delta::kanMX$), and UE1 *tel1* Δ (previously CALy472, *1L::leu2*, *tel1* $\Delta::hphMX4$) strains were generated as previously described in (Shubin et al. 2021). These strains were renamed UE1 and UE11 to highlight the function of a unique end, rather than a 5×UAS landing pad, that was important to this study. A list of all yeast strains and genotypes is presented in Supplemental Table 4.

The UE1 *tel1* Δ *rif1* Δ was generated by transformation. Briefly, transformation was performed by treating 50 mL of logarithmically growing UE1 *rif1* Δ cells with 0.1M lithium acetate (L6883; Sigma-Aldrich) and adding 1 μ g of linear DNA to replace *TEL1* with *tel1* $\Delta::hphMX4$ cassette for one-step integration. Cells were incubated at 30°C for 10 minutes before the addition of 0.5 ml 40% polyethylene glycol (PEG4000, P4338; Sigma-Aldrich), 0.1M lithium acetate. Cells were then incubated for 30 minutes at 30°C followed by a 45 minute heat shock at 42°C. Cells were plated after a water wash onto YPD + HygromycinB, (Corning, at 200ug/ml). Integration at the *TEL1* locus was confirmed by junction polymerase chain reaction (PCR). A list of primers and plasmids used in this study is presented in Supplemental Tables 5 and 6, respectively.

To perform the W303-0,1,2,3 experiment, CVy61 was streaked to single colony on YPD media. One colony was selected (W303-0) and grown in 5 mL YPD to 0.5 OD. This culture was split into two subcultures: 1) growth to 1.0 OD in 100 mL YPD broth for High Molecular Weight genomic DNA extraction (see below) and called W303-0 for nanopore sequencing and 2) growth in 10 mL YPD broth to saturation for 24 hours and passaged as 5 μ L of saturated culture in 10mL fresh YPD. This was repeated a total of 5 times before cells were plated onto YPD media. Three independent colonies were selected (W303-1, W303-2, and W303-3) to inoculate three cultures of 100 mL YPD broth for DNA extraction as

described for subculture #1. We estimate 120 population doublings between W303-0 and W303-1,2,3. See Supplemental Figure S6A for illustration.

To perform the UE1 and UE11 experiments in wildtype and *rif1* Δ backgrounds, frozen stocks of UE1 wildtype, UE11 wildtype, UE1 *rif1* Δ , and UE11 *rif1* Δ were streaked onto YPD media to single colony. One colony each was selected from UE1 wildtype, UE11 wildtype, UE1 *rif1* Δ , and UE11 *rif1* Δ , grown to 0.5 OD in 5 mL YPD broth, followed by growth to 1 OD in 100 mL YPD for DNA extraction and nanopore sequencing as described above. An additional colony was selected from UE1 wildtype and UE1 *rif1* Δ and grown in 10 mL YPD to saturation for 24 hours and passaged as 5 μ L of saturated culture in 10 mL fresh YPD. This was repeated a total of 3 times in liquid culture before cells were grown to 0.5 OD in 5 mL YPD broth, then to 1.0 OD in 10 mL YPD broth for DNA extraction and sequencing of UE1-2 wildtype and *rif1* Δ . We estimate 60 population doublings between UE1-1 and UE1-2. See Supplemental Figure S6B for illustration.

To generate telomerase-null *est2* Δ and *est2* Δ *rad52* Δ cells, the heterozygous diploid strain yAY139 (Ijima and Greider 2003) was sporulated and dissected to yield sister spores for analysis.

Since these mutants will senesce with passaging, we attempted to keep cell division to a minimum, allowing telomeres to shorten, but not form survivors (Chen et al. 2001). For both *est2* Δ and *est2* Δ *rad52* Δ , cells were taken directly from the original dissection plate, and grown to saturation in 10 mL cultures of YPD broth. These mutant cultures were: 1) grown to 1 OD in 100 mL for DNA extraction and sequencing, called passage 1 (P1) or 2) diluted to 10^5 cells/mL in YPD broth every 24 hours before being grown to 1 OD in 100 mL for DNA extraction and sequencing. Dilution and growth for these mutants was continued for two more rounds, collecting cells after each passage called P2 and P3. See Supplemental Figure S10 for illustration. A wildtype haploid spore was also selected from the original

dissection plate of yAY139, grown to saturation in 10 mL YPD, and then grown to 1 OD in 100 mL YPD for DNA extraction and sequencing. This sample was called passage 0 (P0). The sample standard deviation of the telomeres in the P0 wildtype sample was compared to the telomerase null telomere sample standard deviations as a threshold for characterizing potential recombinant telomere length outliers in the mutants. We additionally confirmed this outlier measurement with a qualitative examination of comparisons of mutant and wildtype telomere length distributions. This P0 wildtype sample was used a 0-population doubling initial length in examining the rate of telomere shortening.

High molecular weight genomic DNA extraction

We modified a DNA isolation protocol (Denis 2018) to collect high molecular weight (HMW) genomic DNA from yeast. Cultures were grown at 30°C to 1.0 OD in 100 mL YPD broth, collected in a Sorvall Legend XTR centrifuge with swinging bucket rotor for 5 minutes at 1500×g and washed once with 20 mL TE buffer (10mM Tris, 1mM EDTA, pH8.0).

To generate spheroplasts, cell pellets in a 50 ml conical centrifuge tube (Falcon) were resuspended in 2 mL 1M sorbitol with 0.5 mL of 2.5 mg/mL Zymolyase 100T (Amsbio, 120493-1) and shaken gently (75rpm) at 30°C for 1 hour. Spheroplasts were collected at 300×g for 4 minutes in the swinging bucket centrifuge and the supernatant was carefully discarded.

Pelleted spheroplasts were lysed by resuspension in 1.7 mL lysis buffer (100mM Tris, pH7.5, 100mM EDTA, pH8.0, 500mM NaCl, 1% PVP40) and 250 µL 10% SDS. After inverting the tube gently 10 times, 20 µL of 100mg/mL RNase A (Sigma-Aldrich, R6513) was added and the mixture was shaken at 75 rpm at 37°C for 30 minutes. To digest protein, 25 µL of 20mg/mL Proteinase K (Invitrogen 25530-015) was added, and the mixture was incubated for 2 hours at 50°C with occasional inversion until there was a uniform cloudy suspension.

To precipitate and remove protein, 5 mL TE and 2.5 mL 5M potassium acetate, pH 7.5 were added to the suspension and the tube was inverted 10 times to mix, then placed on ice for 5 minutes. The tube was inverted again and placed on ice for an additional 5 minutes before centrifugation at 4°C for 15 minutes at 4000×g in the swinging bucket centrifuge. The supernatant was decanted into a clean tube and the centrifugation was repeated. The supernatant was again decanted into a clean tube.

DNA was precipitated by adding an equal volume of 100% isopropanol and swirling the tube until the DNA strands are observed. Precipitated DNA was collected from the bottom of the tube using a p1000 wide bore pipette tip and placed in a clean Eppendorf microcentrifuge tube. The DNA was pelleted in a minicentrifuge for 15 seconds and the supernatant carefully decanted. 500 µL 75% ethanol was added to rinse the DNA pellet. After decanting the wash, the residual ethanol was removed with a pipette tip. The DNA pellet was resuspended in 50-200 µL of EB buffer (Qiagen, 10mM Tris-Cl, pH8.5), depending on the size of the pellet. The DNA was allowed to resuspend at 4°C on an inverting rotator overnight. The final concentration of resuspended DNA was measured on a Qubit 3.0 Fluorometer (Thermo Fisher Scientific).

DNA molecular end tagging

To tag the molecular end, HMW genomic DNA (gDNA, see above) was first A-tailed using terminal transferase (NEB M0315) in a reaction by incubating 2 µg of gDNA, 1× terminal transferase buffer, 1× CoCl₂, 5mM dATP, and 20U of terminal transferase enzyme for 1hr at 37C, followed by heating at 70°C for 10 minutes to stop the reaction.

Five primers were designed to make a mixture of TeloTags (see Supplemental Table S6).

To the A-tailed gDNA reaction, the following were added to complete the TeloTag addition:

1× ThermoPol Reaction Buffer Pack (NEB B9004S), 2.5mM dNTP mix, 1mM ATP, 2.5mM TeloTag primer mix, and 4 units of *Sulfolobus* DNA Polymerase IV(NEB M0327S). The TeloTag reaction was incubated in a Veriti 96 well Thermal Cycler (Applied Biosystems) for 1min at 56°C and 10 minutes at 72°C. 400 units of T4 ligase (NEB M0202) was then added to the reaction and allowed to continue incubating at 12°C for 20 minutes. Ampere XP beads (Beckman Coulter) were used to purify the tagged gDNA from the reaction mixture.

Size selection of the tagged gDNA was performed using the Short Read Eliminator XS kit (Circulomics, SS-100-121-01), which retains DNA molecules greater than 10 kb. Once this step is completed, the tagged gDNA is ready for nanopore library preparation (see below).

Southern blotting and densitometry

HMW genomic DNA was extracted and quantitated as described above. Telomere length analysis by Southern Blot was performed on the same strains used for nanopore sequencing according to protocol described in (Kaizer et al. 2015). Briefly, 250 ng of gDNA was digested with PvuII and XhoI for 2-4 hours at 37C and resolved on a 1% agarose gel overnight. Two concentrations of 2-log ladders (NEB) were included as reference for analysis, 10ng for single telomere assays and for bulk telomere assays. After transfer, the membrane was hybridized with ³²P radiolabeled 2-log ladder and PCR fragments unique to either TEL01L, TEL11R (a purified PCR product of the LEU2 gene and CYC1 terminator as described in Shubin, et al.), Y' subtelomeric elements as described in Kaizer, et al. or to CEN4 (Kaizer et al. 2015; Laterreur et al. 2018; Shubin et al. 2021). After washing, the membrane was exposed on a Storage Phosphor screen (GE Healthcare) for 1 day for bulk assays and 4-5 days for single telomere assays. Southern images were captured on a STORM using ImageQuant (GE Healthcare). ImageQuant GEL files were downloaded into Adobe PhotoShop CS6 and saved as TIF files.

Southern blot densitometry was measured to allow for comparison to Nanopore data as it shows both the normal distribution and spread of telomere length for each sample. However, these plots are often inconsistent and are not a direct measure of telomere length as they are generated from the Southern blot image. To somewhat counter this, we measured densitometry using two programs. We first used ImageQuant to generate a distribution fully across each sample lane to be overlaid with Nanopore telomere fragment length data, so that all molecular markers and the resolution cut-off would be visible (Fig. 2C, D, F, G). In ImageQuant, we used the 2-log ladder and the CEN4 band at 1.4kb as molecular markers to determine a linear range and the resulting .csv files were loaded into RStudio to generate telomere length plots. Second, PhotoShop TIF files were inverted and loaded into WALTER ScanToIntensity and IntensityAnalyzer to remove background and generate intensity profiles of the telomere signal alone (Lycka et al. 2021). The reported summary statistics were loaded into RStudio to produce boxplots of telomere length.

Nanopore library preparation and sequencing

Library preparation of 1µg input DNA with a molecular tag was performed using the Native barcoding genomic DNA kit (Oxford Nanopore Technologies EXP-NBD104 and SQK-LSK109). Samples were run on a MinION flow cell (v.9.4.1) or Flongle flow cell (v.9.4.1 pore) using the MK1B or GridION for 24 to 72 hours, and were operated using the MinKNOW software (v.19.2.2). Only reads with barcodes on both ends were selected to pass. The read counts and characteristics are shown in Supplemental Tables S1, S2, S3. We note that this WGS sequencing from a MinION flow cell provides enough coverage (~50-200 reads per telomere) to analyze specific single chromosome ends, but Flongle flow cells produce only ~10 reads at a single chromosome end, which at present are not enough to analyze specific single chromosome ends. On average, we found that a Flongle flow cell costs 33 dollars per 100 telomere reads and 0.35 dollars per 100 total reads (including flow cell and reagent costs), while a MinION flow cell costs 6.9 dollars per 100 telomere reads and 0.04 dollars per 100 total reads. We conclude that it may be more

efficient to highly multiplex samples on a MinION flow cell to generate bulk telomere data than to use a Flongle flow cell to examine individual telomeres.

Nanopore analysis, alignment and telomere length calculation

Base calling to generate FASTQ sequencing files from the electrical signal data (FAST5 files) was performed with GUPPY (v.4.2.3). FASTQ reads containing telomere sequence were selected using TideHunter (v1.4.4) (Gao et al. 2019). Raw base called reads containing the TeloTag were selected using SeqKit (v0.16.1) before alignment (Shen et al. 2016). Nanopore adapters, barcodes, and the TeloTag were removed from reads using Porechop (v0.2.4) before alignment as to not skew length measurement (<https://github.com/rrwick/Porechop>). FASTQ sequence reads were aligned to a custom reference genome modified from sacCer3(Foury et al. 1998) (where 2 kb of the *S. cerevisiae* consensus 13mer telomere repeat sequence (Wahlin and Cohn 2000; Wellinger and Zakian 2012; Li 2018) was added to each chromosome end) using the map-ont preset of minimap2 (v2.18) for alignment of noisy long read sequences. Custom references for WT W303-0, UE1, UE11, and S288C are available below in GitHub and Supplemental Code. Reads with a map quality score of less than twenty, as well as secondary and supplementary reads were filtered out using SAMtools (v1.12), so that only high confidence alignment primary reads were selected to prevent measuring telomere length of any reads that multi-mapped to multiple telomeres (Li et al. 2009). The resulting BAM files were examined in the Integrated Genomics Viewer (IGV) (Robinson et al. 2011; Thorvaldsdottir et al. 2013) to ensure that telomere reads are not clipped during alignment, that there are no large indels in the telomere alignment to skew length measurement, and that the telomere alignment begins at the expected “telomere start” position in the reference. BEDTools (v2.30.0) was used to generate files with the start and end chromosomal position of each read from the aligned reads (Quinlan and Hall 2010). These BED files and a BED file with the start position of each telomere end (available for WTW303, UE1, and UE11 references at below) were read into a custom script in RStudio (available below) (Team 2020) to calculate telomere length for each read and generate plots using the R package ggplot2 (v3.3.3) (Wichman et al. 2016). This R script contains an

additional filter to remove any read name duplications within a BED file, to again ensure that any multi-mapping reads are removed. Statistics were calculated using the R package rstatix (v0.7.0) and ANOM (v0.2)(Pallmann 2017; Kassambara 2021).

DATA ACCESS

The whole genome raw sequencing reads data generated in this study have been submitted to the NCBI BioProject database (<https://www.ncbi.nlm.nih.gov/bioproject/?term=PRJNA730563>) under accession number PRJNA730563. The reference genomes, telomere start position files, and code for the above analysis and plots are hosted on GitHub (https://github.com/timplab/Telomere_Length) and available as Supplemental Code.

COMPETING INTEREST STATEMENT

W.T. has two patents (8,748,091 and 8,394,584) licensed to Oxford Nanopore Technologies.

ACKNOWLEDGEMENTS

This study was supported by the NIH R35CA209974 (to CWG) and NIGMS T32 GM007445 (to the BCMB graduate training program).

We acknowledge Carla Connelly for strain generation and Southern blotting. We thank Dr. Brendan Cormack, Dr. Rebecca Keener, and Dr. Calla Shubin for thoughtful discussions about experimental design and conclusions. We thank Dr. Rebecca Keener also for assistance in generating analysis scripts. We would like to thank Carla Connelly, Dr. Brendan Cormack, Dr. Rebecca Keener, Dr. Paul Hook, Dr. Akshi Jasani, and Thea Egelhofer for critical reading and editing of the manuscript.

REFERENCES

- Abdallah P, Luciano P, Runge KW, Lisby M, Geli V, Gilson E, Teixeira MT. 2009. A two-step model for senescence triggered by a single critically short telomere. *Nat Cell Biol* **11**: 988-993.
- Alexander MK, Zakian VA. 2003. Rap1p telomere association is not required for mitotic stability of a C(3)TA(2) telomere in yeast. *Embo J* **22**: 1688-1696.
- Armanios M. 2013. Telomeres and age-related disease: how telomere biology informs clinical paradigms. *J Clin Invest* **123**: 996-1002.
- Armstrong CA, Tomita K. 2017. Fundamental mechanisms of telomerase action in yeasts and mammals: understanding telomeres and telomerase in cancer cells. *Open Biol* **7**.
- Bah A, Gilson E, Wellinger RJ. 2011. Telomerase is required to protect chromosomes with vertebrate-type T2AG3 3' ends in *Saccharomyces cerevisiae*. *J Biol Chem* **286**: 27132-27138.
- Berlin K, Koren S, Chin CS, Drake JP, Landolin JM, Phillippy AM. 2015. Assembling large genomes with single-molecule sequencing and locality-sensitive hashing. *Nat Biotechnol* **33**: 623-630.
- Berthiau AS, Yankulov K, Bah A, Revardel E, Luciano P, Wellinger RJ, Geli V, Gilson E. 2006. Subtelomeric proteins negatively regulate telomere elongation in budding yeast. *Embo J* **25**: 846-856.
- Bianchi A, Shore D. 2008. How telomerase reaches its end: mechanism of telomerase regulation by the telomeric complex. *Mol Cell* **31**: 153-165.
- Bosco G, Haber JE. 1998. Chromosome break-induced DNA replication leads to non-reciprocal translocations and telomere capture. *Genetics* **150**: 1037-1047.
- Brevet V, Berthiau AS, Civitelli L, Donini P, Schramke V, Geli V, Ascenzioni F, Gilson E. 2003. The number of vertebrate repeats can be regulated at yeast telomeres by Rap1-independent mechanisms. *Embo J* **22**: 1697-1706.
- Chen H, Xue J, Churikov D, Hass EP, Shi S, Lemon LD, Luciano P, Bertuch AA, Zappulla DC, Geli V et al. 2018. Structural insights into yeast telomerase recruitment to telomeres. *Cell* **173**: 331-343.
- Chen Q, Ijima A, Greider CW. 2001. Two survivor pathways that allow growth in the absence of telomerase are generated by distinct telomere recombination events. *Mol Cell Biol* **21**: 1819-1827.
- Claussin C, Chang M. 2016. Multiple Rad52-Mediated Homology-Directed Repair Mechanisms Are Required to Prevent Telomere Attrition-Induced Senescence in *Saccharomyces cerevisiae*. *PLoS Genet* **12**: e1006176.
- Counter CM, Avilion AA, LeFeuvre CE, Stewart NG, Greider CW, Harley CB, Bacchetti S. 1992. Telomere shortening associated with chromosome instability is arrested in immortal cells which express telomerase activity. *Embo J* **11**: 1921-1929.
- Craven RJ, Petes TD. 1999. Dependence of the regulation of telomere length on the type of subtelomeric repeat in the yeast *Saccharomyces cerevisiae*. *Genetics* **152**: 1531-1541.
- d'Adda di Fagagna F, Reaper PM, Clay-Farrace L, Fiegler H, Carr P, Von Zglinicki T, Saretzki G, Carter NP, Jackson SP. 2003. A DNA damage checkpoint response in telomere-initiated senescence. *Nature* **426**: 194-198.
- Denis E, Sanchez, S., Mairey, B., Beluche, O., Cruaud, C., Lemainque, A., Wincker, P., Barbe, V. 2018. Extracting high molecular weight genomic DNA from *Saccharomyces cerevisiae*. *protocol exchange*.
- Enomoto S, Glowczewski L, Berman J. 2002. MEC3, MEC1, and DDC2 are essential components of a telomere checkpoint pathway required for cell cycle arrest during senescence in *Saccharomyces cerevisiae*. *Mol Biol Cell* **13**: 2626-2638.
- Feuerhahn S, Iglesias N, Panza A, Porro A, Lingner J. 2010. TERRA biogenesis, turnover and implications for function. *FEBS Lett* **584**: 3812-3818.
- Foury F, Roganti T, Lecrenier N, Purnelle B. 1998. The complete sequence of the mitochondrial genome of *Saccharomyces cerevisiae*. *FEBS Lett* **440**: 325-331.
- Gao Y, Liu B, Wang Y, Xing Y. 2019. TideHunter: efficient and sensitive tandem repeat detection from noisy long-reads using seed-and-chain. *Bioinformatics* **35**: i200-i207.

- Gilpatrick T, Lee I, Graham JE, Raimondeau E, Bowen R, Heron A, Downs B, Sukumar S, Sedlazeck FJ, Timp W. 2020. Targeted nanopore sequencing with Cas9-guided adapter ligation. *Nat Biotechnol* **38**: 433-438.
- Gilson E, Roberge M, Giralado R, Rhodes D, Gasser SM. 1993. Distortion of the DNA double helix by RAP1 at silencers and multiple telomeric binding sites. *J Mol Biol* **231**: 293-310.
- Gravel S, Larrivee M, Labrecque P, Wellinger RJ. 1998. Yeast Ku as a regulator of chromosomal DNA end structure. *Science* **280**: 741-744.
- Green M, Sambrook J. 2012. *Molecular Cloning: A Laboratory Manual*
- Greider CW. 1996. Telomere length regulation. *Annu Rev Biochem* **65**: 337-365.
- Greider CW. 1998. Telomerase activity, cell proliferation, and cancer. *Proc Natl Acad Sci U S A* **95**: 90-92.
- Greider CW. 2016. Regulating telomere length from the inside out: the replication fork model. *Genes Dev* **30**: 1483-1491.
- Greider CW, Blackburn EH. 1985. Identification of a specific telomere terminal transferase activity in *Tetrahymena* extracts. *Cell* **43**: 405-413.
- Grigorev K, Foox J, Bezdán D, Butler D, Luxton JJ, Reed J, McKenna MJ, Taylor L, George KA, Meydan C et al. 2021. Haplotype diversity and sequence heterogeneity of human telomeres. *Genome Res* doi:10.1101/gr.274639.120.
- Hardy CF, Sussel L, Shore D. 1992. A RAP1-interacting protein involved in transcriptional silencing and telomere length regulation. *Genes Dev* **6**: 801-814.
- Henning KA, Moskowitz N, Ashlock MA, Liu PP. 1998. Humanizing the yeast telomerase template. *Proc Natl Acad Sci U S A* **95**: 5667-5671.
- Ijima A, Greider CW. 2003. Short telomeres induce a DNA damage response in *Saccharomyces cerevisiae*. *Mol Biol Cell* **14**: 987-1001.
- Kaizer H, Connelly CJ, Bettridge K, Viggiani C, Greider CW. 2015. Regulation of Telomere Length Requires a Conserved N-Terminal Domain of Rif2 in *Saccharomyces cerevisiae*. *Genetics* **201**: 573-586.
- Kassambara A. 2021. rstatix: Pipe-Friendly Framework for Basic Statistical Tests.
- Keener R, Connelly CJ, Greider CW. 2019. Tel1 Activation by the MRX Complex Is Sufficient for Telomere Length Regulation but Not for the DNA Damage Response in *Saccharomyces cerevisiae*. *Genetics* **213**: 1271-1288.
- Kovaka S, Fan Y, Ni B, Timp W, Schatz MC. 2021. Targeted nanopore sequencing by real-time mapping of raw electrical signal with UNCALLED. *Nat Biotechnol* **39**: 431-441.
- Laterreur N, Lemieux B, Neumann H, Berger-Dancuse JC, Lafontaine D, Wellinger RJ. 2018. The yeast telomerase module for telomere recruitment requires a specific RNA architecture. *RNA* **24**: 1067-1079.
- Lebel C, Rosonina E, Sealey DC, Pryde F, Lydall D, Maringele L, Harrington LA. 2009. Telomere maintenance and survival in *saccharomyces cerevisiae* in the absence of telomerase and RAD52. *Genetics* **182**: 671-684.
- Lendvay TS, Morris DK, Sah J, Balasubramanian B, Lundblad V. 1996. Senescence mutants of *Saccharomyces cerevisiae* with a defect in telomere replication identify three additional EST genes. *Genetics* **144**: 1399-1412.
- Levy MZ, Allsopp RC, Futcher AB, Greider CW, Harley CB. 1992. Telomere end-replication problem and cell aging. *J Mol Biol* **225**: 951-960.
- Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**: 3094-3100.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, Genome Project Data Processing S. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078-2079.
- Louis EJ, Haber JE. 1990. Mitotic recombination among subtelomeric Y' repeats in *Saccharomyces cerevisiae*. *Genetics* **124**: 547-559.

- Louis EJ, Haber JE. 1992. The structure and evolution of subtelomeric Y' repeats in *Saccharomyces cerevisiae*. *Genetics* **131**: 559-574.
- Lundblad V, Blackburn EH. 1993. An alternative pathway for yeast telomere maintenance rescues est1-senescence. *Cell* **73**: 347-360.
- Lundblad V, Szostak JW. 1989. A mutant with a defect in telomere elongation leads to senescence in yeast. *Cell* **57**: 633-643.
- Lycka M, Peska V, Demko M, Spyroglou I, Kilar A, Fajkus J, Fojtova M. 2021. WALTER: an easy way to online evaluate telomere lengths from terminal restriction fragment analysis. *BMC Bioinformatics* **22**: 145.
- Maciejowski J, de Lange T. 2017. Telomeres in cancer: tumour suppression and genome instability. *Nat Rev Mol Cell Biol* **18**: 175-186.
- Marcand S, Brevet V, Gilson E. 1999. Progressive cis-inhibition of telomerase upon telomere elongation. *Embo J* **18**: 3509-3519.
- Marcand S, Gilson E, Shore D. 1997. A protein-counting mechanism for telomere length regulation in yeast. *Science* **275**: 986-990.
- Matheson K, Parsons L, Gammie A. 2017. Whole-Genome Sequence and Variant Analysis of W303, a Widely-Used Strain of *Saccharomyces cerevisiae*. *G3 (Bethesda)* **7**: 2219-2226.
- Maxwell PH, Coombes C, Kenny AE, Lawler JF, Boeke JD, Curcio MJ. 2004. Ty1 mobilizes subtelomeric Y' elements in telomerase-negative *Saccharomyces cerevisiae* survivors. *Mol Cell Biol* **24**: 9887-9898.
- McEachern MJ, Haber JE. 2006. Break-induced replication and recombinational telomere elongation in yeast. *Annu Rev Biochem* **75**: 111-135.
- McNally EJ, Luncsford PJ, Armanios M. 2019. Long telomeres and cancer risk: the price of cellular immortality. *J Clin Invest* **130**: 3474-3481.
- Moyzis RK, Buckingham JM, Cram LS, Dani M, Deaven LL, Jones MD, Meyne J, Ratliff RL, Wu J-R. 1988. A highly conserved repetitive DNA sequence, (TTAGGG)_n, present at the telomeres of human chromosomes. *Proc Natl Acad Sci USA* **85**: 6622-6626.
- Noma KI. 2017. The Yeast Genomes in Three Dimensions: Mechanisms and Functions. *Annu Rev Genet* **51**: 23-44.
- Pallmann J, Hothorn LA. 2016. Analysis of means: a generalized approach using R. *Journal of Applied Statistics* doi:10.1080/02664763.2015.1117584.
- Pallmann P. 2017. ANOM. Vol v0.2.
- Paschini M, Reyes CM, Gillespie AE, Lewis KA, Glustrom LW, Sharpee TO, Wuttke DS, Lundblad V. 2020. Spontaneous replication fork collapse regulates telomere length homeostasis in wild type cells. *bioRxiv*.
- Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841-842.
- Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP. 2011. Integrative genomics viewer. *Nat Biotechnol* **29**: 24-26.
- Shampay J, Blackburn EH. 1988. Generation of telomere-length heterogeneity in *Saccharomyces cerevisiae*. *Proc Nat Acad Sci USA* **85**: 534-538.
- Shen W, Le S, Li Y, Hu F. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. *PLoS One* **11**: e0163962.
- Shubin CB, Mayangsari R, Swett AD, Greider CW. 2021. Rif1 regulates telomere length through conserved HEAT repeats. *Nucleic Acids Res* **49**: 3967-3980.
- Smogorzewska A, de Lange T. 2004. Regulation of telomerase by telomeric proteins. *Annu Rev Biochem* **73**: 177-208.
- Sobecki M, Souaid C, Boulay J, Guerineau V, Noordermeer D, Crabbe L. 2018. MadID, a Versatile Approach to Map Protein-DNA Interactions, Highlights Telomere-Nuclear Envelope Contact Sites in Human Cells. *Cell reports* **25**: 2891-2903 e2895.

- Stanley SE, Armanios M. 2015. The short and long telomere syndromes: paired paradigms for molecular medicine. *Curr Opin Genet Dev* **33**: 1-9.
- Team RC. 2020. R: A Language and Environment for Statistical Computing.
- Teixeira MT, Armeric M, Sperisen P, Lingner J. 2004. Telomere length homeostasis is achieved via a switch between telomerase- extendible and -nonextendible states. *Cell* **117**: 323-335.
- Teng SC, Zakian VA. 1999. Telomere-telomere recombination is an efficient bypass pathway for telomere maintenance in *Saccharomyces cerevisiae*. *Mol Cell Biol* **19**: 8083-8093.
- Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV): high-performance genomics data visualization and exploration. *Brief Bioinform* **14**: 178-192.
- Wahlin J, Cohn M. 2000. *Saccharomyces cerevisiae* RAP1 binds to telomeric sequences with spatial flexibility. *Nucleic Acids Res* **28**: 2292-2301.
- Watson JM, Riha K. 2010. Comparative biology of telomeres: where plants stand. *FEBS Lett* **584**: 3752-3759.
- Wellinger RJ, Zakian VA. 2012. Everything you ever wanted to know about *Saccharomyces cerevisiae* telomeres: beginning to end. *Genetics* **191**: 1073-1105.
- Wichman H, Navarro D, Pendersen TL. 2016. *ggplot2: Elegant Graphics for Data Analysis*. Springer.
- Xu Z, Duc KD, Holcman D, Teixeira MT. 2013. The length of the shortest telomere as the major determinant of the onset of replicative senescence. *Genetics* **194**: 847-857.

Figure 1.





Figure 2.




Figure 3.




Figure 4.






Figure 5.


A *est2Δ*



B *est2Δrad52Δ*



C *est2Δ*



D *est2Δrad52Δ*






Figure 6.

A *est2*Δ



B *est2*Δ*rad52*Δ





Table 1.							
<i>WT</i>		<i>est2Δ P1</i>		<i>est2Δ P2</i>		<i>est2Δ P3</i>	
Bottom 99% of telomere length							
13L	77.25	10L	93.2	10L	79	10L	77.4
7R	96.06	11L	70.4	11L	71.6	11L	69.4
8R	96.19	2R	72	2R	75.6	8L	85
<i>tell1Δ</i>		<i>est2rad52Δ P1</i>		<i>est2rad52Δ P2</i>		<i>est2rad52Δ P3</i>	
Bottom 99% of telomere length							
13L	69.3	10L	86.45	10L	90.4	10L	76.75
4R	73.4	1L	89.6	1L	81.7	1L	71.9
8R	69.4	3L	171	2R	85	2R	71.95