



Accurate and robust inference of microbial growth dynamics from metagenomic sequencing reveals personalized growth rates

Tyler A Joseph, Philippe Chlenski, Aviya Litman, et al.

Genome Res. published online January 5, 2022

Access the most recent version at doi:[10.1101/gr.275533.121](https://doi.org/10.1101/gr.275533.121)

P<P Published online January 5, 2022 in advance of the print journal.

Accepted Manuscript Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.

Creative Commons License This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <https://genome.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).



To subscribe to *Genome Research* go to:
<https://genome.cshlp.org/subscriptions>

Published by Cold Spring Harbor Laboratory Press

Accurate and robust inference of microbial growth dynamics from metagenomic sequencing reveals personalized growth rates

Tyler A. Joseph¹, Philippe Chlenski¹, Aviya Litman², Tal Korem^{2,3,4*}, and Itsik Pe'er^{1,2,5*}

¹Department of Computer Science, Columbia University, New York, NY, USA

²Department of Systems Biology, Columbia University Irving Medical Center, New York, NY, USA

³Department of Obstetrics and Gynecology, Columbia University Irving Medical Center, New York, NY, USA

⁴CIFAR Azrieli Global Scholars Program, CIFAR, Toronto, Ontario, Canada

⁵Data Science Institute, Columbia University, New York, NY, USA

*These authors contributed equally to this work.

Correspondence: tal.korem@columbia.edu, itsik@cs.columbia.edu.

Running Title: Inference of microbial growth dynamics

Abstract

Patterns of sequencing coverage along a bacterial genome—summarized by a peak-to-trough ratio (PTR)—have been shown to accurately reflect microbial growth rates, revealing a new facet of microbial dynamics and host-microbe interactions. Here, we introduce CoPTR (Compute PTR): a tool for computing PTRs from complete reference genomes and assemblies. Using simulations and data from growth experiments in simple and complex communities, we show that CoPTR is more accurate than the current state-of-the-art, while also providing more PTR estimates overall. We further develop theory formalizing a biological interpretation for PTRs. Using a reference database of 2935 species, we applied CoPTR to a case-control study of 1304 metagenomic samples from 106 individuals with inflammatory bowel disease. We show that growth rates are personalized, are only loosely correlated with relative abundances, and are associated with disease status. We conclude by demonstrating how PTRs can be combined with relative abundances and metabolomics to investigate their effect on the microbiome.

Introduction

Dynamic changes in the human microbiome play a fundamental role in our health. Understanding how and why these changes occur can help uncover mechanisms of disease. In line with this goal, the Integrative Human Microbiome Project and others have generated longitudinal datasets from disease cohorts where the microbiome has been observed to play a role (Buffie et al., 2015; DiGiulio et al., 2015; Lloyd-Price et al., 2019; Zhou et al., 2019; Serrano et al., 2019). Yet, investigating microbiome dynamics is challenging. On one hand, a promising line of investigation uses time-series or dynamical systems based models to investigate community dynamics (Stein et al., 2013; Bucci et al., 2016; Gibbons et al., 2017; Gibson and Gerber, 2018; Shenhav et al., 2019; Joseph et al., 2020). On the other hand, the resolution of such methods is limited by sampling frequency, which is often limited by physiological constraints on sample collection for DNA sequencing. Furthermore, while such methods accurately infer changes in abundance, they do not directly assess growth rates per sample.

Korem et al. (2015) introduced a complementary approach to investigate microbiome dynamics. They demonstrated that sequencing coverage of a given species in a metagenomic sample reflects its growth rate. They summarized growth rates by a metric called the peak-to-trough ratio (PTR): the ratio of sequencing coverage near the replication origin and near the replication terminus. Thus, PTRs provide a snapshot of growth at the time of sampling, and their resolution is not limited by sampling frequency.

Their original method—PTRC—estimates PTRs using reads mapped to complete reference genomes. It has been used as a gold standard to evaluate other methods (Brown et al., 2016; Emiola and Oh, 2018; Gao and Li, 2018). However, most species lack complete reference genomes, reducing PTRC's utility to researchers in the field. Therefore, follow-up work has focused on estimating PTRs from draft assemblies: short sections of contiguous sequences (contigs), where the order of contigs along the genome is unknown. These approaches rely on reordering binned read counts or contigs by estimating their distance to the replication origin. Although less accurate than PTRC, they allow PTRs to be estimated for a larger number of species. iRep (Brown et al., 2016) sorts binned read counts along a 5kb sliding window, then fits a

1 log-linear model to the sorted bins to estimate a PTR. GRiD (Emiola and Oh, 2018) sorts the contigs
2 themselves by sequencing coverage. It fits a curve to the log sequencing coverage of the sorted contigs using
3 Tukey's biweight function. DEMIC (Gao and Li, 2018) also sorts contigs. However, it uses sequencing
4 coverage across multiple samples to infer a contig's distance from the replication origin. Specifically,
5 DEMIC performs a principal component analysis on the log contig coverage across samples. The authors
6 demonstrate that the scores along the first principal component correlate with distance from the replication
7 origin. Ma et al. (2020) provide theoretical criteria for when such an approach is optimal. Finally, other
8 estimators have focused on PTR estimation for specific strains (Emiola et al., 2020), or estimation using
9 circular statistics (Suzuki and Yamada, 2020).

10 Nonetheless, using PTRs has several limitations. From a theoretical perspective, it is not clear what
11 PTRs estimate and how they should be interpreted. Bremer and Churchward (1977) demonstrated that under
12 exponential growth, PTRs measure the ratio of chromosome replication time to generation time, but this is
13 not established under arbitrary models of dynamics. From a practical perspective, estimating PTRs at scale
14 requires running multiple tools across multiple computational environments—a cumbersome task.

15 In the present work we seek to address these issues. Our contributions are threefold. First, we provide
16 theory that shows PTRs measure the rate of DNA synthesis and generation time, regardless of the
17 underlying dynamic model. Second, we derive two estimators for PTRs—one for complete reference
18 genomes and one for draft assemblies. Third, we combine our estimators in a easy-to-use tool called CoPTR
19 (Compute PTR). CoPTR provides extensive documentation, a tutorial, and a precomputed reference
20 databases for its users. We demonstrate that CoPTR is more accurate than the current state-of-the-art, and
21 conclude with a large-scale application to a dataset of 1304 metagenomic samples from a study of
22 inflammatory bowel disease.

Results

CoPTR Overview

The method we developed models the density of reads along the genome in a sample by adapting an argument proposed by Bremer and Churchward (1977). Under an assumption of exponential growth, they showed that the copy number ratio of replication origins to replication termini in a population, R , is given by

$$\log_2(R) = \frac{C}{\tau} \quad (1)$$

where C is the time it takes to replicate a bacterial chromosome, and τ is the (fixed) generation time. We generalize this (see Supplemental Note S1) for dynamic quantities:

$$\log_2(R(t)) = \frac{C}{\tau(t)} \quad (2)$$

The variable τ now depends on collection time t . When a complete reference genome is available the PTR is an estimator for $R(t)$. However, the PTR is only correlated with $R(t)$ on draft assemblies because the assembly may not include the replication origin or terminus. Furthermore, while PTRs correlate with the growth rates (i.e. $\frac{1}{\tau(t)}$), they only measure changes in abundance provided no cells are being removed from the community (Supplemental Note S1.4).

The derivation also suggests that copy number along the chromosome decays log-linearly away from the replication origin (Supplemental Note S2). We used this fact to develop CoPTR (Compute PTR): a maximum likelihood method for estimating PTRs from complete genomes and draft assemblies (Figure 1). CoPTR takes sequencing reads from multiple metagenomic samples and a reference database of complete and draft genomes as input. It outputs a genome-by-sample matrix where each entry is the estimated $\log_2(\text{PTR})$ for each input genome in that sample. It has two modules: CoPTR-Ref that estimates PTRs from complete genomes, and CoPTR-Contig that estimates PTRs from draft assemblies. As such, it combines the

improved accuracy enabled by complete genomes with the flexibility afforded by being able to work against draft and metagenomic assemblies.

For both methods, sequencing reads are first mapped to a reference database. CoPTR-Ref estimates PTRs by applying an adaptive filter to remove regions of ultra-high or ultra-low coverage. It then fits a probabilistic model to estimate the replication origin and the PTR. CoPTR-Contig estimates PTRs by first binning reads into approximately 500 non-overlapping windows. It filters out windows with excess or poor numbers of reads. Coverage patterns across multiple samples are used to reorder bins using Poisson PCA. The reordered bins serve as approximate genomic coordinates which are used to obtain maximum likelihood estimates of PTRs. We chose to reorder bins, rather than contigs, to be more robust to errors in the assembly process, to large changes in coverage patterns along contigs due to mobile genetic elements, and to coverage drops near the edges of contigs.

CoPTR-Ref accurately estimates PTRs using complete reference genomes

We first evaluated CoPTR-Ref on simulated data. Briefly, we simulated read counts based on read density maps generated from high coverage genomic samples of *Escherichia coli*, *Lactobacillus gasseri*, and *Enterococcus faecalis* from Korem et al. (2015) (Supplemental Figure S1). The density maps reflect differences in coverage along a genome due to GC content and mappability. To facilitate comparison with CoPTR-Ref, we also reimplemented PTRC. The new implementation, called KoremPTR, was designed to work with simulated read counts and reads mapped with Bowtie 2 (Langmead and Salzberg, 2012). KoremPTR showed a good correspondence with the original method (Pearson $r > 0.99$; Supplemental Figure S2).

Our simulations demonstrated that CoPTR-Ref requires as few as 5000 reads to achieve greater than 0.95 Pearson's correlation (Supplemental Figure S3). We confirmed our coverage requirements on real data by downsampling the number of reads from the Korem et al. (2015) *E. coli* dataset and comparing estimates from the downsampled data to the estimates from the complete data (Supplemental Figure S4). Notably, CoPTR-Ref more accurately estimated PTRs than KoremPTR (Figure 2A, Supplemental Figure S3).

KoremPTR appeared to underestimate the simulated PTR, causing the difference in accuracy (Figure 2B, Supplemental Figure S3). Nonetheless, PTR estimates by KoremPTR were highly correlated with the ground truth (Pearson $r>0.88$). We saw the same pattern across 6 genomic (bacteria grown in monoculture) and metagenomic datasets (Figure 2C). Both methods were correlated, but CoPTR-Ref estimated larger PTRs than KoremPTR on the same samples.

To evaluate whether variation among representative genomes per 95 % ANI clusters—an operational threshold for defining species (Olm et al., 2020)—affects the accuracy of CoPTR, we mapped the same samples to different strains. We found that PTR estimates were robust to strain variation when the MASH distance (Ondov et al., 2016) between strains was less than 0.05—corresponding to $\approx 95\%$ ANI (Figure 2D). These results indicate that one reference genome per 95% ANI cluster covers the space of genomes for PTR estimation.

We compared $\log_2(\text{PTR})$ estimates to growth rates of *E. coli* grown in a chemostat, and to changes in population size of *E. coli* in an unrestricted growth setting (Supplemental Figure S5) using data from Korem et al. (2015). Our theory suggests that $\log_2(\text{PTR})$ s are correlated with both quantities in these settings. We found a strong correlation ($r>0.9$) between $\log_2(\text{PTR})$ and growth rates, and a strong correlation ($r>0.9$) between $\log_2(\text{PTR})$ s and changes in abundance. We additionally replicated the growth rate experiments using the *E. faecalis* (aerobic and anaerobic growth experiments) and *L. gasseri* datasets, as in Korem et al. (2015). We found a good correlation between estimated growth rates and $\log_2(\text{PTR})$, similar to that of KoremPTR (Supplemental Table S1).

CoPTR-Contig accurately estimates PTRs using MAGs

Because CoPTR-Contig reorders bins, not contigs, we could directly compare CoPTR-Ref to CoPTR-Contig using the same simulation framework (Figure 3A, Supplemental Figure S6). Estimates by CoPTR-Contig were highly correlated (Pearson $r>0.9$) with the simulated ground truth with as few as 5000 reads, but were overall less accurate than CoPTR-Ref. Similar to CoPTR-Ref, we confirmed our coverage

requirements by downsampling sequencing reads from real data (Supplemental Figure S4). Our results highlight the benefit of using the additional information provided by complete reference genomes.

To assess the applicability of our method to metagenomic assemblies, which are of variable quality and contamination levels, we performed simulations investigating their impact on the accuracy of CoPTR-Contig. We found that CoPTR-Contig is robust to the level of genome completeness, providing comparable accuracy with completeness as low as 50%. We further found that CoPTR-Contig's estimates are robust to moderate amounts of up to 5% contamination in the assembly from other species (Supplemental Figure S7).

We then compared CoPTR-Contig to GRiD, DEMIC, and iRep across 5 real genomic and metagenomic datasets of *E. coli* and *L. gasseri* where both complete reference genomes and metagenomic assembled genomes (MAGs) were available (Figure 3B). We considered 10 high-quality MAGs (>90% completeness <5% contamination) from the IGGdb (Nayfach et al., 2019) and computed the correlation between the $\log_2(\text{PTR})$ estimate from each method and the $\log_2(\text{PTR})$ from CoPTR-Ref. For CoPTR-Ref, reads were mapped to a single complete genome (see Methods). All 10 of the *E. coli* MAGs were assigned to the same 95% ANI species cluster, while 8 of the 10 *L. gasseri* MAGs were from one cluster, and the remaining 2 from another. To allow for a fair comparison, we changed the default parameters of each method to allow estimates on each sample—with the exception of DEMIC which provides no command line options to change filtering criteria. We note that almost all the samples we explored were below the minimum recommended coverage for iRep (Figure 3C, Supplemental Figure S8).

We found that CoPTR-Contig significantly outperformed (p -value <0.05 using a two-sided paired t -test; the 2 *L. gasseri* MAGs from a different species cluster were excluded) GRiD on 3 datasets, DEMIC on 2 datasets, and iRep on all 5 datasets. All models performed poorly on the 2 *L. gasseri* MAGs that were from a different 95% ANI cluster (outliers on Figure 3B), recapitulating results from the strain comparison analysis using CoPTR-Ref (Figure 2D). Many of the comparisons between CoPTR-Contig and DEMIC failed to reach significance because DEMIC estimated fewer PTRs overall (Figure 3C, Supplemental Figure S8), resulting in fewer MAGs for comparison (points in Figure 3C). We additionally quantified the accuracy

1 across MAGs by counting the number of MAGs where the correlation between the ground truth and each
2 method was high. We found that CoPTR-Contig had high accuracy across a larger number of MAGs (30
3 MAGs with Pearson $r > 0.9$ for CoPTR-Contig, compared to 19, 11, and 0 for DEMIC, GRiD and iREP,
4 respectively; Supplemental Figure S9).

5 An important aspect affecting the utility of PTR inference methods is the number of PTR estimates they
6 are able to provide for a given sample. We therefore compared the number of estimated PTRs that passed
7 the default filtering criteria of each method (Figure 3C, Supplementary Figure S8). We mapped 10 samples
8 from the IBD dataset (Methods) to 1,009 high-quality MAGs from the IGGdb, and counted the number of
9 PTR estimates. The reported estimates for GRiD are based on GRiD's published minimum coverage
10 requirement: species with $>0.2x$ sequencing coverage. We were unable to run GRiD's high-throughput
11 model on two systems (Ubuntu 18.04.4 LTS and macOS 10.15) to produce estimates on this dataset. We
12 found that CoPTR-Contig produced more PTR estimates overall than the other models we evaluated. We
13 note that this number does not include the additional estimates from complete genomes using CoPTR-Ref.
14 Taken together with the improved accuracy of CoPTR (Figure 3B), these results show that CoPTR
15 outcompetes previous PTR estimation methods in both the number of estimates produced and their
16 accuracy, demonstrating its utility for microbiome analysis.

CoPTR accurately predicts *in situ* growth in a marine microbial population

To evaluate the accuracy of CoPTR in complex natural communities, we used data from a recent study that benchmarked the accuracy of PTRs in estimating *in situ* growth rates of marine bacteria collected from seawater. Long et al. (2021) conducted five growth experiments, each with 4-5 samples collected over 42-44 hours. They used metagenomic sequencing to assemble MAGs and calculate PTRs, and combined total cell counts with relative abundances to estimate absolute abundances and calculate growth rates. The study concluded that PTRs rarely correlated to growth in most marine bacterial populations.

We estimated PTRs for the MAGs provided by Long et al. (2021) using CoPTR (Supplemental Table S2), and compared them to growth rates calculated from absolute abundance estimates (Methods). This is a challenging benchmark, due to low sampling frequency (average of 12 hours between samples), challenges in estimating growth rates where cell death likely plays a role (417 of 1818 calculated growth rates were negative) and low MAG quality (median [IQR] completeness of 66.9% [56.8-84.2%] and redundancy of 2.9% [2.2-4.3%]). Nevertheless, CoPTR showed reasonable accuracy (Figure 4A-B; median [IQR] Pearson r of 0.40 [-0.10-0.64]), with $r > 0.3$ for 29 of 47 MAGs, and high accuracy ($r > 0.6$) for 14 of them. Out of the 9 MAGs with available measurements and completeness $> 90\%$ (Supplemental Figure S10), 7 had positive correlations, with a median [IQR] Pearson r of 0.45 [0.22-0.72], indicating a potential to improve the overall accuracy of estimates with MAGs of higher quality.

We next compared the accuracy of CoPTR to iRep, GRiD and DEMIC. CoPTR outperformed all three methods (two-sided Mann-Whitney U $p = 3.2 \times 10^{-4}$, $p = 0.0047$ and $p = 0.016$ for DEMIC, GRiD and iRep, respectively; Figure 4C). These results again demonstrate the improved accuracy of CoPTR even in this challenging setting.

PTRs recapitulate a signal of antibiotic resistance

We next evaluated if we could use CoPTR to detect a signal of antibiotic resistance in *Citrobacter rodentium*. Korem et al. (2015) generated 86 samples from 3 populations of *in vitro* culture of *C. rodentium*. One population was treated with Erythromycin, a growth inhibiting antibiotic; another was treated with Nalidixic acid to which *C. rodentium* is resistant. The final population was a control and received no treatment.

We wanted to see if we could recapitulate this signal using CoPTR. Similar to the original study, we observed a difference in PTRs between the populations exposed to Erythromycin and Nalidixic acid (Supplemental Table S3). Our results add to the original study by assigning an effect size to each condition. We found that Erythromycin has a strong negative effect size on the $\log_2(\text{PTR})$, while Nalidixic acid has a strong positive effect size. Our results suggest that *C. rodentium* has an increased growth rate in response to Nalidixic acid. However, this decrease did not correspond to an increased rate of population growth ($p=0.706$, two-sided *t*-test).

PTRs are highly personalized

We next sought to demonstrate how PTR measurements can be used in a large-scale study. To this end, we considered 1304 metagenomic samples from 106 individuals in a case-control study of inflammatory bowel disease (IBD) (Lloyd-Price et al., 2019). Individuals in the study had two different subtypes of IBD: Crohn's disease and Ulcerative colitis. We mapped the metagenomic samples to a database from IGGdb (Nayfach et al., 2019) consisting of 2935 complete genomes, assemblies, and MAGs, selected as representative genomes from 95% ANI clusters (Methods). Individuals had between 3 and 23 associated metagenomic samples each, with a median of 11 samples (Supplemental Figure 11A). PTR estimates were sparse among species (Supplemental Figure 11B). Of the species that had at least 1 observed PTR, the median number of PTR estimates was 28; Approximately 20% of species had fewer than 9 observed PTRs.

A large dataset with multiple samples per individual allowed us to investigate sources of variation for PTRs. To this end, we estimated the fraction of variation explained by differences between individuals,

disease-status, age, and sex (Methods). Inter-individual differences in PTRs accounted for the largest fraction of variance among variables explored (Figure 5A, B), consistent with the original study that found inter-individual variation to be largest source of variation among the other multi-omic measure types collected (Lloyd-Price et al., 2019). We repeated the experiment using the top 50% of individuals with the most observations and found similar results (Supplemental Figure S12). PTRs were mostly uncorrelated with relative abundances, suggesting that PTRs tag a signal of biological variation complementary to relative abundances (Figure 5C).

PTRs are associated with IBD

We then asked if we could associate species to disease status through their PTRs (Methods; Figure 6A). We found 1 species that was significantly associated (FDR $q=0.025$, effect size $=-0.1574$) with Crohn's disease (Supplemental Table S4), *Subdoligranulum* sp., and three species with Ulcerative colitis (Supplemental Table S5): *Roseburia intestinalis* ($q = 1.07 \times 10^{-3}$, effect size = 0.094), *Ruminiclostridium* sp ($q = 2.5 \times 10^{-2}$, effect size = -0.138), and *Subdoligranum* sp. ($q = 2.69 \times 10^{-2}$, effect size = -0.168). Vila et al. (2018) also report an increased PTR in *R. intestinalis* in individuals with Crohn's disease and Ulcerative colitis in a separate cohort, using PTRC. We did not observe a significant association between the relative abundance of *R. intestinalis* and disease status, nor did Vila et al. (2018). Altogether, our results provide additional evidence that *R. intestinalis* may play a role in Ulcerative colitis, observable only through analysis of growth dynamics.

For the remaining investigation we focused on *R. intestinalis*. We asked if we could assess the impact of various species on *R. intestinalis* by associating relative abundances across species estimated with MetaPhlAn2 (Truong et al., 2015) with its $\log_2(\text{PTR})$ (Supplementary Table S6). We found two species with a positive association with *R. intestinalis*, and one with a strong negative association (Figure 6B). Finally, we investigated if we could relate metabolomic measurements to $\log_2(\text{PTR})$ s (Supplemental Table S7). We found two metabolites with a positive association with the $\log_2(\text{PTR})$ of *R. intestinalis* (Figure 6C).

One of them—2-hydroxyglutarate—is part of the butanoate metabolic pathway, and *R. intestinalis* is a known butyrate producing bacteria. Altogether, these results demonstrate the utility of PTRs for integrating multi-omic measurements with metagenomic datasets.

Discussion

Peak-to-trough ratios (PTRs) have the potential to be a valuable tool for investigating microbiome dynamics. Here, we provided theory giving PTRs a biological interpretation. We showed that PTRs are correlated with growth rates defined as the reciprocal of generation time, but not always correlated with changes in population size. We introduced CoPTR, a software system combining two methods for estimating PTRs: CoPTR-Ref estimates PTRs with the assistance of a complete reference genome, and CoPTR-Contig estimates PTRs from draft assemblies. CoPTR is easy to use, has extensive documentation, and provides a precomputed reference database for its users.

We showed that CoPTR-Ref is more accurate than KoremPTR, the current gold standard for PTR estimation from complete reference genomes. The difference in performance is likely driven by two key differences between the models of each method. First, KoremPTR applies a strong smoother to binned read counts that reduces the variance in read counts among adjacent bins. The smoother clips read counts near the replication origin and terminus, causing KoremPTR to underestimate PTRs. CoPTR-Ref does not apply a smoother, but instead filters out regions of excess or poor coverage. Second, KoremPTR estimates PTRs by taking the ratio of read counts between the replication origin and terminus. In contrast, CoPTR-Ref uses a probabilistic model to take a maximum likelihood estimate of the PTR under a parametric model.

We also showed that CoPTR-Contig was more accurate than the current state-of-the-art for PTR estimation using draft assemblies, while providing more PTR estimates overall. A potential limitation of CoPTR-Contig is its reliance on multiple samples to reorder binned read counts along the genome. However, this is not a severe limitation. Our simulations show that as few as 5 samples are required to

1 estimate a PTR. Furthermore, since PTRs are not comparable across species, multiple PTRs per species are
 2 required to reach reasonable sample sizes for statistical testing. Thus, for most purposes, the sample
 3 requirement is negligible.

4 When building CoPTR we focused on estimating PTRs per species, rather than per strain. Our goal was
 5 to allow CoPTR to be applied to recent database efforts that combined representative genomes from MAGs,
 6 assemblies, and complete genomes clustered at approximately 95% average nucleotide identity (Almeida
 7 et al., 2019, 2021; Forster et al., 2019; Nayfach et al., 2019; Pasolli et al., 2019; Zou et al., 2019). There are
 8 benefits and drawbacks to this approach. The major benefit is reduction in database size, and therefore in
 9 computational time required for read mapping. Our results showed that PTR estimates from the same
 10 samples mapped to different closely related strains were highly concordant. Thus, there is not much to be
 11 gained from including all strains in the reference database. Nonetheless, the drawback is that CoPTR may
 12 not distinguish differences in PTRs across samples due to differences in strains.

13 We also focused on estimating PTRs from high-quality MAGs (>90% completeness, <5%
 14 contamination). Inference from MAGs is more challenging than other assembly types, due to differences in
 15 assembly completeness and contamination from other species. Many things can go wrong during the
 16 assembly processes. These, in turn, can affect PTR inference. In our opinion, it is better to have fewer
 17 high-quality estimates than more poor-quality ones, and for this reason we have chosen strict inclusion
 18 criteria for MAGs. Nevertheless, even in a complex metagenomic benchmark of oceanic microbes with
 19 lower-quality MAGs, we show that the accuracy of CoPTR significantly outperform other PTR inference
 20 methods. Contrary to previous claims (Long et al., 2021), we demonstrate that PTRs are able to provide
 21 reasonably accurate estimates of *in situ* growth even in this slow-growing, naturally-occurring ecosystem.

22 Our results on the IBD dataset showed that PTRs were highly personalized. Indeed, a large fraction of
 23 variance in PTRs was explained by inter-individual variation. To our knowledge, we are the first study to
 24 show an individualized effect for PTRs. Because of their close connection to growth rates, our results
 25 suggest that species' growth rates are individual-specific and somewhat stable in healthy individuals. We
 26 also showed that some species display differences in PTRs depending on disease status. It would be
 27 interesting to test whether this is a systematic difference in PTRs between cases and controls, or whether

PTRs deviate from a stable baseline during active periods of disease. We could not test this hypothesis here because the metadata do not indicate disease severity at time of sampling. Nonetheless, future work should investigate this further.

There are other benefits to using PTRs as well. Compared to relative abundances, PTRs have a clearer biological interpretation, because an increase in relative abundance does not necessarily correspond to an increase in population size. In contrast, we showed that an increase in PTR in a species corresponds to an increase in the rate of DNA synthesis, and that an increase in the log PTR corresponds to a decrease in generation time. Either of these facts can be used to generate hypotheses about the drivers of differences across conditions. Furthermore, because PTRs provide a snapshot of growth at the time of sampling, they potentially alleviate the need to perform dense-in-time sampling typically needed to detect dynamic changes. This suggests that it may be more cost-effective to sequence more individuals, rather than more samples per individual. Finally, we showed that relative abundances and metabolomic profiles can be used to associate species or metabolites with PTRs. Altogether, our study demonstrates that PTRs can provide new approaches for investigating community interactions, relating multiomic measurements to the microbiome, and for investigating the relationship between microbiome dynamics and disease.

Methods

CoPTR Implementation

Read mapping. Reads are mapped using Bowtie 2 (Langmead and Salzberg, 2012) using the parameter `-k 10` to allow up to 10 mappings per read. We chose this parameter after observing that 99% of reads mapped to 10 or fewer locations on the IGGdb using a subset of 10 samples from the IBD dataset. Reads with fewer than 10 mapping were assigned using a variational inference algorithm described in Supplemental Note 3. We chose variational inference to reassign multi-mapped reads, rather than expectation maximization,

because it provides a more flexible framework for possible extensions to our method. In present work, reads with 10 (or more) mappings were discarded from downstream analysis. However, CoPTR has a command line argument to adjust this setting.

Before reassigning multiply mapped reads, reads are filtered by alignment score. Alignment score is more sensitive than mapping quality, since different alignment scores can result in the same mapping quality. Bowtie 2 assigns penalties to mismatched bases weighted on their base quality score. Bases with a perfect quality score receive a -6 penalty for a mismatch, decreasing as the quality score decreases. For a read of length L , we filtered out reads with a score less than $-6 \times L \times 0.05$. Given a read with perfect quality scores, this corresponding to removing reads with less than 95% identity to the reference sequence. Of course reads do not have perfect quality scores, so this threshold is less strict than 95% identity.

CoPTR-Ref. PTRs from species with complete reference genomes are estimated with CoPTR-Ref. Regions of the genome with excess or poor coverage per sample are first filtered out in two steps. In the first step we apply a coarse-grained filter by binning reads into 500 bins. Let m be the median $\log_2$ read count across nonzero bins, and s the larger of 1 or the standard deviation of the nonzero $\log_2$ read counts. Bins are filtered out if they fall outside the interval $(m - \alpha_{0.025}, m + \alpha_{0.025})$, where $\alpha_{0.025}$ is the two-sided (1-0.025) critical region from an $N(m,s)$ distribution. After the coarse-grained filter, we apply a fine-grained filter by computing read counts across a rolling window encompassing 12.5% of the genome. We apply the same filtering criteria around the center of each window. After filtering, quality is assessed per genome per sample. No estimate is produced if a genome has fewer than 5000 reads, or if more than 25% of binned read counts are zero.

Bins in genomes from the remaining samples are concatenated, and read positions normalized so that they fall in the unit interval $[0,1]$. Let $x \in [0,1]$ be the coordinate of a read, x_i be the coordinate of the replication origin, and $x_i = (x_i + 0.5) \bmod 1$ be the replication terminus. We estimate the $\log_2(\text{PTR})$ and replication origin across all samples by maximizing the likelihood of the model

$$\alpha = \frac{\log_2 r}{x_i - x_t} = \frac{\log_2 p(x_i) - \log_2 p(x_t)}{x_i - x_t}$$

$$\begin{aligned}
x_1 &= \min\{x_i, x_t\} \\
x_2 &= \max\{x_i, x_t\} \\
c(x) &= \begin{cases} \log_2 p(x_i) & \text{if } x = x_i \\ \log_2 p(x_t) & \text{if } x = x_t \end{cases} \\
\log_2 p(x) &= \begin{cases} -\alpha(x - x_1) + c(x_1) & \text{if } x \leq x_1 \\ \alpha(x - x_1) + c(x_1) & \text{if } x_1 < x < x_2 \\ -\alpha(x - x_2) + c(x_2) & \text{if } x \geq x_2 \end{cases}
\end{aligned}
\tag{3}$$

We describe how to compute $\log_2 p(x_i)$, $\log_2 p(x_t)$, and the normalizing constant in Supplemental Note S2. We maximize the likelihood using the SLSQP optimizer in SciPy (Virtanen et al., 2020). We first maximize with respect to each sample separately to get initial estimates of the $\log_2(\text{PTR})$ per sample, then jointly estimate the replication origin given these estimates. Finally, given the estimated replication origin from all samples, each individual $\log_2(\text{PTR})$ is updated once more.

CoPTR-Contig. PTRs from species with draft assemblies are estimated with CoPTR-Contig. Reads across contigs are binned into approximately 500 bins (adjusted such that the average length of each bin is divisible by 100bp). We choose 500 bins, rather than fixed bin size, so that the model would behave similarly across genomes of different lengths. We then apply a similar coarse-grained filter to the $\log_2$ read counts binned into 500 bins. Bins that are filtered are marked as missing for the Poisson PCA step. Genomes in samples with more than 50% missing bins, with fewer than 5000 reads after filtering, with fewer than total 50 bins, or observed in fewer than 5 samples are excluded.

The remaining bins are reordered by applying a Poisson PCA to read counts across samples. Let B be the number of bins, and N the number samples. Let x_{bi} be the read count in bin b from sample i , and let $\Omega = \{x_{bi} : \text{bin } b \text{ is not missing from sample } i\}$. In Poisson PCA, we model the read count in each bin using a matrix $C \approx \exp\{WV\}$, $W \in \mathbb{R}^{B \times k}$, $V \in \mathbb{R}^{k \times N}$ with low-rank structure. Specifically, we assume rank 1 structure where $W \in \mathbb{R}^{B \times 1}$ and $V \in \mathbb{R}^{1 \times N}$. The read count x_{bi} is modeled by

$$x_{bi} \sim \text{Poisson}(\exp(w_b v_i))$$

The parameters W and V are estimated by iteratively maximizing the likelihood

$$L(W, V) = \sum_{(b,i) \in \Omega} \log p(x_{bi}; W, V) \quad (5)$$

with respect to W then V until convergence.

The scores for each bin w_b are used to reorder bins based on their rank, representing approximate distance from the replication origin. After reordering each the top and bottom 5% of bins removed in each sample. The $\log_2(\text{PTR})$ is estimated by maximizing a discretized version of equation 3 using the SLSQP optimizing in SciPy, fixing the replication origin at one end and terminus at the other.

Simulations

To generate realistic simulations we computed read density maps by mapping reads from genomic (monoculture) samples to reference genomes where the strain was known. For each density map, we computed the read count in 100bp bins, then divided by the total number of reads to obtain empirical probabilities that a read originates from a location in the genome. These probabilities are conditioned on the PTR in the sample. We therefore used KoremPTR to estimate the PTR for each sample using the replication origin from the DoriC database (Luo and Gao, 2019), and reweighted the probabilities by the estimated PTR. Specifically, let $p_1, \dots, p_N$ be the unadjusted probabilities that a read originates from a bin, let $\tilde{p}_1, \dots, \tilde{p}_N$ be the probabilities under the model given the replication origin and PTR, and let $\hat{p}_1, \dots, \hat{p}_N$ be the adjusted probabilities. The adjusted probabilities are

$$\log_2 \hat{p}_i = \log_2 p_i - \log_2 \tilde{p}_i + N \quad (6)$$

where N is the normalizing constant.

We generated density maps for *E. coli* from a genomic sample with 894,685 reads (14x coverage), a *L. gasseri* sample with 2,645,206 reads (104x coverage), and *E. faecalis* with 581,836 reads (14.75x

coverage) from Korem et al. (2015). Supplemental Figure S1 displays the adjusted density maps. When simulating data, we performed the reversed adjustment by the simulated replication origin and PTR. Given $\hat{p}_1, \dots, \hat{p}_N$, and theoretical probabilities for the simulated PTR and replication origin $\bar{p}_1, \dots, \bar{p}_N$, we computed the probability that a read is derived from bin i by computing

$$\log_2 p_i = \log_2 \hat{p}_i + \log_2 \bar{p}_i + N$$

To compare CoPTR-Ref and KoremPTR, we performed 100 simulations each for read counts of 1000, 2500, 5000, 10000, and 20000. For each simulation, a random replication origin and PTR is chosen. Reads counts in 100 bp bins are simulated based on the adjusted probabilities described above, then converted to genomics coordinates. The coordinates are provided to CoPTR-Ref and KoremPTR to estimate PTRs.

To evaluate CoPTR-Contig, we performed 20 simulation replicates consisting of 100 samples each, while varying the number of simulated reads. Because PTR estimates can be sparse, we processed samples in batches of 5 to explore how well CoPTR-Contig reordered bins at small sample sizes.

Completeness and contamination experiments. We extended our simulation framework to investigate genome completeness and contamination. To simulate genome completeness, we held out random fragments of the *E. coli* density map in 1% increments selected uniformly at random. The remaining sections of the genome were treated as contigs, and reads were simulated from the contigs. To simulate genome contamination, we simulated reads from two separate genomes: *E. coli* and *L. gasseri*. For a given contamination percentage c , reads were simulated from the *E. coli* genome, setting the completeness percentage to $100-c$. Then, simulated read counts from contigs in *L. gasseri* genome were added until the percentage of contamination by *L. gasseri* was c .

Datasets and reference genomes for benchmarking experiments

We downloaded genomic samples from Korem et al. (2015), and metagenomic samples from the Human Microbiome Project (Lloyd-Price et al., 2017), BioProject PRJNA275349, and the IBD dataset (Lloyd-Price et al., 2019). Vaginal and gut metagenomic samples from the Human Microbiome Project

were selected by mapping reads to reference genomes of *E. coli* and *L. gasseri*, and retaining samples with more than 2500 mapped reads. Gut samples of *L. gasseri* from the IBD dataset were selected based on whether CoPTR had an estimated PTR. Complete accession numbers per experiment are listed in Supplementary Table S8.

To compare estimates across reference genomes, we downloaded reference genomes from NCBI. Accession numbers for genomes and MAGs are listed in Supplemental Table S8. We selected genomes from each of *E. coli*, *L. gasseri*, and *E. faecalis* matching the strains reported by Korem et al. (2015), and performed comparison on genomic samples using these strains. The genomes NC_007779.1, NC_008530.1, and NZ_CP008816.1 correspond to the strains used by Korem et al. (2015). Distances between reference genomes were computed using MASH v2.2 (Ondov et al., 2016).

To compare estimates across MAGs, we downloaded high-quality assemblies from Nayfach et al. (2019). On both complete references and MAGs, we noted that the *L. gasseri* genomes were from two different 95% ANI species clusters: 8 MAGs were from one cluster and 2 MAGs were from the another. To compare PTR estimates from *L. gasseri* MAGs to CoPTR-Ref estimates, we selected a reference genome corresponding to the species cluster with 8 MAGs. We did this by downloading a complete genome in the same species cluster identified by Nayfach et al. (2019), and computing the MASH distance with genomes above. We found one genome with 0 MASH distance to the species cluster which we used for analysis.

To perform the model comparison experiments and the *C. rodentium* experiments, we mapped reads to one genome at a time using Bowtie 2's default parameters, and provided these as input to CoPTR. We used CoPTR's default settings to assess metagenomic samples for quality.

Downsampling experiments

We additionally assessed read count requirements for CoPTR-Ref and CoPTR-Contig using the *E. coli* - *genomic* dataset. Using samples with more than 20000 mapped reads, we computed the Pearson's correlation between $\log_2(\text{PTR})$ s estimated from all reads to $\log_2$ computed by downsampling 2500, 5000,

10000, and 20000 reads respectively. To assess CoPTR-Contig, we downloaded an *E. coli* assembly corresponding to the strain from Korem et al. (2015) from NCBI (NZ_JAGGIP010000010.1).

4 Growth rate and abundance change experiments

We downloaded additional data from Korem et al. (2015) corresponding to the *E. coli* unrestricted growth experiments, and the *E. coli* chemostat experiment. For the *E. coli* chemostat experiment, we computed the Pearson's correlation between $\log_2(\text{PTR})$ at each time t from each method and $\frac{1}{\tau(t)}$. For the *E. coli* unrestricted growth experiments we computed the Pearson's correlation between $\log_2(\text{PTR})$ at each time point i with sample time t_i , and the finite-difference estimate of the change in population size computed using optical densities:

$$\frac{\log(OD(i)) - \log(OD(i-1))}{t_i - t_{i-1}} \quad (8)$$

Additional growth rate experiments using *E. faecalis* and *L. gasseri* datasets were performed following the supplement in Korem et al. (2015) (section "Calculation of temporal growth correlation with PTR," datasets correspond to Fig S2). Genome assembly for *L. gasseri* was performed in PATRIC (Davis et al. 2020) with the SPAdes assembler (Prijbelski et al. 2020). We could not generate a high quality assembly for *E. faecalis*, and instead used an assembly available in PATRIC (1351.4268).

19 Benchmark on marine microbial data

Raw reads were downloaded from the SRA, project ID PRJNA551656, and MAGs from <https://doi.org/10.6084/m9.figshare.9730628>. CoPTR was run with default parameters. Values for GRiD, iRep, and DEMIC were obtained from Table S2 of Long et al. (2021). Observed growth rates were calculated as the slope of log-transformed MAG abundances, also obtained from Table S1 of

Long et al. (2021). As the sampling frequency was low, and as PTR is meant to provide an instantaneous estimate of growth, we used two consecutive data points to estimate growth (Equation 8), and compared them to the PTR at the first of the two, and discarded negative growth rates. We used FastANI (Jain et al., 2018) to ascertain that all MAG pairs had <95% average nucleotide identity (ANI): two pairs, MAGs 6 and 19, and MAGs 8 and 22, had ANI >99%; we therefore discarded MAGs 19 and 22, which had lower completion. Following Long et al., we removed MAGs which were not growing, which we defined as those with maximal observed growth rate of <2/day. Finally, we included only MAGs which had more than 3 pairs of observed growth rates and PTRs.

Antibiotic resistance experiment

We applied CoPTR to a dataset of 86 longitudinal samples from three populations of *C. rodentium*. Samples were taken from three periods of the experiment: a treatment period where the antibiotic was applied, a recovery period when the antibiotic was removed, and a stationary period. The structure of the experiment requires a model that accounts for the sampling time under each period. Let $\mathcal{P} = \{\text{Treatment, Recovery, Stationary}\}$ and for each $p \in \mathcal{P}$ denote T_p as the number of time points. We fit the following model:

$$\log_2(\text{PTR}) = a_{Ery}1_{Ery} + a_{Nal}1_{Nal} = \sum_{p \in \mathcal{P}} \sum_{t=0}^{T_p-1} 1_p(b_p - a_p t) + \epsilon \quad (9)$$

The parameters b_p allow a different mean $\log_2(\text{PTR})$ under each time period; the a_p model directional changes within each period over time. The variables a_{Ery} and a_{Nal} measure the effect of each antibiotic on the $\log_2(\text{PTR})$. While the model is somewhat complex, it is a reflection of the sampling process and dynamics of *in vitro* populations in culture.

1

2 IBD dataset experiments

3 We downloaded 1317 metagenomic samples from a case-control study of 106 individuals with
 4 inflammatory bowel disease (Lloyd-Price et al., 2019) from NCBI (BioProject PRJNA398089). The
 5 samples included all publicly available metagenomic samples from the study excluding technical replicates.
 6 Additional metadata from the study was downloaded from the Integrative Human Microbiome Project data
 7 portal (<https://hmpdacc.org/ihmp/>) to group metagenomic samples by individual. Samples were
 8 mapped to the IGGdb (Nayfach et al., 2019) of representative genomes for human gut species with a
 9 high-quality genome (N=2,935) downloaded from <https://github.com/snayfach/IGGdb>. The
 10 database was indexed, and reads were mapped, using CoPTR's (version 1.1.0) wrapper around Bowtie 2
 11 (version 2.4.1). Sample quality was assessed per genome per sample using CoPTR's default parameters (see
 12 CoPTR Implementation), and estimates produced for each genome that passed CoPTR's quality thresholds.
 13 The resulting PTR table included estimates from 1304 individuals and 660 species: 13 samples had no PTR
 14 estimates.

15

16 **Computing the fraction of variance explained.** Let r_{ij} be the j -th PTR of a species observed in
 17 categorical variable i (i.e. an individual, age group, sex, or disease status). To compute the fraction of
 18 variance explained we fit the random effects model

$$\log_2 r_{ij} = \mu + U_i + \epsilon_{ij} \quad (10)$$

19

$$U_i \sim N(0, \sigma_u^2) \quad (11)$$

20

$$\epsilon_{ij} \sim N(0, \sigma_e^2) \quad (12)$$

21

using the Statsmodels package (Seabold and Perktold, 2010) and reported $\frac{\sigma_u^2}{\sigma_u^2 + \sigma_e^2}$ per species. Because individuals accounted for a large fraction of variation, we selected one PTR at random from each individual to estimate variance components for disease status, age, and sex. For age, we divided individuals into a younger and older group using 18 years as a cutoff resulting in two categories. PTR estimates are sparse across species (Supplemental Figure S11). Therefore, when computing individual variation we only included species that had at least 10 PTR estimates in at least 3 individuals; for all other categories we only included species that had at least 10 PTR estimates in each category.

Correlation with relative abundances. We computed the correlation between $\log_2(\text{PTR})$ and relative abundances from MetaPhlAn2 (Truong et al., 2015). We matched species names from IGGdb to species names in MetaPhlAn2. For each species with more than 25 estimated PTRs, we computed standardized $\log_2(\text{PTR})$ and standardized $\log_2(\text{Rel Abun})$ by subtracting the mean and dividing by the standard deviation, then concatenated the resulting estimates from all species together into a single vector. The Pearson's correlation was computed between the two concatenated vectors.

Associating PTRs with disease status. Because individuals have multiple samples, PTR estimates from the same individual are not independent. Therefore, we tested for a difference in means between cases and controls by taking the mean per individual and adjusting by sample size. We chose this strategy over a linear mixed model because it has higher statistical power. Let r_{ij} be the j -th estimate of a PTR in a species for individual i , let n_i be the total number of PTRs in individual i for that species, and $\bar{r}_i = \frac{1}{n_i} \sum_{j=1}^{n_i} r_{ij}$. We fit the model

$$\sqrt{n_i} \log_2 \bar{r}_i = \sqrt{n_i} + \epsilon_{ij}$$

$$\mu = \text{intercept} + \beta(1_{is\ a\ case})$$

We computed p -values for β separately for each species and disease status, and adjusted for false discoveries using the Benjamini-Hochberg procedure (Benjamini and Hochberg, 1995). We limited our investigation to species with at least 10 PTR estimates in both cases and controls.

Associating PTRs with relative abundances and metabolomics. Because relative abundances and metabolite quantities change per sample, we could not use the same association procedure. We therefore fit the linear mixed model

$$\log r_{ij} = \mu + U_i + \beta x_k + \epsilon_{ij}$$

where μ is a fixed mean, U_i is a random effect for each individual, and x_k is the measurement of interest (a relative abundance or metabolite quantity). For metabolites, we used a log transformation with pseudo count 1 for zeros following the original study (Lloyd-Price et al., 2019). For metabolites, we limited our associations to named metabolites in the Human Metabolome Database. p -values were adjusted for false-discoveries using the Benjamini-Hochberg procedure (Benjamini and Hochberg, 1995).

Software availability

CoPTR is available under a GPL-3.0 License from <https://github.com/tyjo/coptr>, and under Supplementary Code. Documentation for CoPTR is on <https://coptr.readthedocs.io/>.

Acknowledgements

This material is based upon work supported by the National Science Foundation Graduate Research Fellowship to T.A.J. under Grant No. DGE-1644869. T.K. is a CIFAR Azrieli Global Scholar in the Humans & the Microbiome Program. Additional support was provided by NIH/NCI Grant No. U54CA209997 Driving Biological Projects and Columbia University's 2020/2021 Data Science Institute Seed Grant.

1

2 **Competing interests**

3 The authors declare no competing interests.

4

References

- Almeida A, Mitchell AL, Boland M, Forster SC, Gloor GB, Tarkowska A, Lawley TD, and Finn RD. 2019. A new genomic blueprint of the human gut microbiota. *Nature* **568**: 499–504.
- Almeida A, Nayfach S, Boland M, Strozzi F, Beracochea M, Shi ZJ, Pollard KS, Sakharova E, Parks DH, Hugenholtz P, et al.. 2021. A unified catalog of 204,938 reference genomes from the human gut microbiome. *Nature biotechnology* **39**: 105–114.
- Benjamini Y and Hochberg Y. 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)* **57**: 289–300.
- Bremer H and Churchward G. 1977. An examination of the cooper-helmstetter theory of dna replication in bacteria and its underlying assumptions. *Journal of theoretical biology* **69**: 645–654.
- Brown CT, Olm MR, Thomas BC, and Banfield JF. 2016. Measurement of bacterial replication rates in microbial communities. *Nature biotechnology* **34**: 1256.
- Bucci V, Tzen B, Li N, Simmons M, Tanoue T, Bogart E, Deng L, Yeliseyev V, Delaney ML, Liu Q, et al.. 2016. Mdsine: Microbial dynamical systems inference engine for microbiome time-series analyses. *Genome biology* **17**: 121.
- Buffie CG, Bucci V, Stein RR, McKenney PT, Ling L, Gobourne A, No D, Liu H, Kinnebrew M, Viale A, et al.. 2015. Precision microbiome reconstitution restores bile acid mediated resistance to *clostridium difficile*. *Nature* **517**: 205.
- Davis JJ, Wattam AR, Aziz RK, Brettin T, Butler R, Butler RM, Chlenski P, Conrad N, Dickerman A, Dietrich E, et al.. 2020. The PATRIC Bioinformatics Resource Center: expanding data and analysis capabilities. *Nucleic acids research*, 48(D1), D606-D612.
- DiGiulio DB, Callahan BJ, McMurdie PJ, Costello EK, Lyell DJ, Robaczewska A, Sun CL, Goltsman DS, Wong RJ, Shaw G, et al.. 2015. Temporal and spatial variation of the human microbiota during pregnancy. *Proceedings of the National Academy of Sciences* **112**: 11060–11065.
- Emiola A and Oh J. 2018. High throughput in situ metagenomic measurement of bacterial replication at ultra-low sequencing coverage. *Nature communications* **9**: 1–8.
- Emiola A, Zhou W, and Oh J. 2020. Metagenomic growth rate inferences of strains in situ. *Science Advances* **6**: eaaz2299.
- Forster SC, Kumar N, Anonye BO, Almeida A, Viciani E, Stares MD, Dunn M, Mkandawire TT, Zhu A, Shao Y, et al.. 2019. A human gut bacterial genome and culture collection for improved metagenomic analyses. *Nature biotechnology* **37**: 186–192.
- Gao Y and Li H. 2018. Quantifying and comparing bacterial growth dynamics in multiple metagenomic samples. *Nature methods* **15**: 1041–1044.
- Gibbons S, Kearney S, Smillie C, and Alm E. 2017. Two dynamic regimes in the human gut microbiome. *PLoS computational biology* **13**: e1005364.

- 1 Gibson TE and Gerber GK. 2018. Robust and scalable models of microbiome dynamics. *arXiv preprint arXiv:1805.04591* .
- 2 Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, and Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic
- 3 genomes reveals clear species boundaries. *Nature Communications* **9**: 5114.
- 4 Joseph TA, Shenhav L, Xavier JB, Halperin E, and Pe'er I. 2020. Compositional Lotka-Volterra describes microbial dynamics in
- 5 the simplex. *PLOS Computational Biology* **16**: e1007917.
- 6 Korem T, Zeevi D, Suez J, Weinberger A, Avnit-Sagi T, Pompan-Lotan M, Matot E, Jona G, Harmelin A, Cohen N, et al.. 2015.
- 7 Growth dynamics of gut microbiota in health and disease inferred from single metagenomic samples. *Science* **349**:
- 8 1101–1106.
- 9 Langmead B and Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nature methods* **9**: 357.
- 10 Lloyd-Price J, Arze C, Ananthakrishnan AN, Schirmer M, Avila-Pacheco J, Poon TW, Andrews E, Ajami NJ, Bonham KS,
- 11 Brislawn CJ, et al.. 2019. Multi-omics of the gut microbial ecosystem in inflammatory bowel diseases. *Nature* **569**:
- 12 655–662.
- 13 Lloyd-Price J, Mahurkar A, Rahnavard G, Crabtree J, Orvis J, Hall AB, Brady A, Creasy HH, McCracken C, Giglio MG, et al..
- 14 2017. Strains, functions and dynamics in the expanded human microbiome project. *Nature* **550**: 61.
- 15 Long AM, Hou S, Ignacio-Espinoza JC, and Fuhrman JA. 2021. Benchmarking microbial growth rate predictions from
- 16 metagenomes. *The ISME Journal* **15**: 183–195.
- 17 Luo H and Gao F. 2019. Doric 10.0: an updated database of replication origins in prokaryotic genomes including chromosomes and
- 18 plasmids. *Nucleic acids research* **47**: D74–D77.
- 19 Ma R, Cai TT, and Li H. 2020. Optimal estimation of bacterial growth rates based on a permuted monotone matrix. *Biometrika* .
- 20 Nayfach S, Shi ZJ, Seshadri R, Pollard KS, and Kyrpides NC. 2019. New insights from uncultivated genomes of the global
- 21 human gut microbiome. *Nature* **568**: 505–510.
- 22 Olm MR, Crits-Christoph A, Diamond S, Lavy A, Carnevali PBM, and Banfield JF. 2020. Consistent metagenome-derived metrics
- 23 verify and delineate bacterial species boundaries. *Msystems* **5**.
- 24 Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, and Phillippy AM. 2016. Mash: fast genome and
- 25 metagenome distance estimation using minhash. *Genome biology* **17**: 132.
- 26 Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, Beghini F, Manghi P, Tett A, Ghensi P, et al.. 2019. Extensive
- 27 unexplored human microbiome diversity revealed by over 150,000 genomes from metagenomes spanning age, geography,
- 28 and lifestyle. *Cell* **176**: 649–662.
- 29 Prjibelski A, Antipov D, Meleshko D, Lapidus A, and Korobeynikov, A. 2020. Using SPAdes de novo assembler. *Current protocols*
- 30 *in bioinformatics*, 70(1), e102.
- 31 Seabold S and Perktold J. 2010. Statsmodels: Econometric and statistical modeling with python. In *Proceedings of the 9th Python in*
- 32 *Science Conference*, volume 57, p. 61. Austin, TX.

- 1 Serrano MG, Parikh HI, Brooks JP, Edwards DJ, Arodz TJ, Edupuganti L, Huang B, Girerd PH, Bokhari YA, Bradley SP, et al.
2 2019. Racioethnic diversity in the dynamics of the vaginal microbiome during pregnancy. *Nature medicine* **25**: 1001–1011.
- 3 Shenhav L, Furman O, Briscoe L, Thompson M, Silverman JD, Mizrahi I, and Halperin E. 2019. Modeling the temporal dynamics
4 of the gut microbial community in adults and infants. *PLOS Computational Biology* **15**: e1006960.
- 5 Stein RR, Bucci V, Toussaint NC, Buffie CG, Räscher G, Pamer EG, Sander C, and Xavier JB. 2013. Ecological modeling from
6 time-series inference: insight into dynamics and stability of intestinal microbiota. *PLoS computational biology* **9**: e1003388.
- 7 Suzuki S and Yamada T. 2020. Probabilistic model based on circular statistics for quantifying coverage depth dynamics originating
8 from dna replication. *PeerJ* **8**: e8722.
- 9 Truong DT, Franzosa EA, Tickle TL, Scholz M, Weingart G, Pasolli E, Tett A, Huttenhower C, and Segata N. 2015. Metaphlan2 for
10 enhanced metagenomic taxonomic profiling. *Nature methods* **12**: 902–903.
- 11 Vila AV, Imhann F, Collij V, Jankipersadsing SA, Gurry T, Mujagic Z, Kurilshikov A, Bonder MJ, Jiang X, Tigchelaar EF, et al..
12 2018. Gut microbiota composition and functional changes in inflammatory bowel disease and irritable bowel syndrome.
13 *Science translational medicine* **10**.
- 14 Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser W, Bright J,
15 et al.. 2020. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. *Nature Methods* **17**: 261–272.
- 16 Zhou W, Sailani MR, Contrepois K, Zhou Y, Ahadi S, Leopold SR, Zhang MJ, Rao V, Avina M, Mishra T, et al.. 2019.
17 Longitudinal multi-omics of host–microbe dynamics in prediabetes. *Nature* **569**: 663–671.
- 18 Zou Y, Xue W, Luo G, Deng Z, Qin P, Guo R, Sun H, Xia Y, Liang S, Dai Y, et al.. 2019. 1,520 reference genomes from cultivated
19 human gut bacteria enable functional microbiome analyses. *Nature biotechnology* **37**: 179–185.
- 20
21
22
23
24
25
26

Figure Legends

Figure 1: CoPTR Workflow. Sequencing reads from multiple metagenomic samples are mapped to a reference database containing representative strains from complete reference genomes and high-quality assemblies (>90% completeness, <5% contamination). Multi-mapped reads are reassigned to a single genome using a probabilistic model. After read mapping, regions of each genome with ultra-high or ultra-low coverage are filtered using filters designed for complete reference genomes or draft assemblies. Then, PTRs are computed for each genome in each sample. For species with complete reference genomes, PTRs are estimated by maximizing the likelihood of a model describing the density of reads along the genome (CoPTR-Ref). For species with high-quality assemblies, reads are binned across the assembly, bins are reordered based on sequencing coverage across multiple samples using Poisson PCA, and the slope along this order is estimated by maximum likelihood (CoPTR-Contig). CoPTR outputs a table of the $\log_2(\text{PTR})$ per genome in each sample.

Figure 2: CoPTR-Ref is accurate on simulated and real data. (A) Accuracy of CoPTR-Ref and KoremPTR on simulated data based on an *E. coli* genome. Performance was compared by computing the root-mean-square-error (RMSE) of the $\log_2(\text{PTR})$ (y-axis) across 100 replicates while varying the number of reads (x-axis), varying the position of the replication origin, and varying the PTR. **(B)** Ground truth (x-axis) and estimated (y-axis) $\log_2(\text{PTR})$ across 100 simulation replicates with 20000 reads. KoremPTR appears to underestimate the true $\log_2(\text{PTR})$. **(C)** Comparison of KoremPTR $\log_2(\text{PTR})$ (x-axis) and CoPTR $\log_2(\text{PTR})$ (y-axis) on 6 real genomic and metagenomic datasets. **(D)** Evaluation of CoPTR-Ref's $\log_2(\text{PTR})$ estimates using representative genomes from different strains (5 *E. coli* strains, 4 *L. gasseri* strains, and 5 *E. faecalis* strains). Each dataset in panel C was mapped to strains from the same species, and the

Pearson's correlation (y-axis) was computed for each pair of strains. When the distance between strains (x-axis) is small, $\log_2(\text{PTR})$'s are highly correlated.

Figure 3: CoPTR-Contig is accurate on simulated and real data. (A) Comparison of CoPTR-Ref and CoPTR-Contig on simulated data using the *E. coli* density map. Performance was evaluated by computing the correlation (y-axis) between simulated and estimate $\log_2(\text{PTR})$'s across read counts (x-axis), randomly chosen replication origins, and PTRs. CoPTR-Contig shows high accuracy above 5000 reads. **(B)** Comparison of CoPTR-Contig to GRiD, DEMIC, and iRep across 5 genomic (monoculture) and metagenomic datasets (x-axis). For each dataset, reads were mapped to a single reference genome for each species (see Methods). Performance was evaluated by comparing $\log_2(\text{PTR})$ estimates from CoPTR-Ref to the $\log_2(\text{PTR})$ estimate from each method across 10 high-quality metagenome assembled genomes (MAGs; points on the figure). Significance was computed using a two-tailed *t*-test (*: $p < 0.05$; **: $p < 10^{-2}$; ***: $p < 10^{-3}$; ****: $p < 10^{-4}$). **(C)** Number of PTR estimates from species passing the filtering criteria for each model. The mean and standard deviation are reported for the *E. coli* metagenomic gut and genomic datasets across MAGs from (B). Error bars depict one standard deviation. Each model was also applied to 10 samples from the IBD dataset using 1,009 high-quality MAGs from the IGGdb. The total number of PTRs passing filtering criteria for each model is reported.

Figure 4: CoPTR values strongly correlate with observed growth rates. (A, B) $\log_2(\text{PTR})$ estimates (y-axis) versus observed growth rates (x-axis; Methods), for two well predicted MAGs: MAG 6, classified as Flavobacteriales **(A)**, and MAG 13, classified as SAR86 **(B)**. Lines are linear regressions. **(C)** Comparison of per-MAG Pearson r (y-axis) for all MAGs with >3 pairs of measured growth rates and calculated PTRs. CoPTR values are compared to GRiD, DEMIC, and iRep. Significance was computed using a two-sided Mann Whitney U test.

Figure 5: PTRs are highly personalized and uncorrelated with relative abundances. (A) Fraction of variance of $\log_2(\text{PTR})$ explained per species by variation between individuals, disease-statuses, age, and sex. Inter-individual variation accounts for most variation in among $\log_2(\text{PTR})$ s of the variables explored. **(B)** Boxplots of the $\log_2(\text{PTR})$ (x-axis) of *Parabacteroides distasonis* across individuals (y-axis). *P. distasonis* had the smallest p -value when testing for individual differences using the Kruskal-Wallis test on controls. PTRs appear stable within individuals. **(C)** Correlation between standardized $\log_2(\text{PTR})$ and $\log_2(\text{relative abundance})$ on species matched to relative abundances estimated MetaPhlAn2.


Figure 6: Association of $\log_2(\text{PTR})$ s with disease status (A), relative abundances (B), and metabolomics (C). **(A)** $\log_2(\text{PTR})$ s can be used to associate species with disease status. Significance was assessed by a fitting a linear model to $\log_2(\text{PTR})$ per species and correcting for false-discoveries (q -values denote false-discovery rate). PTRs can be combined with relative abundances to assess species interactions **(B)** or the impact of metabolites **(C)**.

Metagenomic Samples


```
@SEQ_ID1
GCTAAGTAAACGTTACCT...
+
00007<70'0'<<'<BFF...
@SEQ_ID2
...
```

Map
reads

Reference
Database




Re-assign multi-mapped reads



Complete Reference Genome


Draft Assembly

CoPTR-Ref




$\log_2(\text{PTR})$

Samples



CoPTR-Contig

1. Compute coverage across
multiple samples




2. Factor using Poisson PCA

$$\approx \exp \left\{ \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_B \end{bmatrix} \begin{bmatrix} v_1 & v_2 & \dots & v_N \end{bmatrix} \right\}$$


Bin Scores





Individual
Scores




3. Reorder bins by scores









4. Fit model













A**B****C**