

Supplementary Table II. Primers used to detect genomic DNA deletion in *ura⁻* mutants

	Forward	Reverse
U3	GCTCCATTGCATTGTCTGTT	TTGACTATGCTAAGCTGCCT
U6	GCAGTTTTGAGTGATAATAT	GATATAGTGTGTTGTTCTAG
U4	TTCAGCAGAAGACTTGCAAC	CATCATCGACTTGCAAGAAC
U1	ATCGAGTCTAAGTCATGCTT	GGTTCTATAACTGGATATTG
U2	TCTCATCTTGTCTAAATGCA	CCTAATTCTCTATTTGTCTC
D1	TAGCTTCAGTGAATTGCTTG	CATACAGTTTTATGATGTCT
D2	AGCACCATGTAGATCAACGA	CGTAATTTTCATAGCTATGAC
D3	AAGTCAATATACCTGACTGT	TGATTCAAGTCTTGTAATCC
D4	AGGTTTATGTGAACGCAATC	CATGCTAACTGAAGTTCCAA
D5	TAAATGATTCACCGACCATG	ATCTCACACCGCATAAACGA
D6	GGCGATATTCAATTGGTTTG	CTCAACACAATGGTAAAAGA
D7	TATCGCAACCTTGCAAGCCT	GACCTCAAACGAATGTATGT
D8	TGAAAGTTCTGGATGTGTTT	TCTAACCTGATCAACGTAAG

Genomic Localization of PCR fragments is shown in Fig. 6D.